

Effects of Water Stress on Germination in Six Provenances of *Pinus brutia* Seeds from Different Bioclimatic Zones in Turkey

Melih BOYDAK, Hüseyin DIRİK, Fahrettin TİLKİ, Mehmet ÇALIKOĞLU
Istanbul University, Faculty of Forestry, Department of Silviculture, 80895, Bahçeköy, İstanbul - TURKEY

Received: 12.07.2002

Abstract: The effects of water potential on germination were studied in six provenances of *Pinus brutia* from different bioclimatic zones in Turkey. Water potentials between 0 and -8 bars were obtained using polyethylene glycol 6000 (PEG-6000) solutions. Seeds were kept for 28 d in light (12 h photoperiod at 1000 lux) and at 20 ± 0.5 °C. A decrease in water potential produced a marked reduction in germination percentage, germination speed and germination value. Significant variations between the provenances were found. The provenances of humid regions were the most susceptible to high moisture stress, and their germination value and germination percentage showed significant reductions at -6 bars. These differences in germination ability of provenances might be attributed to intraspecific variations resulting from the effects of natural selection.

Key Words: *Pinus brutia*, germination, polyethylene glycol, water potential

Türkiye'nin Farklı Biyoklimatik Zonlarından Altı Kızılçam (*Pinus brutia*) Orijinine Ait Tohumların Çimlenmesi Üzerine Su Stresinin Etkileri

Özet: Bu çalışmada, Türkiye'nin farklı biyo-iklim bölgelerine ait altı *Pinus brutia* orijininin çimlenmesi üzerine ozmotik stresin etkisi belirlenmeye çalışılmıştır. PEG-6000 eriyiği kullanılarak 0 ile -8 bar arasında stres düzeyleri oluşturulmuştur. Tohumların çimlendirilmesi 20 ± 0.5 °C sıcaklıkta 12 saat ışık (1000 lüks) altında 28 gün esas alınarak yapılmıştır. Ozmotik potansiyelin azalması ile çimlenme yüzdesi, çimlenme hızı ve çimlenme değeri bütün orjinlerde önemli oranda azalmakla birlikte orjinlerin tutumu arasında önemli farklılıklar görülmektedir. Yağışlı bölgelerden elde edilen orjinler yüksek strese karşı en fazla duyarlılığı göstermekte, az yağışlı bölgeleri temsil eden orjinlere ait tohumlar ise özellikle yüksek stres derecesinde (-6 bar) daha fazla oranda çimlenebilmektedirler. Orijinlerin ozmotik strese karşı gösterdikleri bu farklılığın bir doğal seleksiyon etmeni olarak kuraklığın etkisiyle oluşmuş bir intraspecific varyasyondan ortaya çıktığı ifade edilebilir.

Anahtar Sözcükler: Kızılçam, Çimlenme, Polietilen glikol, su potansiyeli

Introduction

Pinus brutia Tenore, Calabrian pine, is found largely in the Eastern Mediterranean region, and primarily in Turkey (85%) (Nahal, 1984). The natural distribution of *P. brutia* in Turkey indicates that it is well adapted to the Mediterranean-type climate; it is found in the thermomediterranean, mesomediterranean and submediterranean zones (Quezel, 1979).

P. brutia is one of the most commercially important forest species in Turkey, where it is dominant on about 3 million ha (Usta, 1990). It is a medium-lived, intolerant to moderately tolerant tree with rapid juvenile growth. It is fast growing and thus has a short rotation. This species can be managed as even-aged natural stands, or can be

regenerated artificially and managed in plantations. Clearcutting with direct seeding is used on most sites in Turkey including natural fire affected areas (Boydak, 1993). Thus, the seed germination and survival of different provenances are of great importance for natural or artificial regeneration. In addition, the intraspecific variations of drought tolerance can be remarkable.

The climate over most of the *P. brutia* range is semi-humid with hot summers and mild winters. Average annual rainfall varies from 500 mm to 1500 mm (Atalay, 1993; Atalay et al. 1998).

A large number of studies have been carried out on the effects of water stress on the germination of forest tree species including *P. brutia*, and most species and

provenances have shown a different sensitivity to water stress as regards germination and subsequent root growth (Bonner and Farmer, 1966; Barnett, 1969; Djavanshir and Reid, 1975; Kaufmann and Eckard, 1977; Calamassi et al., 1980; Falusi and Calamassi, 1982; Falusi et al., 1983; Dunalp and Barnett, 1984; Thanos and Skordillis, 1987; Falleri, 1994; Lopez et al., 2000). However, in none of the studies was the selection of provenances at the beginning of the research based on a bioclimatic classification.

Soil water supply is an important environmental factor controlling seed germination (Kramer and Kozłowski, 1979). If the water potential is reduced, seed germination will be delayed or prevented depending on the extent of its reduction (Hegarty, 1978). One technique for studying the effect of water stress on germination is to simulate stress conditions using artificial solutions to provide variable water potentials (Larson and Shubert, 1969; Sharma, 1973; Falusi et al., 1983). In the present work, the effects of water stress were examined in six provenances of *P. brutia* using PEG solutions with water potentials ranging from 0 to -8 bars. The purpose of this study was to evaluate the influence of water stress on germination and to determine whether there was a significant intraspecific

variation in drought tolerance between provenances of *P. brutia* seeds from different bioclimatic zones.

Materials and Methods

Seeds were collected in 1998 from six natural stands of *P. brutia* by the Forest Trees and Seeds Improvement Division, Ministry of Forestry, Turkey. The seeds were extracted, cleaned and stored in a dark and cool place at 4 °C until used in 2000.

The climatic conditions prevailing in each of the six natural stands were considered when selecting locations in Turkey (Figure 1 and Table 1). Thus, the Emberger bioclimatic classification method developed for Turkey by Akman and Daget (1971) was used (Figure 2).

Before the germination tests, damaged and insect-infected seeds were discarded, and the empty ones were eliminated using the floating method in distilled water. The water potential of the germination substrates (0, -2, -4, -6 and -8 bars) was determined using PEG-6000 solution, prepared as described by Michel and Kaufman (1973).

Germination tests were performed in 11-cm-diameter glass petri dishes on two layers of filter paper saturated




Figure 1. Location of *P. brutia* provenances studied: T1 (Bucak), T2 (Mustafa Kemalpaşa), T3 (Urla), T4 (Yaras), T5 (Duzlercami) and T6 (Kargi) (adapted from Saatçioğlu, 1976).

Table 1. List of *P. brutia* provenances.

Provenances	Latitude (North)	Longitude	Altitude (m)	Mean precip. (mm)	Mean temp. (°C)	Precip.-temp. index (Q)	Bioclimatic zone
T1 Bucak	37° 24'	30° 37' E	800	712	14.1	74.7	Low rain-cold
T2 M. Kemalpaşa	39° 58'	28° 40' E	250	702	14.6	86.3	Low rain-cool
T3 Urla	38° 14'	26° 36' E	150	695	17.6	87.5	Low rain-warm
T4 Yaras	37° 06'	28° 32' E	750	1209	15.0	133.5	High rain-cool
T5 Duzlercami	36° 59'	30° 33' E	275	1089	18.7	135.7	High rain-warm
T6 Kargı	36° 36'	31° 57' E	350	1080	18.1	155.8	High rain-hot




Figure 2. Location of *P. brutia* provenances on Emberger's Bioclimatic Diagram. Dotted lines show the natural distribution of *P. brutia* in Turkey (adapted from Akman and Daged, 1971).

with water solutions. Four 50-seed replicates for each lot and for each experimental condition were used, for a total of 120 replicates (4 replicates [each has 50 seeds]* 5 water potential levels *6 provenances = 120). Filter papers and solutions were changed every 3 d in order to keep the water potential steady during the whole test period. Experiments were carried out in a temperature-controlled growth chamber at 20 ± 0.5 °C with a photoperiod of 12 h and illuminance of 1000 lux.

Germination counts were performed daily for 28 d and germination was considered to have occurred if the radicle protruded 2 mm from the seed coat. Seeds with abnormal radicles were excluded from the germination counts (Edwards and Wang, 1995). Cumulative

germination percentage, (GP%) was evaluated daily and the final value was obtained after 28 d. Then all cumulative germination percentages of the provenances at every stress level were transformed to relative cumulative germination percentages by considering the control germinations (0 bars) to be 100. These percentages were transformed to $\arcsin P^{1/2}$ for statistical analysis. The results were analysed by factorial analysis and the means were compared with Duncan's test. Mean germination time (MGT) and germination value (GV), the parameters of germination speed, were also calculated for supplementary explanations. MGT, day was computed as follows: $MGT = \sum n_i d_i / n$, where n is the total number of germinated seeds during the germination test, n_i is the

number of germinated seeds on day d_i and i is the number of days during the germination period (between 0 and 28 days) (Yousheng and Sziklai, 1985). In the present study, GV, day was calculated by the formula of Djavanshir and Pourbeik (1976) because it is believed to give a more reliable estimate of subsequent survival for the genus *Pinus*. Thus, GV was computed as follows: $GV = (\sum DGS/N) \times GP \times 10$, where DGS is daily germination speed, which is computed by dividing the cumulative germination percentage by the number of days since the beginning of the test. N is the number of DGS calculated during the test.

Results

Analysis of variance showed highly significant differences among both provenances and water potentials. The interaction between provenances and water stress was not significant (Table 2).

Although lowering the water potential to -8 bars reduced germination for all provenances by more than 50%, the reaction to increased water stress differed among the six provenances: T1 seems to be the most tolerant, with a threshold between -6 and -8 bars, and T5 the least tolerant, with a tolerance threshold between -4 and -6 bars. Considerable tolerance was also observed in T2, with a threshold between -4 and -6 bars (Table 3).

According to Duncan’s test, T1 and T2 were combined in an independent group as the most drought resistant provenances. T2 was also included in another group with T4 and T6, while T6 with T3 were in another different group. T5 was the least drought tolerant provenance. On the other hand, there were no significant differences between 0, -2 and -4 bar stress levels according to the germination percentages of different provenances, while -6 and -8 bar stress levels differed separately (Table 4).

Table 2. Results from the ANOVA on relative cumulative percentage germination.

Source of variation	df	MS	F	P
Provenances (A)	5	60.80	4.79	0.001
Water potentials (B)	4	1067.48	84.11	0.001
Replicate (C)	3	48.21	3.80	0.05
Interaction A x B	20	16.42	1.29	
Interaction A x C	15	13.87	1.09	
Interaction B x C	12	11.45	0.90	
Residual	60	12.69		
Total	119			

Table 3. Effects of water potential on cumulative germination percentage of *Pinus brutia* seeds from six provenances.

Provenances	Water potential (bars)					Average
	0	-2	-4	-6	-8	
T1	74.0	96.0	75.0	70.0	31.0	69.2
T2	91.0	90.5	92.0	66.5	40.5	76.1
T3	90.5	93.0	79.0	47.5	32.0	68.4
T4	62.5	66.5	66.5	41.5	12.0	49.8
T5	94.5	91.0	84.5	46.0	15.0	66.2
T6	93.5	95.5	86.5	61.5	20.5	71.5
Average	84.3	88.7	80.6	55.5	25.2	

Table 4. Effects of water potential on relative cumulative germination percentage of *Pinus brutia* seeds from six provenances.

Provenances	Water potential (bars)					Average
	0	-2	-4	-6	-8	
T1	100	129.7	101.3	94.6	41.9	93.5 a ¹
T2	100	99.5	101.1	73.1	44.5	83.6 ab
T4	100	106.4	106.4	66.4	19.2	80.0 b
T6	100	100.5	95.1	67.6	22.5	77.0 bc
T3	100	102.8	87.3	52.5	34.8	75.5 bc
T5	100	90.5	86.8	48.4	15.8	68.3 c
Average ²	100 a	104.9 a	96.3 a	67.1 b	29.8 c	

¹ Values in the column with the same initial(s) are not significantly different (P < 0.001).

² Values in the row with the same capital initial are not significantly different (P < 0.001).

Relative cumulative percentage germination was significantly reduced between -4 and -6 bars for all provenances (Table 4). Germination of three provenances (T4, T5 and T6) from humid regions decreased from 100% to around 20%, while the other provenances (T1, T2 and T3) continued to germinate more than 34% at the lowest water potential (-8 bars).

MGT differed in the six provenances and was markedly influenced by water potential (Table 5). Even a stress of -2 bars caused a significant increase in MGT (decrease in germination rate) in five provenances (T2, T3, T4, T5 and T6), but in T1 the tolerance threshold was between -4 and -6 bars. Provenance T4 was the most susceptible to high moisture stress since its germination rate was the lowest (highest MGT values), - 8 bars.

Table 5. Effects of water potential on mean germination time (MGT, days) of seeds from six provenances of *Pinus brutia*. See Materials and Methods for calculation of MGT.

Provenances	Water potential (bars)					Average
	0	-2	-4	-6	-8	
T1	16.58	15.45	16.58	21.88	23.23	18.74
T2	10.18	11.66	14.64	18.94	21.86	15.46
T3	9.92	11.67	14.69	20.04	23.61	15.98
T4	12.81	15.74	17.60	21.09	30.25	19.50
T5	11.78	16.24	17.21	21.70	23.86	18.16
T6	12.75	14.88	17.34	22.46	24.12	18.31
Average	12.74	14.27	16.34	21.02	24.49	

GVs of all provenances were affected at -4 bars (Table 6). Each further increment in water stress produced additional significant decreases in GV. The GV for T1, T2 and T3 were higher than those for the other provenances at the lowest water potential (-8 bars), and the lowest tolerance again occurred in T4, T5 and T6.

Discussion

The seeds of *P. brutia* from the six regions used in this work germinate under water stress without any prechilling treatment. Decreasing the water potential in the substrate decreased germination, indicating that water stress inhibits germination.

In this study, germination percentage was adversely affected when moisture stress reached -6 bars, while it was reduced at -8 bars by more than half. This result agrees well with the findings of Falusi and Calamassi (1982), who reported that a decrease in germination capacity in *P. brutia* seed occurred at -4 bars. Similar trends have also been observed in some other conifers; lowering the water potential to -8 bars reduced the germination of *P. pinaster* (Falleri, 1994), *Pinus contorta* and *Picea engelmannii* by approximately 50% (Kaufman and Eckard, 1977). Falusi et al. (1983) observed in *Pinus halepensis* that a reduction of the water potential of the germination substrate even to -2 bars lowers germination percentages considerably, while at -8 bars germination was lowered to approximately 25%. Djavanshir and Reid (1975) reported that a marked reduction of germination capacity occurred at -4 bars in *Pinus ponderosa* seeds. Similar trends were also reported for the seeds of *Pinus palustris* and *Pinus elliotti* (Barnett,

Table 6. Effects of water potential on germination value (GV, %) of seeds from six provenances of *Pinus brutia*. See Materials and Methods for calculation of GV.

Provenances	Water potential (bars)					Average
	0	-2	-4	-6	-8	
T1	21.99	30.20	16.32	9.06	1.86	15.89
T2	41.44	37.97	28.54	12.03	3.77	24.75
T3	40.37	38.92	23.87	5.06	1.84	22.01
T4	15.72	14.36	11.37	3.07	0.30	8.96
T5	37.71	24.81	18.17	3.71	0.40	16.96
T6	33.81	31.26	21.25	6.63	0.56	18.70
Average	31.84	29.59	19.92	6.59	1.45	

1969). In contrast, Thanos and Skordilis (1987) reported that *P. brutia* seeds exhibited water inhibition of germination at values lower than -10 bars, and the absolute values of the water potential required for 50% inhibition of germination in *P. halepensis* were between -14.6 and -19.5 bars at both 15 °C and 20 °C.

The results of this research support the idea that *P. brutia* is well adapted to the Mediterranean-type climate and is a drought-resistance species with respect to several physiological characteristics (Abido, 1983; Grunwald and Schiller, 1988; Dirik, 1994). The drought resistance of *P. brutia* has already been shown in the germination phase.

Drought is a major obstacle to the regeneration of *P. brutia* throughout its natural range. At low elevations, a portion of the seeds dispersed during the dry summer and early autumn germinate after autumn precipitation although the soil does not have enough moisture content. Autumn germination is important in natural regeneration and in broadcast seeding in the thermomediterranean zone (Boydak, 1993). These characteristics of *P. brutia* might be attributed to the fact that the cumulative germination percentage was not adversely affected when moisture stress reached -6 bars; lowering the water potential to -4 bars even increased the germination percentage in some provenances (Table 3). However, GV and MGT were not affected in the same way. In another study, water priming with aerated solutions of polyethylene glycol improved both final germination and the speed of germination in *P. brutia* (Dirik et al., 1999).

The response of germination to water stress differed among the six provenances. This intraspecific variation agrees with the experimental data reported previously for *P. brutia* (Calamassi et al., 1980; Falusi and Calamassi,

1982), *Pinus elderica* and *P. halepensis* (Calamassi et al., 1980; Falusi et al., 1983; Thanos and Skordilis, 1987), *P. taeda* (Dunalp and Barnett, 1984) and *Cedrus libani* (Dirik, 2000).

According to the results, T1 and T2 were the most stress-tolerant provenances on a water stress gradient. T4 and T6 followed these provenances. However, when we take into account the lowest water stress level (-8 bars), T1, T2 and T3, which have precipitation-temperature indexes (Q) below 100, have higher germination capacities than the other seed provenances. In addition, the relative germination capacities of T1 and T2 were highest at -6 and -8 bars. Climatic differences among the provenances might account for this behaviour; i.e. the higher amount of rain in the area of T4, T5 and T6 and the lower coefficient of precipitation-temperature index (Q) of T1, T2 and T3.

GVs decreased with decreasing water potential, an observation consistent with studies on *P. elderica* (Djavanshir and Reid, 1975), *P. halepensis* (Falusi et al., 1983), and *P. pinaster* (Falleri, 1994). As with germination percentages, the GV for T1, T2 and T3

were higher than those for the other provenances at the lowest moisture stress (-8 bars).

MGT differed among the six provenances, and the differences were markedly increased with decreasing water potential. Even moderate water stresses (-2 bars) caused an increase in MGT. Especially at lower values of water potential, the germination rate significantly decreased. Moderate water stress seems to have a stronger effect on germination capacity than higher water stress. These results are consistent with those of other studies on the genus *Pinus* (Falusi and Calamassi, 1982; Falusi et al., 1983; Falleri, 1994) and *Liquidambar* (Bonner and Farmer, 1966). However, at the lowest bars (-8 bars), there were no significant differences among the provenances in terms of germination rate.

The higher germination percentage and value in T1, T2 and T3 (Q < 100) at the lowest water potential (-8 bars) may be related to their better adaptation to water deficits compared to the other three provenances (Q > 100). Furthermore, differences under water stress also show that provenances are characterised by a significantly different tolerance to drought.

References

- Abido, M.S. 1983. Drought tolerance comparison of Aleppo pine and brutia pine seedlings. Thesis summary. Forestry Abstract 44.
- Akman, Y. and Ph. Daget. 1971. Quelques aspects synoptiques des climats de la Turquie. T 5, 3: 270-300.
- Atalay, İ. 1993. Ecological conditions and natural occurrence areas of *Pinus brutia* in Turkey, in: International Symposium on *Pinus brutia* Ten. 1993. Ministry of Forestry, Ankara, pp. 57-66.
- Atalay, İ., I. Sezer and H. Çukur. 1998. Kızıldağ (*Pinus brutia* Ten.) ormanlarının ekolojik özellikleri ve tohum nakli açısından bölgelere ayrılması (The ecological properties of Red pine (*Pinus brutia* Ten.) and their regioning in terms of seed transfer). Orman Ağaçları ve Tohum Islahı Araştırma Müdürlüğü 6. Ankara.
- Barnett, J.P. 1969. Moisture stress affects germination of longleaf and slash pine seeds. For. Sci. 15: 275-276.
- Bonner, F.T. and R.E. Farmer. 1966. Germination of sweetgum in response to temperature, moisture stress and length of stratification, For. Sci. 12: 40-43.
- Boydak, M. 1993. Kızıldağın silvikültürel özellikleri, uygulanabilecek gençleştirme yöntemleri ve uygulama esasları, in: Uluslararası Kızıldağ Bildirileri, Marmaris, Turkey, pp. 146-158.
- Calamassi, R., M. Falusi and A. Tocci. 1980. Variazione geografica e resistenza a stress idrico in semi di *Pinus halepensis* Mill. *Pinus brutia* Ten. and *Pinus elderica* Medw, Annali dell'Istituto Sperimentale per la Selvicoltura XI: 195-230.
- Dirik, H. 1994. Üç yerli çam türünün (*Pinus brutia* Ten., *Pinus nigra* Arn. ssp. *pallasiana* Lamb. Holmboe, *Pinus pinea* L.) kurak peryottaki transpirasyon tutumlarının ekofizyolojik analizi (Analyse écophysiological des comportements de transpiration de trois espèces forestiers (*Pinus brutia* Ten., *Pinus nigra* Arn. ssp. *pallasiana* Lamb. Holmboe, *Pinus pinea* L.)), İ.Ü. Orm. Fak. Derg. Seri A 44: 111-121.
- Dirik, H., M. Çalikoğlu and F. Tilki. 1999. Kızıldağ (*Pinus brutia* Ten.) tohumlarında ozmotik stres ile koşullandırmanın çimlenme üzerindeki etkileri (Effects of osmotic priming on germination of Calabrian pine (*Pinus brutia* Ten.) seeds), İ.Ü. Orm. Fak. Derg. Seri A, 49: 75-89.
- Dirik, H. 2000. Effet du stress hydrique osmotique sur la germination des graines chez les provenances de Cedre du Liban (*Cedrus libani* A. Rich.) d'origine turque, Annals of Forest Science 57: 361-367.
- Djavanshir, K. and C.C.P. Reid. 1975. Effect of moisture stress on germination and radicle development of *Pinus eldarica* Medw. and *Pinus ponderosa* Laws, Canadian Journal of Forest Research 5: 80-83.
- Djavanshir, K. and H. Pourbeik. 1976. Germination value – a new formula, Silvae Genetica 25: 79-83.
- Dunalp, J.R. and J.P. Barnett 1984. Manipulating loblolly pine (*Pinus taeda* L.) seed germination with simulated moisture and temperature stress. In: Seedling Physiology and Reforestation Success (Eds.: M.L. Duryea and G.N. Brown). Martinus Nijhoff/Dr. W. Junk. Publ., Boston, pp. 61-74.

- Edwards, D.G.V. and B.S.P. Wang. 1995. A Training guide for laboratory analysis of forest tree seeds. Pacific Forestry Center, Information Report BC X-356, Canada.
- Falleri, E. 1994. Effect of water stress on germination in six provenances of *Pinus pinaster* Ait, Seed Science and Technology 22: 591-599.
- Falusi, M. and R. Calamassi. 1982. Effetti degli stress idrici su germinazione e crescita della radice in cinque provenienze di *Pinus brutia* Ten. (Effects of moisture stress on germination and root growth in provenances of *Pinus brutia* Ten.), Ann. Acc. It. Sc. For. XXXI: 99-118.
- Falusi, M., R. Calamassi and A. Tocci. 1983. Sensitivity of seed germination and seedling root growth to moisture stress in four provenances of *Pinus halepensis* Mill., Silvae Genetica 32: 4-9.
- Grunwald, C. and G. Schiller. 1988. Needle xylem water potential and water saturation deficit in provenances of *Pinus halepensis* Mill. and *P. brutia* Ten., Forêt Méditerranéenne X: 407-414.
- Hegarty, T.W. 1978. The physiology of seed hydration and dehydration, and the relation between water stress and control of germination: a review, Plant Cell Environment 1: 101-109.
- Kaufmann, M.R. and A.N. Eckard. 1977. Water potential and temperature effects on germination of Engelmann spruce and Lodgepole pine seeds, Forest Science 23: 27-33.
- Kramer, P.J. and T.T. Kozlowski. 1979. Physiology of woody plants. Academic Press, New York, 811 p.
- Larson, M.M. and G.N. Shubert. 1969. Effect of osmotic water stress on germination and initial development of Ponderosa pine seedlings, Forest Science 15: 30-36.
- Lopez, M., J.M. Humara, A. Casares and J. Majada. 2000. The effect of temperature and water stress on laboratory germination of *Eucalyptus globulus* Labill. seeds of different sizes, Annals of Forest Science 57: 245-250.
- Michel, B.E. and M.R. Kaufmann 1973. The osmotic potential of polyethylene glycol 6000, Plant Physiology 51: 914-916.
- Nahal, I. 1984. Le pin brutia (*Pinus brutia* Ten. subs *P. brutia*). Deuxième partie, Forêt Méditerranéenne VI: 5-17.
- Quezel, P. 1979. La région méditerranéenne française et ses essences forestières, signification écologique dans le contexte circum-méditerranéen. Forêt Méditerranéenne I: 7-18.
- Saatçioğlu, F. 1976. Silvikültür I. İ.Ü Orm. Fak. Yayınları. No: 222/2187, İstanbul.
- Sharma, M.L. 1973. Simulation of drought and its effect on germination of five pasture species. Agronomy Journal 65: 982-987.
- Thanos, C.A. and A. Skordillis. 1987. The effects of light, temperature and osmotic stress on the germination of *Pinus halepensis* and *P. brutia* seeds, Seed Science and Technology 15: 163-174.
- Usta, H.F. 1990. Kızılgam (*Pinus brutia* Ten.) ağaçlandırmalarında hasılat araştırmaları. A study on the yield of *Pinus brutia* Ten. Plantations, OAE Yay. Teknik Bulten 219. 138 p.
- Yousheng, C. and O. Sziklai 1985. Preliminary study on the germination of *Toona sinensis* (A. Juss.) Roem. seed from eleven Chinese provenances. Forest Ecology and Management 10: 269-281.