

## Age-specific life tables of *Aonidiella aurantii* (Maskell) (Hemiptera: Diaspididae) and its parasitoid *Aphytis melinus* DeBach (Hymenoptera: Aphelinidae)

Khalid MOHAMMED<sup>1,2</sup> , İsmail KARACA<sup>3,\*</sup> , Manjree AGARWAL<sup>1</sup> , James NEWMAN<sup>1</sup> , YongLin REN<sup>1</sup> 

<sup>1</sup>School of Veterinary and Life Sciences, College of Science, Health, Engineering and Education, Murdoch University, Murdoch, Western Australia

<sup>2</sup>Department of Plant Protection, College of Agriculture and Forestry, University of Mosul, Mosul, Nineveh, Iraq

<sup>3</sup>Department of Plant Protection, Faculty of Agricultural, Isparta University of Applied Sciences, Isparta, Turkey

Received: 11.05.2019 • Accepted/Published Online: 30.10.2019 • Final Version: 01.04.2020

**Abstract:** Biological parameters of the California red scale (*Aonidiella aurantii* [Maskell] [Hemiptera: Diaspididae]) were determined under laboratory conditions at 3 different temperatures (20, 23, and 27 °C) on butternut squash (*Cucurbita moschata* Duchesne ex Lamarck) (Cucurbitaceae), while the biological parameters of its parasitoid *Aphytis melinus* DeBach (Hymenoptera: Aphelinidae) were conducted at 27 °C. The survival of *A. aurantii* ranged between 80.0% and 88.3%. The highest mortality was recorded during the adult stage, with mortalities ranging between 12% and 20%. On *C. moschata*, the total development time was  $93.1 \pm 9.73$ ,  $81.8 \pm 7.13$ , and  $65.7 \pm 6.37$  days, while the adult longevity was  $54.65 \pm 0.71$ ,  $47.05 \pm 0.97$ , and  $39.35 \pm 1.07$  days at 20, 23, and 27 °C, respectively. The oviposition period of *A. aurantii* was  $44.3 \pm 0.51$ ,  $40.65 \pm 0.41$ , and  $34.5 \pm 0.45$  days at 20, 23, and 27 °C, respectively. Average fecundity was  $73.25 \pm 1.827$ ,  $109.7 \pm 3.569$ , and  $129.35 \pm 4.564$  individuals at 20, 23, and 27 °C, respectively. For *A. melinus*, adult longevity was  $19.24 \pm 0.73$  days, and average fecundity  $62.7 \pm 2.81$  eggs at 27 °C. The preoviposition period was  $0.82 \pm 0.05$  days, oviposition period was  $15.7 \pm 0.52$  days, and postoviposition period was  $2.21 \pm 0.09$  days. The intrinsic rate of increase ( $r_m$ ) of *A. melinus* ( $0.188 \text{ ♀/♀/day}$ ) was significantly greater than that of *A. aurantii* (0.080) at 27 °C. These laboratory results demonstrated that *A. melinus* is an effective parasitoid for decreasing *A. aurantii* populations. Fecundity of *A. aurantii* and *A. melinus* was determined with the Enkegaard equation. The best-fit parameters of fecundity were calculated as  $a = 0.410$ ,  $b = 0.099$ ;  $a = 0.624$ ,  $b = 0.098$ ;  $a = 0.661$ ,  $b = 0.091$ ;  $a = 1.190$ ,  $b = 0.179$  for *A. aurantii* at 20, 23, and 27 °C, and *A. melinus* at 27 °C, respectively.

**Key words:** *Aphytis melinus*, biological control, California red scale, life table, population ecology

### 1. Introduction

Australia has an important role in citrus (*Citrus* spp., Rutaceae) production in the world, and this production has increased (FAO, 2017). Many pest species such as *Ceratitidis capitata* (Wiedemann) (Diptera: Tephritidae), *Aonidiella aurantii*, *Sasseita oleae* (Gomez-Menor Ortega) (Hemiptera: Coccidae), *Toxoptera citricada* (Kirkaldy) (Hemiptera: Aphididae), *Phyllocnistis citrella* Stainton (Lepidoptera: Gracillariidae), et al. were observed in WA (Western Australia) citrus orchards (Sonia, 2006). Growers have followed various methods to avoid damage caused by these pests. Biological control is one of these methods (FAO, 2017; Pekas, 2011a; Uygun and Satar, 2008). There are 32 species in the genus *Aonidiella* Berlese & Leonardi, which is a genus of scale insects in the family of Diaspididae (Hemiptera), the armoured scale insects (Ben-Dov, 2006). California red scale *Aonidiella aurantii* (Maskell) occurs on numerous host plants throughout the world. They attack different

crops such as fruit trees and ornamental plants all over the world, and cause heavy damage to the plants. Individual species infest leaves, fruits, branches, main stems, trunks, and roots. They are distributed throughout the world except in the cold extremes of the Arctic and Antarctic regions (Miller, 2005). California red scale *Aonidiella aurantii* is one of the most important pests infesting citrus trees in different parts of the world (Garcerá et al., 2008; Badary and Abd-Rabou, 2010; Hill, 2008; Karaca and Uygun, 1992; Uygun et al., 1995). California red scale feeds on various parts of host plants, such as twigs, leaves, and fruit (Beardsley and Gonzalez, 1975), harming them by inserting its mouthparts deep into plant tissue and sucking sap from parenchyma cells and injecting toxic saliva into the plants during the feeding process. Severe infestations of the scale cause leaf drop, defoliation, and dieback of twigs and branches. Young trees may die out in the absence of effective control (Hely et al., 1982; Smith et al., 1997).

\* Correspondence: [ikaraca98@gmail.com](mailto:ikaraca98@gmail.com)

The control of *A. aurantii* has encountered many difficulties, which has raised interest in alternative control methods (Vacas et al., 2010). *Aonidiella aurantii* has numerous predators and parasitoids (Hely et al., 1982; Forster et al., 1995). Among the parasitoids that feed on *A. aurantii*, *Aphytis melinus* DeBach (Hymenoptera: Aphelinidae) is known to be relatively important (Erler and Tunç, 2001). It was introduced into Australia in 1961 from the University of California, Riverside (Furness et al., 1983; Malipatil et al., 2000). It is now widely distributed in the citrus orchards of Victoria, South Australia, Western Australia, and the inland New South Wales citrus districts, and in Alice Springs in the Northern Territory (Smith et al., 1997). *Aphytis melinus* is commercially reared for release in citrus orchards to control *A. aurantii* (Grafton-Cardwell et al., 2006). Regarding parasitoids, females are more important to produce, because they are responsible for parasitising the host (via oviposition and host-feeding) and, thus, keep the pest's population under an economic threshold. A plethora of experiments have been conducted to determine the environmental conditions which best contribute to increases in the production of females (Ode and Hardy, 2008).

For the success of crop protection programs, it is essential to know some biological information about the pest. This information can be obtained by constructing a life table providing how different mortality factors act in sequence on the successive developmental stages. The intrinsic rate of increase ( $r_m$ ) is of value as the mean of describing the potential growth of population under given climatic and food conditions; it is an important parameter in inductive strategic and management models of pest populations (Abou Hatab, 1999; Bayoumy et al., 2009). Few studies have been conducted on the biological parameters of the California red scale and its parasitoid *A. melinus* (Heimpel, 1997; Badary and Abd-Rabou, 2010). Therefore, the aims of the present work were to simultaneously study the life history and behaviour of *A. aurantii* and its parasitoid *A. melinus* at 3 constant temperatures (20, 23, and 27 °C) under laboratory conditions.

## 2. Materials and methods

### 2.1. Rearing of *Aonidiella aurantii* and *Aphytis melinus*

Specimens of *A. aurantii* were initially collected in the field in 2016 from a population of *A. aurantii* in a citrus orchard located in WA (32.30°S, 116.01°E; 69 m AMSL), Australia. Stock cultures of *A. aurantii* were maintained on butternut squash (*Cucurbita moschata* Duchesne ex Lamarck), in climate chamber laboratory conditions [27 °C, 65 ± 5% RH, Light 16 h (L16) : Dark 8 h (D8) photoperiod] in the Murdoch University insect culture room.

The adult wasps of *A. melinus* used to start the colony were provided by Biological Services commercial

insectary (Adelaide, Australia). *Aphytis melinus* adults used in the experiments were reared following the method developed by Opp and Luck (1986) for rearing *A. melinus* in the climate chamber (26 °C, 40%–60% RH, L16 : D8 photoperiod). This method is commonly used to produce these insects in commercial insectaries. The production method is based on rearing the pest *A. aurantii* on butternut squash. When the host reached the third instar, which is the preferred age for the parasitoid to laid eggs and maximise progeny production, the infested squash with third instar scales was exposed to releases of 2-day-old adult parasitoids (male and female) in a ventilated cage (30 cm W × 30 cm H × cm D) with honey distributed on plastic trays. Adult parasitoids emerged 11–14 days later. Parasitoids thus produced participated in the experiments 1 day after emergence.

### 2.2. Survival, longevity, and fecundity of *Aonidiella aurantii*

Uniform groups of *A. aurantii* were obtained by leaving butternut squash undisturbed for 24 h in the *A. aurantii* colony maintained in the insect culture room at Murdoch University. These squashes were then removed from the colony, and each 10 randomly selected settled crawlers (those with the stylet inserted into the fruit and already forming the waxy cover) were enclosed by a micro plastic cage (small plastic thimble, 3 cm diameter with fine holes on its top) affixed with the help of modelling clay to the surface of the butternut squash. The squash fruit and nymphs were marked to track them throughout the experiment.

An age-specific life table of *A. aurantii* was constructed at 3 different constant temperatures (20, 23, and 27 °C). The infested fruits were kept in a ventilated polystyrene box (30 cm × 30 cm × 30 cm). Each treatment of 2 squash fruits (6 replicates) were kept on a plastic tray in identical climate cabinets (HWS Ningbu Southeast Equipment Co. Ltd., China) in which the experiment was conducted under controlled climatic conditions (20, 23, and 27 °C, 65 ± 5% RH, and L16 : D8 photoperiod). The development of the individuals of *A. aurantii* was observed daily using a stereomicroscope (×20) until the death of the adult females, and the following observations were recorded.

### 2.3. Survival, longevity, and fecundity of *Aphytis melinus*

Newly emerged adult parasitoids were shifted to an Agilent glass vial (1.5 mL). The vial was then plugged with a cotton bung. A drop of honey was supplied for each vial. After 24 h, each female parasitoid was confined under a micro plastic cage (small plastic thimble, 3 cm diameter with fine holes on its top) affixed with the help of modelling clay to the surface of butternut squash containing 10 female individuals female of the 3rd instar stage of *A. aurantii*. A virgin female and male of *A. melinus* were confined into each of these cages, as the parasitoid prefers to oviposition on initial stages of this host (Heimpel, 1997). A

stereoscopic magnifying glass was used to observe mating. The parasitoids were transferred every 24 h to a new cage and observations continued until the female parasitoid died (Qiu et al., 2007). Determining the number of eggs laid without removing the armour scale of the nymphs is impossible, as the process leads to the death of the nymph (Foldi, 1990). A parallel experiment, identical to that used in the longevity and oviposition experiment, was therefore undertaken. Nonparasitised individuals were inverted to check for egg remains or unhatched parasitoid eggs, indicating unsuccessful parasitism or infertile eggs, respectively (Urbaneja et al., 2007). Observations were recorded to get the information on preoviposition, oviposition, and postoviposition periods and the number of eggs laid each day. Parasitised nymphs were kept to determine when the adult parasitoid emerged in order to estimate developmental time and survivorship (a number determined from the number emerged divided by total number parasitised). The experimental unit was a group of 10 female parasitoids. Each experimental unit was replicated 3 times.

**2.4. Design and statistical analysis**

A completely randomised design was used in all of the experiments. An analysis of variance (ANOVA) was subsequently performed, and means of survivorship, developmental times, longevity, and fecundity were compared using Tukey's multiple comparison tests (P < 0.05). Subsequently, all data were analysed using Levene's test to ensure homogeneity of variance.

**2.5. Life table parameters**

The main purpose of age-specific fecundity table studies is to predict the rate of population growth of *A. aurantii* and *A. melinus*. The fertility life table was constructed by making a list of the data collected on the developmental and reproductive biology of the species as per formulae provided by Birch (1948). Daily adult numbers from observation from the first day of female exclusion until complete female death, daily survival rate ( $I_x$ ), and female fecundity ( $m_x$ ) were calculated according to the equation of the life tables to predict the growth rate of the population under laboratory conditions at 27 °C. From the fertility and survival rate, several population growth parameters including the net reproductive rate ( $R_0$ ), mean generation time ( $T_0$ ), and intrinsic rate of natural increase ( $r_m$ ) were calculated using the formulas suggested by Carey (1993), where  $x$  is the age of individuals in days,  $I_x$  is the age-specific survival, and  $m_x$  is the age-specific number of female offspring.

Age-specific life table parameters of the insects were calculated according to the Euler-Lotka equation (Birch, 1948). These parameters are:

$$\text{Net reproductive rate } R_0 = \sum I_x m_x;$$

$$\text{Intrinsic rate of natural increase } (r_m);$$

$$l = \sum I_x m_x \cdot e^{-r_m \cdot x}$$

Mean generation time,  $T_0 = \ln R_0 / r_m$ ;  
 Gross reproduction rate,  $GRR = \sum m_x$ ;  
 Finite rate of increase,  $\lambda = e^{r_m}$ .

Age-specific eggs laid by a female were described by the Enkegaard equation:

$$F(x) = a \cdot x \cdot e^{(-b \cdot x)} \text{ (Enkegaard, 1993; Hansen et al., 1999).}$$

Where  $F(x)$  is the daily age-specific fecundity rate (eggs/female/day),  $x$  is the female's age in days, and  $a$  and  $b$  are constants. Day 1 is the first day of the oviposition period. Analyses were done by using JMP (v. 5), MS Excel (v. 2003), SPSS (v. 24.0), and CurveExpert pro (v. 1.6.7) software.

**3. Results**

**3.1. Survival, development, and oviposition of *Aonidiella aurantii***

Table 1 shows the survival rates of the immature state of *A. aurantii* at different temperatures (20, 23, and 27 °C). The rates were similar but varied between 80.0% and 88.0% for these 3 temperatures. The average mortality recorded at the 3 different temperatures was 0.16%. The highest mortality occurred during the preadult stage, with values ranging between 0.12% and 0.20% (depending on the temperature). Based on these results, we assume that there were no significant differences in the survival of *A. aurantii* reared on *C. moschata* at these different temperatures.

The obtained data in Table (2) indicated that the duration of each developmental stage was affected by temperature. These results indicated that 27 °C was an adequate tested temperature for *A. aurantii*, resulting in the highest oviposition (129.35 ± 4.56 eggs/female), the shortest incubation period (4.45 ± 0.14 days), and greatest adult longevity (39.35 ± 1.07 days).

**3.2. Survival, development, and oviposition of *Aphytis melinus***

*A. melinus* mortality was concentrated in older individuals. The adult *A. melinus* individuals started to die from day 10 until day 18 and showed a sharp drop on days 13 and 16 (Figure 1).

The oviposition period of *A. melinus* lasted an average of 15.7 ± 0.52 days (Table 3). The average oviposition rate was 3.95 ± 0.22 eggs female<sup>-1</sup> day<sup>-1</sup>. Each female laid an average total of 62.7 ± 2.81 eggs. The preoviposition period was 0.82 ± 0.05 day; adult longevity was an average of 19.24 ± 0.73 days, while the egg-pupal stage was 11.25 ± 0.12 day.

**3.3. Life table parameters**

Age-specific fecundity rates ( $m_x$ ) of *A. aurantii* reared on fruits of *C. moschata* at 3 constant temperatures (20, 23, and 27 °C) and *Aphytis melinus* (27 °C) are given in Table 4 and Figure 2.


**Table 1.** Effect of temperature on the survival of *Aonidiella aurantii* reared on *Cucurbita moschata* in the laboratory at 20, 23, and 27 °C, 65.0 ± 5.0% RH and 16 L : 8 D photoperiod.

Temperature	n	1st instar %	2nd instar %	3rd instar female %	Gravid female %	Survival, %
20 °C	60	98.33	95.00	95.00	83.33	83.3 a
23 °C	60	100.00	98.33	598.33	88.33	88.3 a
27 °C	60	100.00	98.33	98.33	80.00	80.0 a

**Table 2.** Duration of the California red scale *Aonidiella aurantii* stages at 3 constant temperatures (20, 23, and 27 °C, 65.0 ± 5.0% RH, and 16 L : 8 D photoperiod) reared on fruits of *Cucurbita moschata*.

Developmental stages	Duration (days) ± SE		
	Mean ± SE at 20 °C	Mean ± SE at 23 °C	Mean ± SE at 27 °C
1st instar	12.15 ± 0.38 a*	10.85 ± 0.32 b	8.65 ± 0.27 c
2nd instar	12.65 ± 0.26 a	10.95 ± 0.17 b	8.45 ± 0.24 c
3rd instar	13.05 ± 0.22 a	12.3 ± 0.20 a	9.11 ± 0.44 b
Preoviposition period	8.85 ± 0.26 a	6.15 ± 0.19 b	4.45 ± 0.14 c
Oviposition period	44.35 ± 0.51 a	40.65 ± 0.41 b	34.5 ± 0.45 c
Total eggs female <sup>-1</sup>	73.25 ± 1.83 c	109.7 ± 3.57 b	129.35 ± 4.56 a
Adult longevity	54.65 ± 0.71 a	47.05 ± 0.97 b	39.35 ± 1.07 c
Adult lifespan	93.1 ± 9.73 a	81.8 ± 7.13 b	65.7 ± 6.37 c

\*Means within each row followed by different letters differ significantly based on Tukey’s test (P < 0.05). Data are means ± standard errors.



**Figure 1.** Survivorship of female *Aphytis melinus* at 27 °C, 65 ± 5% RH, and a photoperiod of 16 L : 8 D.

**Table 3.** Oviposition periods, longevity, fecundity, and oviposition rates of *Aphytis melinus* at 27 °C, 65.0 ± 5.0% RH and a photoperiod of 16 L : 8 D.

Developmental stages	Duration of the stages, days <sup>1</sup>
Egg-pupal stage	11.25 ± 0.12
Preoviposition	0.82 ± 0.05
Oviposition	15.7 ± 0.52
Postoviposition	2.21 ± 0.09
Adult longevity	19.24 ± 0.73
Oviposition	Number of eggs
Total eggs female <sup>-1</sup>	62.7 ± 2.81
Eggs female <sup>-1</sup> day <sup>-1</sup>	3.95 ± 0.22

<sup>1</sup> Data are reported as the mean ± standard error.

It was observed that *A. aurantii* had a higher net reproduction rate ( $R_0$ ) and mean generation time ( $T_0$ ), and conversely, intrinsic rate of increase rate ( $r_m$ ) than *A. melinus* (Tables 4). Parameters show that the *A. aurantii* population grows 61.02 times at 27 °C in one generation. The net reproduction rate for each female of a generation was considerably higher for *A. aurantii* than *A. melinus*, reaching 28.14 at 27 °C (Table 4), which indicates its high reproductive capacity when fed butternut squash.


Moreover, the net reproduction rate obtained for *A. melinus* indicates a low substitution potential for each female having *A. aurantii* as a host under laboratory conditions. The intrinsic rate of increase ( $r_m$ ), defined as a



**Table 4.** Life table parameters and statistics of *Aonidiella aurantii* and *Aphytis melinus* parasitizing on butternut squash (*Cucurbita moschata*) under laboratory conditions.

Parameter	<i>Aonidiella aurantii</i>			<i>Aphytis melinus</i>
	20 °C	23 °C	27 °C	27 °C
Net reproductive rate ( $R_0$ )	30.79 a*	48.78 b	61.02 c	28.14 a
Mean Generation time ( $T_0$ )	62.31 a	55.53 b	51.39 c	19.06 d
Intrinsic rate of increase ( $r_m$ )	0.055 a	0.070 b	0.080 c	0.188 d
Finite rate of increase ( $\lambda$ )	1.056 a	1.072 b	1.083 c	1.207 d
Gross reproduction rate (GRR)	37.09 a	55.43 b	65.16 c	32.15 d

\*Means within each row followed by different letters differ significantly based on Tukey's test ( $P < 0.05$ ). Data are means  $\pm$  standard errors.



**Figure 2.** Age-specific fecundity ( $m_x$ ) of *Aonidiella aurantii* reared on fruits of *Cucurbita moschata* at 3 constant temperatures (20, 23, and 27 °C) and *Aphytis melinus* (A.m) (27 °C).


population's capacity to multiply, was higher for *A. melinus* (0.188) than *A. aurantii* which was 0.080 at 27 °C (Table 4).

The parameters (a, b, and  $R^2$ ) of the Enkegaard regression model applied on the age-specific fecundity rate ( $m_x$ ) of *A. aurantii* and *A. melinus* are given in Figure 3. The relationship between days and fecundity was higher at all temperatures.

**4. Discussion**

Since the 1960s, there have been intensive studies on mortality and growth of *A. aurantii* conducted worldwide (DeBach et al., 1971; Abdelrahman, 1974; Karaca et al., 1987; Badary and Abd-Rabou, 2010). Information about the biology of *A. aurantii* is important for improving the mass production of certain parasitoids used in biological

control, particularly *A. melinus*, a species that is used to control *A. aurantii* (Olivas et al., 2011). The use of constant temperature provides a useful reference point for the performance of *A. aurantii*. The survival values of *A. aurantii* varied between 80% and 88%. These values were approximately similar to those obtained by Karaca (1990) at 26 °C on lemon, orange, and grapefruit, and higher than the survival of *A. aurantii* on mandarin, in a study on the development, survival, and longevity of *A. aurantii* on different citrus species, in which survival was only 0.90, 0.81, 0.85, and 0.61 on lemon, orange, grapefruit and mandarin, respectively. This could be due to the variety of the fruit species and laboratory conditions being different from those used in this study. The average mortality is approximately similar to that obtained by Vanaclocha



**Figure 3.** Enkegaard distribution of *Aonidiella aurantii* and *Aphytis melinus*.

et al. (2012) on lemon at 26 °C, for which the mortality percentage was 15%, and to those obtained by Karaca (1990) on lemon and orange, grapefruit, and higher than those in mandarin, where the highest mortality (39%) was recorded in adults reared on mandarin fruits.

The analysis of variance in Table 2 revealed that the total life cycle and development rates of the female of *A. aurantii*

were significantly different depending on the difference in the temperature used, except for the duration of the third instar which did not differ significantly between 20 °C and 23 °C. In addition, the preoviposition period, oviposition period, longevity, and the mean number of eggs per female of *A. aurantii* differed significantly depending on the difference in temperature. In general, the duration of each developmental stage recorded in this study was consistent with those determined for *A. aurantii* by Badary and Abd-Rabou (2010), and less than those determined by Karaca (1990). In other species of *Aonidiella*, the average developmental time ranged between 45 days and 94 days depending on the temperature used for *A. orientalis* (Flaih, 2007), 65 days for *A. citrina* in California; the reproductive period lasts 60 days under a constant temperature of 27.8 °C (Nel, 1933). Temperature is the most important abiotic factor affecting insect growth, development rate, oviposition, and survival of *A. aurantii*. With high temperatures within optimal limits, all processes occur significantly faster, which results in rapid ageing of the females with a significant increase in the total number of eggs laid and reduced longevity. Likewise, the number of eggs laid per day increases (Pekas, 2011b). This last parameter, together with the increased rate of oviposition and nymphal development, increases the potential pest status of *A. aurantii* in warm citrus-growing areas.

*Aphytis melinus* females showed gradual mortality over time, which can be attributed to death by natural ageing of the individuals. The developmental period of *Aphytis* spp. is usually short and depends on climatic conditions, principally temperature and humidity (Yu and Luck, 1988). For example, at 26.7 °C, *A. melinus* completes its development in almost a fortnight, whereas it takes 1 month to complete development at 17 °C. Nutrition also contributed to longevity in the ectoparasitoid species *A. melinus*. Higher longevity was observed in females that were fed than those not allowed to host-feed (Heimpel et al., 1997). As a result, greater availability of females could occur over time when the temperature is appropriate and there is food, but this is not necessarily associated with an increase in the parasitoidism rate. *Aphytis melinus* has been found to have 2–3 generations to 1 generation of its host *A. aurantii* (Yu and Luck, 1988). The duration of the longevity period of *A. melinus* is approximately consistent with that obtained by Vanaclocha et al. (2012), who determined that at 26.7 ± 1.5 °C the duration of the longevity period of *A. melinus* was 20.5 days, while the total number of eggs per female was about 11.7 eggs fewer than the number produced by *A. melinus* at 26.7 ± 1.5 °C [39]. The diet of adult parasitoids has an effect on lifetime reproductive success (Flaih, 2007). Adult females of many species obtain materials required for egg maturation by feeding upon host insects (host-

feeding), and materials necessary for adult maintenance and survival are acquired by host-feeding and by feeding upon any of a number of sugar sources (Jervis, 1996; Heimpel and Collier, 1996).

These results confirm the potential of *A. melinus* to control *A. aurantii*. However, this characteristic must be compared with that of other species, such as other *Aphytis* spp., which can use the same field substrate. Studies carried out by Orphanides (1984) point out that there is interspecific competition and competitive displacement between *Aphytis* spp. which are *A. aurantii* parasitoids; thus, other species or subspecies could affect biological control success in the field. Comparing the life table parameters of *A. aurantii* and *A. melinus*, it can be seen that there are significant differences in all 5 of the parameters studied (Table 4). The net reproductive rate (number of females for each female of a generation) is significantly higher for *A. aurantii* (61.02) than that of *A. melinus* (28.14). Generation time, the mean time between 2 successive generations, is significantly longer in *A. aurantii* (51.39 days) than in *A. melinus* (19.06

days) (Table 4); this is interpreted as favourable for increasing the numbers and efficiency of the parasitoid (La Rossa et al., 2002). The intrinsic rate of increase ( $r_m$ ), which indicates the ability of a population to increase in abundance from generation to generation, is an essential indicator of the potential of a parasitoid to control its host (Mercado et al., 2014; Persad and Khan, 2002). The  $r_m$  parameter of *A. melinus* (0.188) is significantly greater than that of *A. aurantii* (0.080) at 27 °C (Table 4).

The relationship between days and fecundity was demonstrated well by using the Enkegaard regression model; at 20, 23, and 27 °C for *A. aurantii* ( $R^2 = 0.833$ ,  $a = 0.410$ ,  $b = 0.099$ ;  $R^2 = 0.799$ ,  $a = 0.624$ ,  $b = 0.098$ ;  $R^2 = 0.715$ ,  $a = 0.661$ ,  $b = 0.091$ , respectively) and at 27 °C for *A. melinus* ( $R^2 = 0.841$ ,  $a = 1.190$ ,  $b = 0.179$ ). Most of the eggs were laid within the first half of the oviposition period (Figure 3). The influence of increased temperature was clearly seen for *A. aurantii* by increased the fecundity peak at an earlier age. Thus, the fecundity curve turns to the left at 27 °C earlier than at other temperatures; this is clearly seen for *A. melinus* as well (Figure 3).

## References

- Abdelrahman I (1974). Studies in ovipositional behaviour and control of sex in *Aphytis Melinus* Debach, a parasite of California red scale, *Aonidiella Aurantii* (Mask.). Australian Journal of Zoology 22 (2): 231-247. doi: 10.1071/ZO9740231
- Abou Hatab EE (1999). Ecological and biological studies on the red scale, *Aonidiella aurantii* (Mask) (Homoptera: Diaspididae). MSc, Mansoura University, Mansoura, Egypt.
- Badary H, Abd-Rabou S (2010). Biological studies of the California red scale, *Aonidiella aurantii* (Maskell) (Hemiptera: Diaspididae) under different host plants and temperatures with an annotated list of natural enemies of this pest in Egypt. Egyptian Academic Journal of Biological Sciences 3: 235-242.
- Bayoumy M, Abdel-Salam A, Abdel-Baky N, Schopf A, Abdel-Kareim AS (2009). Age-specific life tables of the aphelinid wasp, *Aphytis diaspidis* and its diaspidid hosts, *Hemiberlesia lataniae* and *Quadraspidiotus perniciosus* with special reference to the intrinsic rate of increase. Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie 17: 235-240.
- Beardsley Jr JW, Gonzalez RH (1975). The biology and ecology of armored scales. Annual Review of Entomology 20 (1): 47-73. doi: 10.1146/annurev.en.20.010175.000403
- Ben-Dov Y (2006). Taxonomy of *Aonidiella yehudithae* sp. nov. and *Lindingaspis misrae* (Laing) comb. nov. with a key to species of *Aonidiella* Berlese & Leonardi (Hemiptera: Coccoidea: Diaspididae). Zootaxa 1: 51-57. doi:10.11646/zootaxa.1190.1.2
- Birch L (1948). The intrinsic rate of natural increase of an insect population. The Journal of Animal Ecology 1: 15-26. doi: 10.2307/1605
- Carey JR (1993). Applied Demography for Biologists: With Special Emphasis on Insects. Oxford, UK: Oxford University Press.
- DeBach P, Rosen D, Kennett CE (1971). Biological control of coccids by introduced natural enemies. In: Biological Control. Boston, MA, USA: Springer, 165-194. doi: 10.1007/978-1-4615-6531-4\_7
- Enkegaard A (1993). The poinsettia strain of the cotton whitefly, *Bemisia tabaci* (Homoptera; Aleyrodidae), biological and demographic parameters on poinsettia (*Euphorbia pulcherrima*) in relation to temperature. Bulletin Entomological Research 83: 535-546.
- Erler F, Tunç IA (2001). Survey (1992–1996) of natural enemies of Diaspididae species in Antalya, Turkey. Phytoparasitica 29 (4): 299-305. doi: 10.1007/BF02981846.
- Flaih SK (2007). Biological studies of the oriental yellow scale insect *Aonidiella orientalis* (Newstead) (Homoptera: Diaspididae) and the efficacy of three introduced Hymenopterous Parasitoids. MSc, Agriculture College, University of Baghdad, Baghdad, Iraq [in Arabic].
- Foldi I (1990). The scale cover. Armoured Scale Insects, their Biology, Natural Enemies and Control. Amsterdam: Elsevier, pp. 43-54.
- Food and Agriculture Organisation of the United Nations (2017). Citrus Fruit - Fresh and Processed Statistical Bulletin. Rome: FAO.
- Forster LD, Luck RF, Grafton-Cardwell EE (1995). Life stages of California red scale and its parasitoids. Oakland, CA, USA: University of California Publications, Division of Agriculture and Natural Resources.

- Furness GO, Buchanan GA, George RS, Richardson NL (1983). A history of the biological and integrated control of red scale, *Aonidiella aurantii* on citrus in the Lower Murray Valley of Australia. *Entomophaga* 28 (3): 199-212. doi: 10.1007/BF02372170
- Garcerá C, Chueca P, Santiago S, Moltó E (2008). Biological efficacy of two organophosphate insecticides against California red scale (*Aonidiella aurantii* Maskell) related to deposition parameters under laboratory conditions. *Integrated Control in Citrus Fruit Crops* 38: 67-74.
- Grafton-Cardwell EE, Faber BA, Haviland DR, Morse JG, O'Connell NV et al. (2006). UC IPM pest management guidelines: citrus, insects, mites and snails. Oakland, CA, USA: University of California Agriculture and Natural Resources Publication 3441.
- Hansen DL, Brødsgaard HF, Enkegaard A (1999). Life table characteristics of *Macrolophus caliginosus* preying upon *Tetranychus urticae*. *Entomologia Experimentalis et Applicata* 93: 269-275.
- Heimpel GE, Collier TR (1996). The evolution of host-feeding behaviour in insect parasitoids. *Biological Reviews* 71(3): 373-400. doi: 10.1111/j.1469-185X.1996.tb01279.x
- Heimpel GE, Rosenheim JA, Kattari D (1997). Adult feeding and lifetime reproductive success in the parasitoid *Aphytis melinus*. *Entomologia Experimentalis et Applicata* 83 (3): 305-315. doi: 10.1046/j.1570-7458.1997.00185.x
- Hely PC, Pasfield G, Gellatley JG (1982). *Insect Pests of Fruit and Vegetables in NSW*. Melbourne: Inkata Press.
- Hill DS (2008). *Pests of Crops in Warmer Climates and Their Control*. Dordrecht, The Netherlands: Springer Science & Business Media.
- Jervis MA (1996). Parasitoid adult feeding behaviour and biocontrol-a review. *Biocontrol News Inform* 17: 11-26.
- Karaca İ (1990). Doğu Akdeniz bölgesi turuncgillerinde zararlı olan *Aonidiella* (Homoptera: Diaspididae) türleri, yayılışları ve değişik turuncgill çeşitlerinde *Aonidiella aurantii* (Mask.) nin populasyon gelişmesinin saptanması. PhD, Çukurova University, Adana, Turkey (in Turkish).
- Karaca İ, Uygun N (1992). Population development of California red scale, *Aonidiella aurantii* (Maskell) (Homoptera : Diaspididae) on different citrus varieties. *Proceedings of the Second Turkish National Congress of Entomology*; Adana, Turkey, pp. 9-19.
- Karaca İ, Şekeroğlu E, Uygun N (1987). Life table of California red scale, *Aonidiella aurantii* (Mask.) (Homoptera: Diaspididae) in laboratory conditions. *Proceedings of the First Turkish National Congress of Entomology*; İzmir, Turkey, pp. 127-138.
- La Rossa FR, Crespo DC, Lecuona RE (2002). Population parameters of *Spalangia endius* Walker (Hymenoptera: Pteromalidae) on pupae of *Musca domestica* L. (Diptera: Muscidae). *Neotropical Entomology* 31 (4): 597-600. doi: 10.1590/S1519-566X2002000400012
- Malipatil MB, Smith D, Dunn KL (2000). An illustrated guide to the parasitic wasps associated with citrus scale insects and mealybugs in Australia. *Australian Journal of Entomology* 39: 353-357. doi: 1440-6055.2000.0188a.x
- Mercado VT, Fernández ES, Giliomee JH (2014). Life table parameters of the woolly whitefly *Aleurothrixus floccosus* (Hemiptera: Aleyrodidae) and its parasitoid *Cales noacki* (Hymenoptera: Aphelinidae). *European Journal of Entomology* 111 (2): 251-256. doi: 10.14411/eje.2014.020
- Miller DR (2005). Selected scale insect groups (Hemiptera: Coccoidea) in the southern region of the United States. *Florida Entomologist* 88 (4): 482-501. doi:10.1653/0015-4040(2005)88[482:SSIGHC]2.0.CO;2
- Nel RGA (1933). Comparison of *Aonidiella aurantii* and *Aonidiella citrina*, including a Study of the Internal Anatomy of the Latter. *Hilgardia* 7 (11): 417-466.
- Ode PJ, Hardy ICW (2008). Parasitoid sex ratios and biological control. In: Wajnberg E, Bernstein C, van Alphen J, editors. *Behavioural Ecology of Insect Parasitoids: From Theoretical Approaches to Field Applications*. Oxford, UK: Blackwell, 30: 253-291.
- Olivas J, Lucas A, Calvo J, Belda JE (2011). Biological control of diaspidid scales *Aspidiotus nerii* Bouche and *Aonidiella aurantii* (Maskell) by means of commercial *Aphytis melinus* DeBach (Hym., Aphelinidae) releases: basis for an IPM strategy in Spanish citrus. *IOBC/WPRS Bulletin* 62: 311-316.
- Opp SB, Luck RF (1986). Effects of host size on selected fitness components of *Aphytis melinus* and *A. lingnanensis* (Hymenoptera: Aphelinidae). *Annals of the Entomological Society of America* 79 (4): 700-704. Doi: 10.1093/aesa/79.4.700
- Orphanides GM (1984). Competitive displacement between *Aphytis* spp. [Hym. Aphelinidae] parasites of the California red scale in cyprus. *Entomophaga* 29 (3): 275-281. doi: 10.1007/BF02372114
- Pekas A (2011a). Biological pest control in citrus an alternative to chemical pesticides with benefits for essential oil quality. IFEAT International Conference; Barcelona, Spain, pp. 115-124.
- Pekas A (2011b). Factors affecting the biological control of California red scale *Aonidiella aurantii* (Hemiptera: Diaspididae) by *Aphytis* (Hymenoptera: Aphelinidae) in eastern Spain citrus: host size, ant activity, and adult parasitoid food sources (Doctoral dissertation). PhD, Universitat Politècnica de València, València, Spain. doi: 10.4995/Thesis/10251/10293
- Persad A, Khan A (2002). Comparison of life table parameters for *Maconellicoccus hirsutus*, *Anagyrus kamali*, *Cryptolaemus montrouzieri* and *Scymnus coccivora*. *BioControl* 47 (2): 137-149. doi: 10.1023/A:1014581616965
- Qiu BL, De Barro PJ, Ren SX, Xu CX (2007). Effect of temperature on the life history of *Eretmocerus* sp. nr. *furuhashii*, a parasitoid of *Bemisia tabaci*. *BioControl* 52 (6): 733-746. doi: 10.1007/s10526-006-9064-y



- Smith D, Beattie GA, Broadley R (1997). Citrus Pests and Their Natural Enemies: Integrated Pest Management in Australia. Queensland: Department of Primary Industries. URL: <http://hdl.handle.net/10462/pdf/9446>
- Broughton S (2006). Development of an Integrated Pest Management Strategy for Citrus in Western Australia: Final project. South Perth, WA, Australia: Horticulture Australia Limited and Department of Agriculture and Food Western Australia.
- Urbaneja A, Sánchez E, Stansly PA (2007). Life history of *Eretmocerus mundus*, a parasitoid of *Bemisia tabaci*, on tomato and sweet pepper. *BioControl* 52 (1): 25-39. doi: 10.1007/s10526-006-9014-8
- Uygun N, Satar S (2008). The current situation of citrus pests and their control methods in Turkey. *IOBC-WPRS Bulletin* 38: 2-9.
- Uygun N, Karaca İ, Şekeroğlu E (1995). Population dynamics of *Aonidiella aurantii* (Maskell) (Diaspididae) and its natural enemies on citrus from 1976 to 1993 in the Mediterranean region of Turkey. *Israel Journal of Entomology* 29: 239-246.
- Vacas S, Alfaro C, Navarro-Llopis V, Primo J (2010). Mating disruption of California red scale, *Aonidiella aurantii* Maskell (Homoptera: Diaspididae), using biodegradable mesoporous pheromone dispensers. *Pest Management Science* 66 (7): 745. doi: 10.1002/ps.1937
- Vanaclocha P, Vacas S, Alfaro C, Primo J, Verdú MJ et al. (2012). Life history parameters and scale-cover surface area of *Aonidiella aurantii* are altered in a mating disruption environment: implications for biological control. *Pest Management Science* 68 (7): 1092-1097. doi: 10.1002/ps.3273
- Yu DS, Luck RF (1988). Temperature-dependent size and development of California red scale (Homoptera: Diaspididae) and its effect on host availability for the ectoparasitoid, *Aphytis melinus* DeBach (Hymenoptera: Aphelinidae). *Environmental Entomology* 17 (2): 154-161. doi: 10.1093/ee/17.2.154