

Comparative wood anatomy of *Rhodothamnus* species

Bedri SERDAR*

Department of Forest Botany, Faculty of Forestry, Karadeniz Technical University, 61080 Trabzon, Turkey

Received: 04.04.2012

Accepted: 06.12.2012

Published Online: 15.05.2013

Printed: 30.05.2013

Abstract: In this study, the comparative wood anatomy of the European [*Rhodothamnus chamaecistus* (L.) Reichb.] and Anatolian (*Rhodothamnus sessilifolius* P.H.Davis) species of *Rhodothamnus* were studied. The wood anatomy of the taxa shows evidence of adaptation to growing in alpine habitats. The woods of the species exhibit primitive wood anatomical characteristics and share similar qualitative anatomical features. However, some of the quantitative anatomical characteristics of the taxa show significant differences, such as the distinctness of the growth ring and the bar number of the scalariform perforation plate. The present study describes and compares the anatomical properties of the wood of the *Rhodothamnus* species.

Key words: *Ericaceae*, *Rhodothamnus*, wood anatomy

1. Introduction

The genus *Rhodothamnus* Reichb. has 2 species in the world: *Rhodothamnus chamaecistus* (L.) Reichb. is confined to the eastern Alps of southern Europe (Greguss, 1959), and the other species, *Rhodothamnus sessilifolius*, is endemic to the north-eastern corner of Turkey (Stevens, 1978). *R. chamaecistus* is confined to crevices in calcareous (limestone or dolomite) rocks (Stevens, 1978), whereas on Tiryal Mountain, *R. sessilifolius* grows on igneous dacite rock outcrops that form cliffs or ridges (Terzioğlu & Milne, 2002). An emended description for *R. sessilifolius* was given by Terzioğlu and Milne (2002) based upon specimens collected from north-eastern Turkey in 2000, which was the first gathering of this species since 1960. *R. sessilifolius* was first collected on Tiryal Mountain above Murgul on 23 June 1957 by Davis and Hedge (Davis 29974 & Hedge), and then again from an adjacent mountain range (Şavval Tepe) in July 1960 (Stevens, 1978). The species has since then been known only from these 2 localities in the north-eastern corner of Turkey, in Artvin Province. The type locality is a large rock at 2150 m on Tiryal Mountain.

Comparative wood anatomy consists of 2 main efforts: wood identification and evolutionary studies. Evolutionary studies can be divided into 2 main areas: systematic wood anatomy and ecological wood anatomy (Olson, 2005; Güvenç & Kendir, 2012; Eo & Hyun, 2013; Tiwari et al., 2013).

The 2 species of the genus are closely related to each other by means of morphology (Davis, 1962). The goal of

the present study is to define the wood anatomical traits in order to contribute to their identification.

2. Materials and methods

The wood samples that were studied were taken from KATO herbarium (*R. sessilifolius*, KATO: 13360 and *R. chamaecistus*, KATO: 10501) materials. Wood samples, from stems about 1.5–3.0 mm diameter in each case, were boiled in water and stored in 50% aqueous ethanol, sectioned using a freezing sliding microtome at a thickness of about 20–25 µm, and then stained with a safranin and alcian blue combination (Ives, 2001). The permanent slides were examined and photographed with an Olympus BX 50 research microscope (Bs200Prop Image Processing and Analysis Systems). Wood portions from each species were macerated using Schultzze's method (Normand, 1972) and stained with safranin. All wood terms used conform to the usage of the International Association of Wood Anatomists (IAWA) Committee on Nomenclature (1989).

3. Results and discussion

In the present study, the anatomical features of the wood of the *Rhodothamnus* species were studied and are given in detail in the following text.

Rhodothamnus chamaecistus: Wood diffuse porous with distinct growth rings in contrast to *R. sessilifolius*. Vessels evenly distributed without any tendency to a specific pattern, many to numerous (370–1010 vessel/mm²), very small (9.33–16.79 µm, 9.33–22.39 µm in tangential and

* Correspondence: bserdar@ktu.edu.tr

radial diameters, respectively), angular in cross-section, exclusively solitary, and mean number of vessels per group 2.32, with thin walls (1.75 μm) (Figure). Vessel elements short (288–720 μm), perforation plates scalariform with many to numerous bars (13–53 per perforation plate), intervessel pits mostly scalariform, sometimes tending to opposite. Helical thickening not observed in vessel lateral walls and ligulate ends. Fibre tracheids 254–628 μm long, 9.33–14.93 μm wide with rather thin walls (1.86–4.67 μm), with bordered pits on radial and tangential walls. Axial

parenchyma is not abundant, apotracheal diffuse, and scanty paratracheal. Rays homogeneous and uniseriate, composed of only upright cells, ray 86–202 μm high (Figure).

Rhodothamnus sessilifolius: Wood diffuse porous with indistinct growth rings. Vessels evenly distributed without any tendency to a specific pattern, many to numerous (720–1150 vessel/ mm^2), very small (9.33–20.52 μm , 11.19–20.52 μm in tangential and radial diameters, respectively), angular in cross-section,




Figure. a, b, and c- *Rhodothamnus chamaecistus*; d, e, and f- *R. sessilifolius*. a- TS, wood diffuse porous, growth ring boundaries distinct with thick-walled and radially flattened latewood fibres. b- RLS, rays homogeneous, composed of only upright cells. c- TLS, uniseriate rays. d- TS, wood diffuse porous, growth ring boundaries indistinct. e- RLS, scalariform perforation plate. f- TLS, scalariform intervessel pits. Scale bars: a, d = 50 μm ; b, c, e, f = 10 μm . Abbreviations: TS: transverse section, RLS: radial longitudinal section, TLS: tangential longitudinal section.

exclusively solitary, and mean number of vessels per group 1.12, with thin walls (0.90 μm) (Figure). Vessel elements short (163–432 μm), perforation plates scalariform with many to numerous bars (11–60 per perforation plate), intervessel pits mostly scalariform, sometimes tending to opposite, vessel-ray pits similar to intervessel pits (Figure). Helical thickening not observed in vessel lateral walls and ligulate ends. Fibre tracheids 163–542 μm long, 7.46–13.06 μm wide with rather thin walls (1.86–4.66 μm), with bordered pits on radial and tangential walls. Axial parenchyma is not abundant and scanty paratracheal. Rays homogeneous and uniseriate, composed of only upright cells, ray 103–448 μm high.

The growth rings are very narrow to uncountable in the *Rhodothamnus* species. The width of the growth rings is variable according to the growth rate in many dicotyledonous woods. In members of Ericaceae, the growth rings are narrower in dwarf shrubs and shrubs, while wider in small trees and trees. Many authors regard this as a reflection of the slow growth associated with the shrublet/shrub habit (Suzuki & Ohba, 1988; Noshiro et al., 1995a, 1995b).

These 2 dwarf shrub species of *Rhodothamnus* differ in features of growth rings, the number of vessels/ mm^2 , and the dimensions of the vessels, fibres, and rays (Table). Variation in vessel diameter with relation to seasonality was evident in these species, which had very narrow and short vessels. This characteristic might reflect the habit and/or result from growing in dry sites at high altitudes. Narrowness of vessels is inversely correlated with the number of vessels/ mm^2 . Furthermore, vessel number was found to increase with altitude in *Rhododendron* (Merev & Yavuz, 2000). According to Carlquist (1977), short and narrow vessel elements are theorised to resist high tension in the water column. The safety and efficiency of water transport are strongly related to vessel diameter and

vessel density, and decreasing vessel diameter increases the safety of water conduction (Zimmermann, 1983; Baas & Wheeler, 1996).

Forsaitth (1920) considered that narrower vessels and uniseriate or narrower rays were all reduced in size by the influence of alpine conditions. This implies that wood structure can vary according to ecological factors, and is therefore not solely determined by the plant's genetically determined habit.

In the present study, the 2 dwarf shrubs had fibre with distinctly bordered pits. The taxa examined here both had uniseriate ray tissues that comprised upright cells only (Figure). The longest dimensions of the ray parenchyma cells are oriented in the axial direction of the plant; these were observed both in the radial and tangential sections.

The following wood anatomical characters are considered primitive in the Baileyan sense (Baas et al., 2000): scalariform perforation plates, fibre with distinctly bordered pits, apotracheal parenchyma, and heterocellular rays. In *Rhodothamnus*, lateral wall pitting is scalariform to opposite, which might account for the presence of scalariform perforation plates. Perforation plates comprising numerous bars per perforation plate and scalariform lateral wall pitting are primitive characteristics. Furthermore, according to classical evolutionary theory (Baileyan sense), a higher number of bars indicates a more primitive species, whereas fewer bars tend to indicate one that is more evolved.

Acknowledgements

I would like to extend my thanks to Dr Nesime Merev for her thoughtful reviews of this manuscript and to VI Gorissen for supplying the material. I would also like to thank the anonymous reviewers for their valuable help on improving an earlier draft of this paper.

Table. Selected qualitative and quantitative data (min–max) showing differences between *Rhodothamnus sessilifolius* and *Rhodothamnus chamaecistus*.

Feature	<i>R. sessilifolius</i>	<i>R. chamaecistus</i>
Growth ring	indistinct	distinct
Vessel elements' length (μm)	163–432	288–720
Number of vessels/ mm^2	720–1100	370–1010
Bar number per perforation plate	11–60	13–53
Fibre length (μm)	163–542	254–628
Ray height (μm)	103–448	86–202

References

- Baas P & Wheeler E (1996). Parallelism and reversibility in xylem evolution a review. *IAWA Journal* 17: 351–364.
- Baas P, Wheeler E & Chase M (2000). Dicotyledonous wood anatomy and APG system of Angiosperm classification. *Botanical Journal of the Linnean Society* 134: 3–17.
- Carlquist S (1977). Wood anatomy of Tremandraceae: phylogenetic and ecological implications. *American Journal of Botany* 64: 704–713.
- Davis PH (1962). *Rhodothamnus sessilifolius* P.H. Davis. *Icones Plantarum* 36: 3575.
- Eo JK & Hyun JO (2013). Comparative anatomy of the needles of *Abies koreana* and its related species. *Turkish Journal of Botany* 37: 553–560.
- Forsyth CC (1920). Anatomical reduction in some alpine plants. *Ecology* 1: 124–135.
- Greguss P (1959). *Holzanatomie der europäischen Laubhölzer und Straucher*. Budapest: Akademiai Kiado.
- Güvenç A & Kendir G (2012). The leaf anatomy of some *Erica* taxa native to Turkey. *Turkish Journal of Botany* 36: 253–262.
- IAWA Committee on Nomenclature (1989). IAWA list of microscopic features for hardwood identification. *IAWA Bulletin* 10: 221–332.
- Ives E (2001). *A Guide to Wood Microtomy*. Suffolk Offset: Suffolk, UK.
- Merev N & Yavuz H (2000). Ecological wood anatomy of Turkish *Rhododendron* L. (Ericaceae): intraspecific variation. *Turkish Journal of Botany* 24: 227–237.
- Normand D (1972). *Manuel D'Identification des Bois Commerciaux*. Tom 1. Paris: Nogent, Sur/Marne.
- Noshiro S, Joshi L & Suzuki M (1995a). Ecological wood anatomy of Nepalese *Rhododendron* (Ericaceae) I. Interspecific variation. *Journal of Plant Research* 108: 1–9.
- Noshiro S, Joshi L & Suzuki M (1995b). Ecological wood anatomy of Nepalese *Rhododendron* (Ericaceae) II. Intraspecific variation. *Journal of Plant Research* 108: 217–233.
- Olson ME (2005). Commentary: typology, homology, and homoplasy in comparative wood anatomy. *IAWA Journal* 26: 507–522.
- Stevens PF (1978). *Rhodothamnus* Reichb. In: Davis PH (ed.) *Flora of Turkey and the East Aegean Islands*, Vol. 6, p. 95. Edinburgh: Edinburgh University Press.
- Suzuki M & Ohba H (1988). Wood structural diversity among Himalayan *Rhododendron*. *IAWA Bulletin* 9: 317–326.
- Terzioğlu S & Milne RI (2002). An emended description for *Rhodothamnus sessilifolius* P.H. Davis (Ericaceae). *Edinburgh Journal of Botany* 59: 291–294.
- Tiwari SP, Kumar P, Yadav D & Chauhan DK (2013). Comparative morphological, epidermal and anatomical studies of *Pinus roxburghii* needles at different altitude from North West Indian Himalaya. *Turkish Journal of Botany* 37: 65–73.
- Zimmermann MN (1983). *Xylem Structure and the Ascent of Sap*. Berlin: Springer-Verlag.