

## A new species from Türkiye, *Centaurea turhanii* (Cardueae, Asteraceae)

Tuna UYSAL<sup>1\*</sup>, Meryem BOZKURT<sup>1</sup>, Ergin HAMZAOĞLU<sup>2</sup>

<sup>1</sup>Department of Biology, Faculty of Science, Selçuk University, Konya, Türkiye

<sup>2</sup>Department of Mathematics and Science Education, Gazi Faculty of Education, Gazi University, Ankara, Türkiye

Received: 16.04.2023 • Accepted/Published Online: 16.11.2023 • Final Version: 16.01.2024

**Abstract:** In this article, a new cornflower, *Centaurea turhanii* Uysal & Hamzaoglu (Asteraceae) from Türkiye was described and reported. Morphological, karyological, and molecular analyses were used to reveal the taxonomic status of the new species with the species thought to be related to taxa. A modified determination key was submitted and its ecology and phenology were given.

**Key words:** Endemic, karyomorphology, molecular, SEM, Türkiye

### 1. Introduction

An important and relatively large genus for The Flora of Turkey (Wagenitz, 1975), *Centaurea* L. (Asteraceae) shows a broad distribution across Eurasia, especially in Irano-Turanian and Mediterranean Regions (Susanna and García-Jacas, 2007). Even the genus was taxonomically problematic, the number of species in the genus is increasing day by day, especially in Asia minor. This shows that the genus is still evolving in terms of speciation. Various methods and approaches have been applied by different researchers for a long time to solve taxonomic problems within the genus *Centaurea* (Romaschenko et al., 2004; Garcia-Jacas et al., 2000; 2001; 2006; Uysal et al., 2009, 2010, 2015, 2016, 2017a; Hilpold et al., 2014; Negaresh and Rahiminejad 2018). Although many species have been transferred to different genera (Wagenitz and Hellwig, 2000; Greuter, 2003; Greuter and Raus, 2009; Negaresh, 2019), Türkiye is still the main diversity center of the genus *Centaurea* (Wagenitz, 1986). The discovery of many new species in the time is strong evidence to support this (Yıldırım, 2012; 2018; Köse and Alan, 2013; Bancheva et al., 2014; Bancheva and Kaya, 2015; Bona, 2015, 2016; Negaresh et al., 2015; Yüzbaşıoğlu et al., 2015; Kültür et al., 2016; Pınar, 2016, Uysal et al., 2016, 2017b; Uysal and Hamzaoglu, 2017; Behçet et al., 2017; Armağan and Uysal, 2018; Negaresh and Rahiminejad, 2018; Şirin et al., 2019, 2020, 2022; Hamzaoglu and Koç 2020; Duman et al., 2021; Özbek, 2021; Uysal et al., 2022). According to the List of Plants of Türkiye (Uysal, 2012), while 194 taxa in Türkiye represent the genus *Centaurea*, this number has currently reached 221 with the latest additions and new

records (Uysal, 2012; Hamzaoglu and Koç, 2020; Duman et al., 2021; Özbek, 2021; Şirin et al., 2022; Uysal et al., 2022). Of these taxa, 134 are endemic and the endemism rate of the genus for Türkiye is approximately 60%. This study introduced a new species of *Centaurea* to the scientific world as based on morphological, karyological, and molecular data.

### 2. Materials and methods

**Morphology:** Interesting *Centaurea* specimens were collected around Kayseri in 2019. Interesting specimens of the genus were examined using the Flora of Turkey and the East Aegean Islands and the neighboring country floras (Wagenitz, 1975, 1980; Dostál J (1976); Tzvelev, 2001). They were compared with the specimens found in KNYA, GOET, E and K herbariums. In the morphologic works, fifteen specimens were used and they were deposited KNYA and GAZI herbarias. Acronyms are according to Thiers (2023).

**SEM micromorphology:** For achene micromorphology, achenes of new and closely related species (at least 10) were first passed through a series of different concentrations of alcohol (70%, 80%, 96%, and 100%) (Kashi et al., 2014). Then, after the surfaces of the achenes were coated with gold, images of the achene surface at different magnifications were taken under a high vacuum in a ZEISS EVO LS-10 scanning electron microscope (SEM).

**Karyology:** Mature seeds of the new species and related taxa were selected and germinated for karyomorphological analyses. Root meristems from germinating seeds were

\* Correspondence: [tuyisal@selcuk.edu.tr](mailto:tuyisal@selcuk.edu.tr)

pretreated with 0.002 M 8-hydroxyquinoline at 4 °C for 8 h. Then, the material was fixed with Carnoy for 24 h in the fridge at 4 °C. Before staining, the material was hydrolyzed with 5 M HCl for 1 h at room temperature, stained with 1% aceto-orcein and mounted in 45% acetic acid. At least 10 metaphases were examined per taxa; the best metaphase plates were photographed (100×) with a digital camera (Olympus DP-72), mounted on an Olympus BX53 microscope. Karyotype asymmetry, ideograms, and karyograms of these taxa were made using the KAMERAM analysis system. While chromosome terminology was made according to the design proposed by Levan et al. (1964), karyotype asymmetries were calculated according to the indices given in the literature (Romero Zarco 1986; Paszko, 2006; Peruzzi and Eroğlu 2013).

**DNA isolation, PCR amplification, and data analysis:** DNA isolation of the new species and some taxa (Table 1) were performed on leaves from herbarium specimen (Doyle and Doyle, 1987; Soltis et al., 1991; Cullings, 1992). The ITS region was amplified according to the method given by Garcia-Jacas et al. (2006). For ITS region amplification, ITS1-ITS4 primers were used for PCR amplifications (White et al., 1990). ITS sequences of outgroups and new species-related taxa were downloaded from the gene bank (Table 1). All sequences obtained were aligned with the Bioedit 7.2.6.1 program (Hall, 1999). The PAUP 4.0b10 program was used for parsimony analysis and TrNef+I was found to be the most suitable model in AIC (Swofford 2002). Moreover, retention (RI), consistency (CI), and homoplasy (HI) indices were calculated. We

**Table 1.** Locality and Genebank numbers of *Centaurea* and its related taxa.

Section	Taxa	Locality	Genebank Number ITS
<i>Pseudoseridia</i>	<i>Centaurea cheirolopha</i> (Fenzl) Wagenitz		DQ319101
<i>Pseudoseridia</i>	<i>Centaurea lycopifolia</i> Boiss. & Kotschy ex Boiss. & Kotschy		DQ319130
<i>Pseudoseridia</i>	<i>Centaurea hermanni</i> F.Herm.		DQ319118
	<i>Centaurea odyssei</i> Wagenitz		DQ319140
	<i>Centaurea xylobasis</i> Rech.f.		DQ319176
	<i>Centaurea kirmacii</i> Uysal & Armağan	Türkiye; Muğla: Kavaklıdere, Menteşe village, 3.5 km Southern of Gökçukurplateu, above limestones, 1620 m, 27 vii 2017, Armağan 7797	OQ746369 (In this study)
<i>Cheirolepis</i>	<i>Centaurea drabifolia</i> Sm.		DQ319111
<i>Cheirolepis</i>	<i>Centaurea deflexa</i> Wagenitz		DQ319105
<i>Microlophus</i>	<i>Centaurea thracica</i> (Janka) Janka ex Gugler		DQ319171
	<i>Centaurea kotschyi</i> (Boiss. & Heldr.) Hayek		DQ319127
	<i>Centaurea cheirolepidoides</i> Wagenitz		DQ319100
	<i>Centaurea ensiformis</i> P.H.Davis		DQ319112
	<i>Centaurea paphlagonica</i> (Bornm.) Wagenitz		DQ319142
	<i>Centaurea nallihanense</i> Uysal & Hamzaoğlu		KX158188
	<i>Centaurea isaurica</i> Hub.-Mor.		DQ319124
	<i>Centaurea derderiifolia</i> Wagenitz		Q319106
	<i>Centaurea nivea</i> (Bornm.) Wagenitz		DQ319139
	<i>Centaurea cankiriensis</i> A.Duran & H.Duman		DQ319096
<i>Ptosimopappus</i>	<i>Centaurea ptosimopappa</i> Hayek		DQ319152
<i>Ptosimopappus</i>	<i>Centaurea ptosimopappoides</i> Wagenitz		DQ319153
	<i>Centaurea turhanii</i> Uysal & Hamzaoğlu	Türkiye, Kayseri, Akkışla, between Ortaköy and Seydinali villages, 1460 m a.s.l., gypseous steppe slopes, 28 vi 2019, Hamzaoğlu 7600 (KNYA!).	OQ746370 (In this study)
<i>Cheirolepis</i>	<i>Centaurea saligna</i> (K.Koch) Wagenitz		DQ319159

Table 1. (Continued).

	<i>Centaurea leptophylla</i> (K.Koch) Tchich.		AM087154 AM087155
<b>Microlophus</b>	<i>Centaurea behen</i> L.		AY826250
<b>Microlophus</b>	<i>Centaurea rigida</i> Banks & Sol.		DQ319158
<b>Microlophus</b>	<i>Centaurea polypodiifolia</i> Boiss.		DQ319148
	<i>Centaurea antitauri</i> Hayek		DQ319084
	<i>Centaurea kirikkalensis</i> Özbek	Türkiye; Kırıkkale: Delice, above Baraklı village, <i>Quercus pubescens</i> Willd. clearings and steppe, 1100–1150 m, a.s.l., 03 vi 2016, Hamzaoğlu 7239	OQ778863 (In this study)
<b>Rhizocalathium</b>	<i>Centaurea rhizantha</i> C.A.Mey.		DQ319157
<b>Rhizocalathium</b>	<i>Centaurea rhizocalathium</i> (K.Koch) Tchich.	Türkiye; Artvin: Yusufeli-Çevrelüköyüarası, İspir yol ayrımından 1 km, yol kenarı eğimli taşlık kayalık alanlar, 06 vi 2020, T.Uysal 4062 (KNYA!).	OQ778864 (In this study)
<b>Rhizocalathium</b>	<i>Centaurea armena</i> Boiss.		DQ319085
Outgroup	<i>Rhaponticoides pythiae</i> (Azn. & Bornm.) M.V.Agab. & Greuter		ON694102

used MrBayes 3.2 (Ronquist et al., 2012) to perform the Bayesian phylogenetic analyses. In the Bayesian analyses, random starting trees were used, which were run for  $7.1 \times 10^4$  generations, comprising 2 independent runs that consisted of four metropolis-coupled chains. Tracer version 1.7 software was used to analyze the trace files created by the Bayesian Markov chain Monte Carlo studies (Rambaut et al., 2018) and, after checking them for convergence, the first 1000 samples (20%) were discarded as burn-in. FigTree v1.4.0 software (Rambaut, 2006) was used as the graphic viewer of the phylogenetic tree.

### 3. Results

***Centaurea turhanii*** Uysal & Hamzaoğlu **sp. nova** (Figure 1; Table 2)

**Type:**— Türkiye, Kayseri, Akkışla, between Ortaköy and Seydinali villages, 1460 m a.s.l., gypseous steppe slopes, 28 vi 2019, Hamzaoğlu 7600 (**holo.** GAZI, **iso.** ANK, GAZI, HUB, KNYA).

**Diagnosis:** Although the new species is not morphologically closely related to any known species in the genus, diagnosis will be made in the focus of these two species, since *C. ptosimopappa* Hayek is located in the closest position in the phylogenetic tree. In addition, the species *C. leptophylla* (K.Koch) Tchich., and *C. malatyensis* which were thought to show a similarity at first glance in terms of habitat and habitat form as well as appendage structure, were also compared with the new species. Therefore, the new species could be related to *C. leptophylla*, *C. malatyensis*, and *C. ptosimopappa* but it is different from them by having a shorter stem (not more

than 8 cm long), floccose-tomentose leaves (not glabrous or tomentose to villous), ovate-oblong involucre (not narrowly ovoid or obovoid-oblong), alone with ciliated middle phyllaries (not all ciliated), shorter flowers (not more than 12 mm long), more large with oblong-elliptic achenes (not ovate-oblong and not less than 5 mm long) and nondeciduous simple scabrous pappus (not deciduous, biserial and plumose or barbellate).

#### **Description:**

Perennial herbs with woody rootstock, sterile shoots usually present. Stem decumbent, 3–8 cm long, densely grey floccose-tomentose, terete, simple or with a few branches in upper part, 0.9–1.5 mm diameter at base. Leaves densely grey floccose-tomentose, elliptic-oblancheolate, attenuate at base, long petiolate, apex acute, with a cartilaginous mucro; lower and median entire, denticulate or with few coarse teeth or lobes, 5–9 × 0.5–1.5 cm; upper linear-elliptic to linear-oblancheolate, usually entire. Capitula solitary at end of branches. Involucre oblong-ovate, 14–18 × 8–12 mm. Phyllaries with 5–6 series, sparsely floccose-tomentose; outer and median ones ovate-triangular; inner ones linear-lanceolate. Appendages small, usually a simple spine, mostly very deciduous, spinule simple or lanceolate, rarely ciliate, sparsely floccose-tomentose, patent or reflexed; outer linear, simple, without cilia, only with a terminal spinule; median ones linear-lanceolate, 1–2 cilia (0.5–1.5 mm) on each side, ending in a 2–3.5 mm spinule. Flowers bright yellow in vivo, pale yellow in sicco; central florets hermaphroditic, 23–25 mm long, corolla 17–19 mm long, 5-lobed, lobes 4–6 mm long, with brownish nerves, anther tube straw-colored; peripheral florets sterile, finely




Figure 1. Holotype of *Centaurea turhanii* (A); Habit (B); Capitula (C); Achene (D).

Table 2. Diagnostic characters of *Centaurea turhanii* sp. nov. and related species, *C. leptophylla*, *C. ptosimopappa*, *C. malatyensis* and *C. cheirolopha*.

→Taxa ↓Characters	<i>Centaurea turhanii</i>	<i>C. leptophylla</i> (the measurements base on both our samples (O.Tugay 5172 & T.Uysal) and the type specimen (GOET; Barcoding number 001277)	<i>C. ptosimopappa</i>	<i>C. malatyensis</i>	<i>C. cheirolopha</i>
Stem	3-8 cm long, decumbent, densely grey floccose-tomentose, terete, simple or with a few branches in upper part, 0.9-1.5 mm diameter at base	up to 25 cm, erect or ascending, simple	up to 180 cm, erect and branched from the base	7-15 cm, decumbent, simple or with few branches in the lower part and with 1(-2) capitula	Up to 25 cm, erect or ascending, simple or with few branches.

Table 2. (Continued).

<p><b>Leaves</b></p>	<p>Basal leaves soft, densely <b>grey floccose-tomentose</b>, elliptic-ob lanceolate, long petiole; lower and median entire, denticulate or with few coarse teeth or lobes, 5–15 mm broad; upper similar, usually entire.</p>	<p>soft, <b>glabrous</b>, linear-lanceolate, shortly petiole, 2–4 mm broad; upper similar, entire.</p>	<p>firm and leathery, lanceolate-spathulate to obovate, upper similar, entire.</p>	<p>Basal leaves short petiole (20–25 mm), oblanceolate, 5–6 mm wide, entire, tomentose to villous, lateral nerves indistinct, median and upper leaves sessile, linear to lanceolate; median leaves 4–6 × 0.5–0.7 cm; upper leaves 3–4.5 × 0.4–0.6 cm with entire margin</p>	<p>Leaves bicolored, green, and glabrescent above, grey-tomentose below, of very variable form, basal and lower usually lyrate with large lanceolate or hastate terminal segment and 1–3 pairs of lateral segments or undivided, lanceolate, entire or denticulate; median and upper leaves lanceolate to linear-lanceolate, sometimes narrowly decurrent.</p>
<p><b>Involucre</b></p>	<p><b>Ovate-oblong</b>, 14–18 × 8–12 mm; Phyllaries with 5–6 series, sparsely floccose-tomentose; outer and median ovate-triangular; inner linear-lanceolate.</p>	<p>narrowly obovoid-oblong, 12–15 × 9–12 mm; Phyllaries with 5–6 series, glabrous; outer and median ovate-triangular; inner linear-lanceolate.</p>	<p>narrowly ovoid 18–25 × 8–16 mm; Phyllaries very numerous, glabrous.</p>	<p>Ovoid, 37–47 × 12–17 mm; villous in upper parts, greenish-yellow</p>	<p>Ovoid, 15–20 × 9–12(–15) mm,</p>
<p><b>Appendage</b></p>	<p>small, usually a simple spine, mostly very deciduous, spinule simple or lanceolate, rarely ciliate, sparsely floccose-tomentose, patent or reflexed; outer linear, simple, without cilia, only with a terminal spinule; median linear-lanceolate, <b>1–2 cilia (0.5–1.5 mm) on each side</b>, ending in a 2–3.5 mm spinule</p>	<p>small, with a simple spine, mostly very deciduous, spinule simple or lanceolate, without cilia, glabrous, patent or reflexed; outer linear, simple, without cilia, only with a terminal spinule; median linear-lanceolate, <b>without cilia</b>, ending in a 2.5–3.5 mm spinule</p>	<p>a minute, deciduous, spinule simple and too short, <b>without cilia</b>, glabrous, patent, or reflexed; all similar, ending in a 0.3–0.5 mm spinule.</p>	<p>triangular, not concealing basal part of phyllaries, not decurrent, entire in the outer and the median phyllaries with a spinule 2–3 mm long, irregularly lacerate in the inner phyllaries ending in a spinule 0.5–1.5 mm long.</p>	<p>Appendages small, brown, broadly triangular, spreading or reflexed, with 5–7 digitate 1–5 mm spinules.</p>
<p><b>Flowers</b></p>	<p>yellow, not radiant, <b>9–12 mm</b> long, with 5 linear-filiform 4–6 mm lobes; anther tubes yellow</p>	<p>yellow, not radiant, <b>18–20 mm</b> long, with 5 linear-filiform 5–6 mm lobes; anther tubes yellow.</p>	<p>yellow, not radiant, <b>20–25 mm long</b>, with 5 linear-filiform 5–6 mm lobes; anther tubes yellow.</p>	<p>yellow, marginal not radiant, ca. 35 mm long, corolla tube glabrous, 18–22 mm long, lobes 6–8 mm, linear with brown stripes along corolla</p>	<p>Flowers yellow</p>

Table 2. (Continued).

<b>Achene</b>	5–6(–7) × 1.8–2.3 mm, <b>oblong-elliptic</b> , brownish, glabrous, notched and rounded at base.	4.5–5 × 2.2–2.5 mm, narrowly ovoid- <b>oblong</b> , creamish-brown in mature, superficially with sparsely pilose hairy.	4–5 × 2 mm, narrowly ovoid to oblong, brown, with sparsely pilose hairy.	6–6.5 × 3.5–4.5 mm, obovoid with striations, light brown, covered with adpressed whitish hairs;	4–5.5 mm;
<b>Pappus</b>	<b>simple</b> , scabrous, brownish to purplish, 2-3(–4) mm long, <b>not deciduous</b>	<b>double</b> , scabrous, outer series whitish-cream, 8–10 mm, barbellate, inner series 0.5 mm, similar to ascale, <b>deciduous</b> .	<b>Simple</b> , scabrous, creamish-brown, c. 4 mm, deciduous.	<b>simple</b> , 18–22 mm, plumose, brownish	0.5-1.5 mm, inner row not distinctly different.

dissected, not radiant, nearly equal to central ones, 15–17 mm long, 4–5-lobed, lobes 4–6 mm long. Achenes 5–6 × 1.8–2.3 mm, oblong-elliptic, brownish, glabrous, notched and rounded at base. Pappus scabrous, brownish to purplish, simple, 2-3(–4) mm long, persistent.

**Phenology:**—The new species blooms in June and starts fruiting in mid-July.

**Group B (Volume V, Flora of Turkey (Wagenitz, 1975)**

- 1. Flowers pink or purple
- 1. Flowers yellow
- 3. All leaves linear-lanceolate to linear or elliptic-oblancheolate, entire
- 4. Leaves linear lanceolate, glabrous 172. *leptophylla*
- 4. Leaves elliptic-oblancheolate, densely tomentose or grey floccose-tomentose
- 5. Pappus plumose, 18–22 mm 172.a. *malatyensis*
- 5. Pappus scabrous, 2–4 mm 172.b. *turhanii*
- 3. All leaves pinnatifid to lyrate

**Etymology:** The species is named in honor of our precious friend and scientist Prof. Dr. Turhan ÇETİN (Ankara, Türkiye), who has participated in many field studies with us and made geographical observations and created maps. He is very fruitful a scientist who has studies on Türkiye's natural environment and botanical tourism.

**Distribution, habitat, and ecology:** *Centaurea turhanii* grows on gypseous steppes between Ortaköy and Seydinali villages in the Akkişla district of Kayseri Province at altitudes of 1370–1590 m a.s.l. (Figure 2), and it is associated with *Gypsophila turcica* Hamzaoğlu, *Thesium stelleroides* Jaub. & Spach, *Thymus cappadocicus* Boiss., *Isatis glauca* Willd. ex DC. subsp. *sivasica* (P.H.Davis) Yild., *Matthiola anchoniifolia* Hub.-Mor., *Ferula szowitsiana* DC., and *Paracaryum racemosum* Britten. The area where the species grows is not threatened other than intensive grazing, and it is part of the Irano-Turanian floristic region.

Due to the lack of detailed observations and studies on the distribution area of the species and the number of individuals for several years, no suggestions were made regarding the threat category and protection status.

**Achene structure:** Achenes of *Centaurea turhanii* are glabrous, oblong-elliptical, 5.9–6.3 × 2.7–2.9 mm, with ornamentation irregular sulcate. Cells are distinct and cell boundaries are thin (Figure 3). Achenes of *C. ptosimopappa* are sparsely pilose, ovoid to oblong, 5.24 × 2.05 mm, with ornamentation irregular sulcate. Ripe achene lacks the pappus (deciduous). Cells and cell walls are distinct (Figure 3).




Figure 2. Distribution map of *C. turhanii* and related taxa.




Figure 3. Scanning electron micrographs of achene in *Centaurea turhanii* (A–D) and *C. ptosimoppappa* (E–H).




Figure 4. Mitotic metaphase chromosomes, idiograms and karyograms of *Centaurea turhanii* (A–C) and *C. ptosimoppappa* (D–F).

**Table 3.** The chromosome features of *Centaurea* taxa.


Taxa	2n	R SC-LC (µm)	R- LC / SC	p (µm) mean (±SD)	q (µm) mean (±SD)	CL (µm) mean (±SD)	TCL (µm)	CI mean (±SD)	KF
<i>C. turhanii</i>	18	3.05– 4.92	1.613	1.38 (±0.43)	2.40 (±0.39)	3.78 (±0.73)	33.995	36 (±0.06)	6m + 10sm + 2st
<i>C. ptosimopappa</i>	18	2.40– 4.14	1.725	1.24 (±0.30)	1.80 (±0.36)	3.05 (±0.59)	27.415	41 (±0.05)	10m + 8sm
<i>Centaurea ptosimopappa</i> (a) (Koçyiğit& Bona 2013)	18								3m+2msm+4sm
<i>Centaurea ptosimopappa</i> (b) (Koçyiğit& Bona 2013)	18								4m+5msm
<i>C. leptophylla</i> (Uysal et al. 2015)	36	1.36– 2.43	1.78	0.67 (±0.15)	1.07 (±0.22)	1.75 (±0.33)	31.5	38 (±0.05)	2m+7sm

R: Range, SC: Shortest chromosome length, LC: Longest chromosome length, p: Mean length of the long short arm, q: Mean length of the long arm, CL: Mean chromosome length, TCL: Total haploid complement length, CI: Mean centromeric index, SD: Standard deviation, KF: Karyotype formula, m: Metacentric, sm: Submetacentric, st: Subtelocentric.

**Table 4.** The karyotype indices of *Centaurea* taxa

Taxa	A <sub>1</sub>	A <sub>2</sub>	CV <sub>CL</sub>	CV <sub>CI</sub>	AI	M <sub>CA</sub>
<i>C. turhanii</i>	0.429	0.194	19.418	15.968	3.101	26.98
<i>C. ptosimopappa</i>	0.302	0.193	19.269	12.58	2.424	18.42
<i>C. leptophylla</i> (Uysal et al 2015)	0.36	0.19	18.95	12.99	2.46	22.99

A<sub>1</sub>: Intrachromosomal asymmetry index, A<sub>2</sub>: Interchromosomal asymmetry index, CV<sub>CL</sub>: Coefficient of variation of chromosome length, CV<sub>CI</sub>: Coefficient of variation of centromeric index, AI: Karyotype asymmetry index, M<sub>CA</sub>: Mean centromeric asymmetry.



**Figure 5.** Majority rule consensus tree obtained from Parsimony and Bayesian analysis of the ITS data for *C. turhanii* and related taxa (tree length: 608, CI: 0.732, RI: 0.762, HI: 0.268). Values in the branches are shown as maximum Bayesian posterior probabilities/Parsimony bootstrap.



**Karyology:** According to our counts, *Centaurea turhanii* is diploid with chromosome number  $2n = 18$ . The karyotype formula of the species is  $2n = 6m + 10sm + 2st$  (Figure 4). According to the classification based on the length of the chromosomes (Lima De Faria 1980), the new species has four pairs of medium and five pairs of small chromosomes and chromosomes get shorter gradually from medium to small (Table 3). When we compare our results with the information from the literature, although there is agreement in terms of the chromosome number of *C. ptosimopappa*, there are significant differences in terms of karyomorphology. In the study conducted by Koçyiğit and Bona (2013), different karyotype formulas of two different populations of *C. ptosimopappa* were specified. Interestingly, there is no match between the related study and the karyotype formulas determined in our study. (Figure 4 and Table 3). Chromosomal features of the new species and related taxa are given in Tables 3 and 4.

**Phylogeny:** Phylogenetic analyses included 32 taxa, one of which was an outgroup. The aligned data matrix consists of 608 base pairs long and 533 of them are constant. While the informative character number for parsimony was 28 and the uninformative was 47. The tree created by the PAUP program has reliable tree scores [consistency index (CI: 0.732), retention index (RI: 0.762), and homoplasy (HI: 0.268)]. Parsimony analysis was executed with the TrNef+I model according to chosen AIC criteria. Trees revealed by parsimony and Bayesian analysis formed similar topologies for the studied *Centaurea* taxa (Figure 5). Therefore, a common tree was created by combining the analysis. In the combined tree based on ITS regions, the new species is in a different clade from all studied *Centaurea* taxa in terms of its phylogenetic history. Indeed, we concluded that the new species is located phylogenetically differently from the taxa belonging to the sections analyzed.

#### 4. Discussion

*Centaurea turhanii* has been characterized by achene surface micromorphology, karyomorphology, nrDNA ITS, and morphological studies. *Centaurea turhanii* which was published as a new species to science is a unique species both morphologically and phylogenetically, it does not have any close relatives in the genus in terms of the characteristics given. Previously, the involucre bract and appendages have played a vital role morphologically in the taxonomy of the genus (Wagenitz, 1975), and the subgroups were classified according to these characters. However, in the ongoing studies, it has been pointed out that the classification based on phyllaries and appendages is unreliable in *Centaurea*. In particular, the analyses based on molecular studies suggested that there is a mismatch between morphology and molecular data and that the

reason for this is rapid evolution due to hybridization (Garcia-Jacas et al., 2006; Hilpold et al., 2014). Similarly, it is declared that sectional classification had difficulties in establishing natural and clear division based on bract appendices as depended on the entanglement of sections, as previously suggested by Ertuğrul et al. (2004) and Hilpold et al. (2014). The virtual absence of congruence between morphology and molecular data suggests that the morphological characters that have been used taxonomically are not reliable with respect to the true relationships. Some morphological traits may have developed convergently in several cases– a high plasticity of morphological characters is also likely. The development of equal or similar morphological character states may have partly been due to an adaptive response to similar ecological conditions, for example, the development of small divided tomentose leaves as a response to dry conditions (cf. Larcher, 2003), or of spiny bract appendages as a response to herbivores (cf. Cooper and Owen-Smith, 1986; Hilpold et al., 2014). Because of these problems, we will focus on the groups and species to which the new species is relative according to the information from the molecular and karyological data, rather than morphology.

In terms of involucre bracts, the new species exhibits an intermediate state on the point of main types of appendages. While the outer phyllaries are in the form of a simple spinule, the appendages of some middle phyllaries irregularly contain one or two pairs of cilia at the base or mostly simple spinule. The inner phyllaries are in the form of a narrowly triangular simple membranous spine. Based on this morphology, especially in group B, in which the appendage has been reduced to a terminal spiny or spinule (Wagenitz, 1975). One of the sections defined in Group B is *Microlophus* and the section is represented by six species in the Flora of Turkey. Another section thought to be related according to phylogenetical analyses is *Ptosimopappus* and this section includes only two endemic species. The known species of these sections in the mentioned groups consist of taller plants with long branches and erect stems. Even if it is not completely morphologically, the new species can be more associated with the taxa of the *Ptosimopappus* too, considering the mainly achene and pappus structures. However, a very short and creeping stem is not a usual for the species of neither A nor B groups. Besides with this general comparison, the new species display morphologically an interesting similarity with *C. leptophylla* which are a local endemic species spreading in Çoruh Valley of Artvin Province. Although the two species are similar in terms of growing on rocky slopes, creeping habitat form and especially appendages, they differ in terms of leaf structure, indumentum and pappus characteristics. In addition, the new species is similar to *Centaurea malatyiensis*. However, it is easily distinguished

from the *C. malatyensis* by its smaller capitula and very short and scabrous pappus. Moreover, the two species (new species and *C. malatyensis*) are different in their ecology; the new species is distributed on gypseous steppes while the other species (*C. malatyensis*) prefers serpentine cliffs. The new species clearly looks different when compared to *C. cheirolopha*. Not only *C. turhanii* is different in terms of leaf hairy and color (not bicolor) but also it has short creeping stem (not longer than 10 cm and not erect-ascending stem). Additionally, while *C. turhanii* is rarely including irregularly one or two pair cilia at the base of medium phyllaries, *C. cheirolopha* has 5–7 pair digitate cilia in all phyllaries (not all phyllaries have several pairs of cilia). Lastly, it includes longer and scabrous pappus hairs (not 0.5–1.5 mm and not plumose) (Table 2).

So far, it has been reported that achene surface morphology is effective in the differentiation of taxa belonging to the genus *Centaurea* (Uysal et al., 2016; Güzel, 2022). The cell boundaries of *C. malatyensis* are relatively expanded. The cell boundaries of *C. leptophylla*, *C. ptosimopappa*, and *C. cheirolopha* species are thinner than *C. malatyensis*. However, even the boundaries of these species are thicker compared to the new species. It is seen that the new species differs from *C. ptosimopappa*, *C. malatyensis*, *C. leptophylla*, and *C. cheirolopha* in terms of achene shape, surface ornamentation and hairy status (Kültür et al., 2016; Aydın et al., 2019; Güzel 2022).

The chromosome number determined had been reported before for many *Centaurea* species in several studies (Georgiadis and Phitos, 1976; Phitos and Damboldt, 1976; Romaschenko et al., 2004; Uysal, 2008, 2009; Uysal et al., 2015, 2017). Additionally, the basic chromosome numbers of some taxa belonging to *Microlophus* were reported as  $x = 8$  and  $x = 17$  (Garcia-Jacas et al., 1998; Romashchenko et al., 2004; Uysal et al., 2009; Tasar et al., 2018). These results disagree with number of the new species. The basic chromosome number of the *C. thracica* (Janka) Janka ex Gugler was reported as  $x = 9$  (Damboldt and Matthäs, 1975; Constantinidis et al., 2002). However, the taxonomic position of this species is controversial according to phylogeny studies (Garcia-Jacas et al., 2006). Based on the information we obtained from the literature, with exceptions, we reconfirmed the  $x = 9$  basic chromosome number commonly seen in the genus *Centaurea*. According to our karyological comparisons among *C. turhanii*, *C. ptosimopappa*, and *C. leptophylla*, the new species has distinctive karyomorphological features (Karyotype formula and asymmetry indices; Tables 3 and

4) and is clearly different karyologically from its relatives. Unlike of *C. leptophylla*, the new species is a diploid like *C. ptosimopappa*. The karyotype formula is unique and consist of mostly submetacentric chromosomes as well as subtelocentric. Subtelocentric chromosomes are not seen in related taxa.

Romaschenko et al. (2004) reported the chromosome number of *C. cheirolopha* as  $2n = 2x = 18$  (diploid). *C. cheirolopha* is the mostly similar species in terms of karyomorphology with the new species. However, the sixth and ninth chromosome pairs of *C. cheirolopha* are metacentric. In particular, *C. cheirolopha* differs from the new species because it has metacentric chromosome pairs.

As related to chromosomal counts and analyses of *C. ptosimopappa*, there is a minor conflict among present and previous reports (Tables 3 and 4) Even so, the karyotype formulas of this taxon are not hold related to the new species.

According to the ITS based phylogenetic tree; the new species is taken place relatively far from the better resolved clade containing *Microlophus* species (PP: 0.69) and in the immediate near of the clade containing *Ptosimopappus* species (PP: 1/BS: 94), which can be considered as a polytomic clade. For this reason, it can be said that the species does not display high closeness to any members of mentioned groups but partly more closely related to species of the sections *Microlophus* and *Ptosimopappus*. However, based on the current data, it is thought that it would not be correct to evaluate the species in one of these two sections, and it may be considered to determine a common new section for *C. leptophylla*, *C. malatyensis* and *C. turhanii* species in a future study.

#### Additional specimens examined

##### *Centaurea ptosimopappa* Hayek

C6 Osmaniye, Zorkumroad, 1050 m, roadside, 05 July 2003, A. Duran 6306 & M. Sağıroğlu (KNYA!); C6 Osmaniye, between Zorkum, 300–400 m, roadside, 12 September 2003, A. Duran 6344 & M. Sağıroğlu (KNYA!).

##### *Centaurea leptophylla* (K.Koch) Tchich.

Artvin; Yusufeli, Sarıgöl girişi 790 m, 08 viii 2007, O. Tugay 5172 & T.Uysal (KNYA!).

#### Acknowledgements

We are grateful to the foundation TÜBİTAK (the project number 117Z222) and S.U. BAP (the project number 19401172) because of their financial support.

## References

- Armağan M, Uysal T (2018). *Centaurea kirmacii* (Asteraceae), a new species from southwestern Anatolia, Turkey. *Phytotaxa* 362 (2): 233-238. DOI: <https://doi.org/10.11646/phytotaxa.362.2.10>
- Aydın R, Dural H, Yılmaz Çıtak B (2019). Investigation of morphological, anatomical and micromorphological characteristics of *Centaurea cheirolopha* (Fenzl) Wagenitz and *Centaurea lycopifolia* Boiss. & Kotschy (Asteraceae) taxa. *Erzincan University Journal of Science and Technology* 12 (3): 1341-1355. DOI: [10.18185/erzifbed.480745](https://doi.org/10.18185/erzifbed.480745)
- Bancheva S, Kaya Z, Binzet R (2014). *Centaurea aytugiana* (Asteraceae), a new species from north Anatolia, Turkey. *Novon* 23 (2): 133-138. DOI: <https://doi.org/10.3417/2011026>
- Bancheva S, Kaya Z (2015). *Centaurea raimondoi*, a new species from Asteraceae. *Flora Mediterranea* 25: 305-310. DOI: <https://doi.org/10.7320/FIMedit25.305>
- Behçet L, İlçim A, Yapar Y (2017). *Centaurea bingöelensis* (Asteraceae), a new species from Turkey. *Turkish Journal of Botany* 41 (2): 180-188. DOI: <https://doi.org/10.3906/bot-1512-14>
- Bona M (2015). *Centaurea goksivriensis* (Asteraceae), a new species from Turkey. *Phytotaxa* 203 (1): 63-68. DOI: <https://doi.org/10.11646/phytotaxa.203.1.6>
- Bona M (2016). *Centaurea amanosensis* (Asteraceae), a new species from Turkey. *Plant Biosystems* 150 (5): 1083-1086. DOI: <https://doi.org/10.1080/11263504.2014.1001465>
- Constantinidis T, Bareka EP, Kamari G (2002). Karyotaxonomy of Greek serpentine angiosperms. *Botanical Journal of the Linnean Society* 139 (1): 109-124. <https://doi.org/10.1046/j.1095-8339.2002.00044.x>
- Cooper SM, Owen-Smith N (1986). Effects of plant spinescence on large mammalian herbivores. *Oecologia* 68: 446-455.
- Cullings KW (1992). Design and testing of a plant-specific PCR primer for ecological and evolutionary studies. *Molecular Ecology* 1 (4): 233-240. <https://doi.org/10.1111/j.1365-294X.1992.tb00182.x>
- Damboldt O, Matthäs C (1975). Chromosome numbers of some Mediterranean and C. European *Centaurea* species (Asteraceae). *Plant Systematics and Evolution* 123 (2): 107-115.
- Doyle JJ, Doyle JL (1987). A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemical Bulletin*, Botanical Society of America 19: 11-15.
- Duman H, Uzunhisarcıklı ME, Noyan Bahadır Y (2021). A new species of *Centaurea* (Asteraceae) from northern Turkey. *Nordic Journal of Botany* 39 (10): 1-8. DOI: <https://doi.org/10.1111/njb.03244>
- Dostál J (1976). *Centaurea* L. In: Tutin TG, Heywood VH, Burges NA, Moore DM, Valentine DH, Walters SM, Webb DA (editors). *Flora Europaea*, Vol. 4, Cambridge: Cambridge University Press, pp. 254-301.
- Ertuğrul K, Uysal T, Garcia-Jacas N, Susanna A, Garnatje T (2004). The systematic position of *Centaurea ensiformis* and *Centaurea isaaurica* from Turkey and the evolution of some characters in *Centaurea*. *Israel Journal of Botany* 52: 257-263.
- Garcia-Jacas N, Susanna A, Mozaffarian V (1998). New chromosome counts in the subtribe Centaureinae (Asteraceae, Cardueae) from West Asia. III. *Botanical Journal of the Linnean Society* 128 (4): 413-422. DOI: [10.1111/j.1095-8339.1998.tb02129.x](https://doi.org/10.1111/j.1095-8339.1998.tb02129.x)
- Garcia-Jacas N, Susanna A, Mozaffarian V, Ilarslan R (2000). The natural delimitation of *Centaurea* (Asteraceae: Cardueae): ITS sequence analysis of the *Centaurea jacea* group. *Plant Systematics and Evolution* 223: 185-199.
- García-Jacas N, Susanna A, Garnatje T, Vilatersana R (2001). Generic delimitation and phylogeny of the subtribe Centaureinae (Asteraceae): a combined nuclear and chloroplast DNA analysis. *Annals of Botany* 87 (4): 503-515. <http://dx.doi.org/10.1006/anbo.2000.1364>
- García-Jacas N, Uysal T, Romaschenko K, Suárez-Santiago VN, Ertuğrul K et al. (2006). *Centaurea* revisited: A molecular survey of the *Jacea* group. *Annals of Botany* 98: 741-753. <http://dx.doi.org/10.1093/aob/mcl157>
- Georgiadis T, Phitos D. (1976). Contribution à l'étude cytotonomique du genre *Centaurea* L. (sectio Acrolophus (Cass.) DC.) en Grèce [Contribution to the cytotonomic study of the genus *Centaurea* L. (section Acrolophus (Cass.) DC.) in Greece]. *Biologie et Écologie Méditerranéenne* 3: 13-16.
- Greuter W (2003) The Euro+Med treatment of Cardueae (Compositae)—generic concepts and required new names. *Willdenowia* 33: 49-61. <http://dx.doi.org/10.3372/wi.33.33104>.
- Greuter W, Raus T (2009). Med-ChecklistNotulae, 28. *Willdenowia* 39 (2): 335-345. DOI: <https://doi.org/10.3372/wi.39.39211>
- Güzel Y (2022). *Centaurea ptosimopappa* var. *elegans* Güzel. A New *Centaurea* L. (Asteraceae) taxon from Türkiye with taxonomical notes on sect. *Ptosimopappus* and with lectotypification. *ANADOLU Journal of Aegean Agricultural Research Institute* 32 (2): 287-300. DOI: [10.18615/anadolu.1216309](https://doi.org/10.18615/anadolu.1216309)
- Hall TA (1999) BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT/2K/XP. *NucleicAcidsSymp. Ser.* 41: 95-98 (1999).
- Hamzaoğlu E, Koç M (2020). A new record for the flora of Turkey, *Centaurea gulissashwillii* (sect. *Centaurea*, Asteraceae). *Türler ve Habitatlar* 1 (2): 106-113.
- Hilpold A, Vilatersana R, Susanna A, Meseguer Andrea S, Boršić I et al (2014). Phylogeny of the *Centaurea* group (*Centaurea*, Compositae): geography is a better predictor than morphology. *Molecular Phylogenetics and Evolution* 77: 195-215. <http://dx.doi.org/10.1016/j.ympev.2014.04.022>
- Kashi AM, Tahermanesh K, Chaichian S, Joghataei MT, Moradi F et al (2014). How to Prepare Biological Samples and Live Tissues for Scanning Electron Microscopy (SEM). *Galen Medical Journal* 3 (2): 63-80. DOI: [10.31661/gmj.v3i2.267](https://doi.org/10.31661/gmj.v3i2.267)
- Koçyiğit M, Bona M (2013). Chromosome numbers of five Turkish *Centaurea* L. (Asteraceae) species. *Plant Biosystems*. 147 (4): 970-978. <http://dx.doi.org/10.1080/11263504.2013.789454>
- Köse YB, Alan S (2013). *Centaurea baseri* (Compositae), a new species from Turkey. *Phytotaxa* 117 (1): 23-29. DOI: <http://dx.doi.org/10.11646/phytotaxa.117.1.3>

- Kültür Ş, Bona M, Nath EÖ (2016). A new species of *Centaurea* (Asteraceae) from East Anatolia, Turkey. *Phytotaxa*. 247 (1): 85-91. DOI: <https://doi.org/10.11646/phytotaxa.247.1.6>
- Larcher W. 2003. *Physiological Plant Ecology*, fourth ed. Springer, Berlin.
- Lima De, Faria A (1980). Classification of genes, rearrangements and chromosomes according to the chromosome field. *Hereditas* 93: 1-46.
- Negaresh K, Kaya Z, Rahiminejad MR (2015). *Centaurea semikoviiana* (Asteraceae, Cardueae) a new species from central Anatolia, Turkey. *Annales Botanici Fennici* 52 (5/6): 321–327. DOI: <https://doi.org/10.5735/085.052.0507>
- Negaresh K, Rahiminejad MR (2018). A revision of *Centaurea* sect. *Cynaroides* (Asteraceae, Cardueae–Centaureinae). *Phytotaxa* 363 (1): 1-131. DOI: <https://doi.org/10.11646/phytotaxa.363.1.1>
- Negaresh K (2019). A contribution to the taxonomy of the genus *Rhaponticoides* (Asteraceae, Cardueae). *Phytotaxa* 422 (1): 9–32.
- Özbek UM (2021). *Centaurea kirikkalensis* (Asteraceae), a new species from Turkey. *Nord J Bot* 39 (7): 1-9. DOI: <https://doi.org/10.1111/njb.03235>
- Paszko B (2006). A critical review and a new proposal of karyotype asymmetry indices. *Plant Systematics and Evolution* 258: 39-48. <https://doi.org/10.1007/s00606-005-0389-2>
- Peruzzi L, Eroğlu HE (2013). Karyotype asymmetry: again, how to measure and what to measure?. *Comparative Cytogenetics* 7 (1): 1-9. doi: 10.3897/CompCytogen.v7i1.4431
- Phitos D, Damboldt J (1976). Ein Beitrag zur Kenntnis der Gattung *Centaurea* L. in Griechenland [A contribution to the knowledge of the genus *Centaurea* L. in Greece]. *Veröffentlichungen des Geobotanischen Institutes ETH, Stiftung Rübél, Zürich* 56: 183-189.
- Pınar SM (2016). *Centaurea sintenisiana* Gand. A new record for the family Asteraceae (Compositae) from Turkey. *Yüzüncü Yıl University Journal of the Institute of Natural & Applied Sciences* 21 (2): 75-82.
- Rambaut A (2006). FigTree: Tree figure drawing tool version 1.4.0. University of Edinburgh: Institute of Evolutionary Biology, (<http://tree.bio.ed.ac.uk/software/figtree/>).
- Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA (2018). Posterior summarization in Bayesian phylogenetics using Tracer 1.7. *Systematic Biology* 67 (5): 901-904. <https://doi.org/10.1093/sysbio/syy032>.
- Romero-Zarco C (1986). A new method for estimating karyotype asymmetry. *Taxon* 35(3): 526-530. <https://doi.org/10.2307/1221906>
- Romaschenko K, Ertugrul K, Susanna A, Garcia-Jacas N, Uysal T et al (2004). New chromosome counts in the *Centaurea jacea* group (Asteraceae, Cardueae) and some related taxa. *Botanical Journal of the Linnean Society* 145 (3): 345-352. <http://dx.doi.org/10.1111/j.1095-8339.2004.00292.x>
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A et al (2012). MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61 (3): 539-542. <https://doi.org/10.1093/sysbio/sys029>
- Soltis DE, Soltis PS, Collier TG, Edgerton ML (1991). Chloroplast DNA Variation within and among Genera Of The Heuchera Group (Saxifragaceae): Evidence for chloroplast transfer and paralogy. *American Journal of Botany* 78: 1091-1112. <https://doi.org/10.2307/2444898>.
- Susanna A, García-Jacas N (2007). Tribe Cardueae Cass. In: Kadereit, J.W. & Jeffrey, C. (Eds.) *Kubitzki's The families and genera of vascular plants*, vol. 8. Springer, Berlin Heidelberg, pp. 123-147.
- Swofford DL (2002). PAUP/. *Phylogenetic Analysis Using Parsimony (and other methods)*, ver. 4.0b10. Sinauer, Sunderland.
- Şirin E, Çeçen Ö, Bozkurt M, Ertuğrul K (2019). *Centaurea uysalii* (Cyanus/Asteraceae), a new species from Turkey. *Turkish Journal of Botany* 43 (6): 809-816. DOI: <https://doi.org/10.3906/bot-1901-12>
- Şirin E, Uysal T, Bozkurt M, Ertuğrul K (2020). *Centaurea akcaadaghensis* and *C. ermenekensis* (Asteraceae), two new species from Turkey. *Mediterranean Botany* 41 (2): 173-79. DOI: <https://doi.org/10.5209/mbot.68628>
- Şirin E, Uysal T, Bozkurt M, Ertuğrul K (2022). Karyomorphological, molecular and morphological evidence for a new species, *Centaurea mengensis* (Asteraceae), from Turkey. *Nordic Journal of Botany* e03389. DOI: <https://doi.org/10.1111/njb.03389>
- Tasar N, Dogan G, Kiran Y (2018). Karyological investigation on seven *Centaurea* L. (Asteraceae) taxa from Turkey. *Cytologia* 83 (3): 317-321. DOI: 10.1508/cytologia.83.317
- Thiers B. (2023) Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/science/ih/>.
- Tzvelev NN (2001). [*Centaurea* L.] In: Bobrov, E.G. & Czerepanov, S.K. (Eds.). *Flora of USSR*. Vol. 28 (Translated from Russian: Smithsonian Institution Libraries). Amerind Publishing, Washington, D.C., pp. 529-530.
- Uysal T (2008). *Centaurea ertugruliana* (Asteraceae), a new species from Turkey. *Annales Botanici Fennici* 45: 137-140.
- Uysal T, Ertuğrul K, Susanna A, Garcia-Jacas N (2009). New chromosome counts in the genus *Centaurea* (Asteraceae) from Turkey. *Botanical Journal of the Linnean Society* 159 (2): 280-286. DOI: 10.1111/j.1095-8339.2008.00939.x
- Uysal T, Arslan E, Tugay O, Ertugrul K (2010). Determination of the relationship between some *Centaurea* species based on SDS-PAGE. *Turkish Journal of Biology* 34 (2): 125-131. DOI: <https://doi.org/10.3906/biy-0809-15>
- Uysal T. (2012). [*Centaurea* L.]. In: Güner A, Aslan S, Ekim T, Vural M, Babaç MT (edlr.). *Türkiye Bitkileri Listesi (Damarlı Bitkiler)*. Nezahat Gökyiğit Botanik Bahçesi ve Flora Araştırmaları Derneği Yayını, İstanbul, pp. 127-140.
- Uysal T, Bozkurt M, Şimşek Sezer EN, Ertuğrul K, Tugay O (2015). Karyological studies of four endemic *Centaurea* L. species. *Caryologia* 68 (4): 339–346. DOI: <https://doi.org/10.1080/00087114.2015.1109922>
- Uysal T, Hamzaoğlu E, Ertuğrul K, Bozkurt M (2016). A new species of *Centaurea* (Asteraceae) from Turkey. *Phytotaxa* 275(2): 149-158. DOI: <https://doi.org/10.11646/phytotaxa.275.2.6>

- Uysal T, Bozkurt M, Tugay O, Ertuğrul K, Şimşek Sezer EN et al. (2017a) Karyomorphology of Turkish species in *Centaurea* sections *Centaurea* and *Phalolepis* (Asteraceae) and implications for taxonomy, Plant Biosystems - An International Journal Dealing with all Aspects of Plant Biology, 151 (6): 949-964. DOI: 10.1080/11263504.2016.1218967
- Uysal T, Dural H, Tugay O (2017b). *Centaurea sakariyaensis* (Asteraceae), a new species from Turkey. Plant Biosystems 151 (1): 126-130. DOI: <https://doi.org/10.1080/11263504.2015.1108940>
- Uysal T, Hamzaoğlu E (2017). A new *Centaurea* L. (Asteraceae) species from Turkey. Plant Biosystems 151: 813-821. DOI: <https://doi.org/10.1080/11263504.2016.1211196>
- Uysal T, Bozkurt M, Hamzaoğlu E (2022). A New species from Türkiye, *Centaurea ankarica* (Cardueae, Asteraceae). Türler ve Habitatlar 3 (2): 81-93. doi.org/10.53803/turvehab.1167966
- Wagenitz G (1975). *Centaurea* L. In: Davis, P.H. (Ed.) Flora of Turkey and the East Aegean Islands, vol. 5. Edinburgh University Press, Edinburgh, pp. 465-585.
- Wagenitz G (1980). [*Centaurea* L.] In: Rechinger, K.H. (Ed.). Flora Iranica. Vol. 139b. Akademische Druck- und Verlagsanstalt, Graz, pp. 356-362.
- Wagenitz G (1986). *Centaurea* L. in South-West Asia: Patterns of distribution and diversity. Proceedings of the Royal Society Edinburgh 89 B: 11-21. <http://dx.doi.org/10.1017/s0269727000008861>
- Wagenitz G, Hellwig FH (2000). The genus *Psephellus* Cass. (Compositae, Cardueae) revisited with a broadened concept. Willdenowia 30 (1): 29-44. doi.org/10.3372/wi.30.30102
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In PCR Protocols: A Guide to Methods and Applications (eds Innis, M. A. et al.) 315-322 (Academic Press, 1990).
- Yıldırım Ş (2012). The heaven of gypsophilous phytodiversity of Turkey: Keven, Sivrihisar, Eskişehir, Turkey, 13 taxa as new. Ot Sistemik Botanik Dergisi 19 (2): 1-51.
- Yıldırım Ş (2018). Five new species, a new subspecies and a new combination and status from gypsaceous areas of Ankara province, Turkey. Ot Sistemik Botanik Dergisi 25: 1-22.
- Yüzbaşıoğlu İS, Bona M, Genç M (2015). A new species of *Centaurea* sect. *Pseudoseridia* (Asteraceae) from north-eastern Turkey. PhytoKeys 53: 27-38. DOI: <https://doi.org/10.3897/phytokeys.53.5250>