

Magnetite nanoparticles–based hydroxyl radical scavenging activity assay of antioxidants using N, N-dimethyl-*p*-phenylenediamine probe

Ziya CAN¹, Büşra KESKİN^{1,2}, Ayşem ARDA^{1*}, Erol ERÇAĞ³, Mustafa Reşat APAK^{1,4}

¹Department of Chemistry, Faculty of Engineering, İstanbul University-Cerrahpaşa, İstanbul, Turkey

²Department of Chemistry, Institute of Graduate Studies, İstanbul University-Cerrahpaşa, Turkey

³Department of Chemistry, Faculty of Arts and Sciences, Tekirdağ Namık Kemal University, Tekirdağ, Turkey

⁴Turkish Academy of Sciences (TÜBA), Ankara, Turkey

Received: 02.06.2020 • Accepted/Published Online: 27.07.2020 • Final Version: 26.10.2020

Abstract: Excessive amounts of reactive oxygen species (ROS), unless counterbalanced by antioxidants, can cause cellular damage under oxidative stress conditions; therefore, antioxidative defenses against ROS must be measured. With the development of nanotechnology, nanoparticles have found numerous applications in science, health, and industries. Magnetite nanoparticles (Fe_3O_4 :MNPs) have attracted attention because of their peroxidase-like activity. In this study, hydroxyl radicals ($\cdot\text{OH}$) generated by MNPs-catalyzed degradation of H_2O_2 converted the N,N-dimethyl-*p*-phenylenediamine (DMPD) probe into its colored DMPD $^{\cdot+}$ radical cation, which gave an absorbance maximum at $\lambda = 553$ nm. In the presence of antioxidants, $\cdot\text{OH}$ was partly scavenged by antioxidants and produced less DMPD $^{\cdot+}$, causing a decrease in the 553 nm-absorbance. Antioxidant concentrations were calculated with the aid of absorbance differences between the reference and sample solutions. The linear working ranges and trolox equivalent antioxidant capacity coefficients of different classes of antioxidants were determined by applying the developed method. In addition, binary and ternary mixtures of antioxidants were tested to observe the additivity of absorbances of mixture constituents. The method was applied to real samples such as orange juice and green tea. Student t-test, F tests, and the Spearman's rank correlation coefficient were used for statistical comparisons.

Key words: Magnetite (Fe_3O_4) nanoparticles, reactive oxygen species (ROS), N,N-Dimethyl-*p*-phenylenediamine, colorimetric probe, antioxidant activity

1. Introduction

Reactive oxygen species (ROS) is a collective term expressing both radicalic oxygen species [e.g., hydroxyl ($\cdot\text{OH}$) and superoxide radicals ($\text{O}_2^{\cdot-}$)] together with other nonradical derivatives of oxygen, such as, singlet oxygen ($^1\text{O}_2$) and hydrogen peroxide (H_2O_2). Due to beneficial and harmful health effects, the amount of ROS in the body plays a central role in biological processes. Oxidative damage of ROS to cells and biomolecules [1,2] may be compensated for by intrinsic and extrinsic antioxidants. If the delicate balance between ROS generation and antioxidative defenses of the organism is not conserved, oxidative stress conditions may occur [1]. Oxidative stress is related to the general aging process and cell death, affecting all organ systems [3]. ROS also plays a part in many age-related diseases, such as Alzheimer [4], Parkinson [5], and cardiovascular diseases [6]. Among all ROS, $\cdot\text{OH}$ is considered as one of the most reactive species, because it can rapidly attack biomolecules with a high oxidation potential and cause irreversible damage [7]. Therefore, the measurement of ROS scavenging activity of different classes of antioxidants is important. Quantitative analysis of total antioxidant activity can be carried out using globally adopted molecular spectroscopic methods. Among the free radical scavenging assays, those using artificial radicals such as ABTS (2,2'-azinobis (3-ethylbenzothiazoline-6-sulfonic acid) [8,9] and DPPH (2,2-diphenyl-1-picrylhydrazyl) [10, 11], or ORAC as the peroxy radical scavenging method (Oxygen Radical Absorbance Capacity) [12,13] have emerged as the most popular ones. The transition metal ion reducing capacity of antioxidants can be measured with CUPRAC (Cupric Reducing Antioxidant Capacity) [14,15] and FRAP (Ferric Ion Reducing Antioxidant Power) assays [16-19]. Many alternative spectrophotometric [20–23] and voltammetric methods [24,25] were later developed, based on different mechanisms of antioxidant and radical scavenging action. With the advancement of nanoscience, many new methods are on the rise due to extraordinary physico-chemical properties of nanoparticles, such as large surface-to-volume ratio for enhanced catalysis and surface plasmon resonance absorption giving rise to high sensitivity.

* Correspondence: auzer@istanbul.edu.tr

The peroxidase-like activity of MNPs has already been established [26] and applied to many different areas such as the removal of organic pollutants [27,28], hydrogen peroxide [29–31], and analysis of home-made explosives (e.g., TATP) [32]. The peroxidase-like activity of Fe_3O_4 NPs is based on the simultaneous existence of $\text{Fe}^{2+}/\text{Fe}^{3+}$ ion pair in the structure, having a similarity to Fenton's reagent [26].

In the developed method, hydroxyl radicals were produced through MNPs-catalyzed degradation of H_2O_2 , owing to the peroxidase-like activity of Fe_3O_4 . Since a part of the obtained hydroxyl radicals were scavenged with antioxidants, the remaining $\cdot\text{OH}$ could oxidize DMPD to the colored DMPD^+ radical cation. Both the reference (the solution containing no antioxidant) and samples were measured against water at 553 nm. For the calculations, the difference between the reference and sample absorbances (ΔA) was used. It is better to monitor a Fenton-type reaction of $\cdot\text{OH}$ generation with MNPs because Fe_3O_4 can be magnetically separated at any stage while soluble Fe(II) -catalyzed degradation of H_2O_2 in homogenous solution is less controllable. The developed method was applied to different classes of antioxidants and their binary/ternary mixtures. In addition, it was applied to real samples such as, orange juice and green tea, and its statistical comparisons were made with the aid of Student t-test, F test, and Spearman's rank correlation tests.

2. Materials and methods

2.1. Instrumentation and chemicals

Shimadzu UV-1800 (Kyoto, Japon) spectrophotometer was used for absorption measurements. Hellma GmbH & Co. KG (Müllheim, Germany) quartz cuvettes with 1 cm optical thickness were used for the UV-Vis measurements.

All reagents were of analytical reagent grade unless otherwise stated. Quercetin (QR), gallic acid (GA), ferulic acid (FA), caffeic acid (CFA), catechin (CAT), epicatechin, trolox (TR), para-coumaric acid (p-CUM), L-ascorbic acid (AA), L-cysteine (CYS), morin (MR), rutin (RT), hydrogen peroxide (30% by mass), and N,N-Dimethyl-p-phenylenediamine (DMPD) were purchased from Sigma-Aldrich Chemie GmbH (Steinheim, Germany), and all other reagents from Merck KGaA (Darmstadt, Germany) and Sigma-Aldrich Chemie GmbH. Commercial orange juice and green tea samples were obtained from local markets.

2.2. Preparation of solutions

Except for the ascorbic acid, all antioxidant solutions were prepared in ethanol. Ascorbic acid, mixtures containing AA, and diluted orange juice samples were prepared in 1:1 (v/v) ethanol:water.

H_2O_2 solution ($4 \times 10^{-4} \text{ mol L}^{-1}$), pH 3.6 acetic acid buffer solution (containing a total $\text{CH}_3\text{COOH}/\text{CH}_3\text{COONa}$ concentration of 2 mol L^{-1}), Fe_3O_4 NPs solution (400 mg L^{-1}), and N,N-dimethyl-p-phenylenediamine (DMPD) solution ($2 \times 10^{-3} \text{ mol L}^{-1}$) were prepared in ultrapure water.

To prepare the green tea sample, 50 mL boiled water was added to 0.5 g of green tea and let to stand for 15 min for infusion [25]. Then, the final solution was passed through filter paper and diluted to 50 mL with ultrapure water. This stock solution was appropriately diluted and the proposed method was applied.

For the orange juice sample, orange juice was centrifuged at 10000 rpm for 20 min and the supernatant filtered with the aid of a microfilter. The proposed method was applied to this sample with appropriate dilution. The results of real samples were given in trolox equivalents (TE).

2.3. Standard addition of AA to orange juice

To determine the recovery of AA from orange juice samples, the standard addition method was applied. The analyses of 1:200 diluted orange juice sample, $2 \times 10^{-5} \text{ mol L}^{-1}$ AA solution at final concentration, and AA-added solution were carried out separately with the proposed method.

2.4. Synthesis of Fe_3O_4 NPs

The coprecipitation method was carried out for synthesizing Fe_3O_4 NPs [29]. In the beginning, 10 mL of FeCl_2 (2 mol L^{-1}) solution in 2 mol L^{-1} HCl were mixed with 50 mL of FeCl_3 (1 mol L^{-1}) solution. After deoxygenation with nitrogen gas, the mixture solution was added drop wise to 500 mL of oxygen-free aqueous ammonia solution (0.7 mol L^{-1}). The final solution was stirred under a nitrogen atmosphere for 30 min. Fe_3O_4 NPs were separated and washed with water.

2.5. Application of the proposed method

One milliliter of $4 \times 10^{-4} \text{ mol L}^{-1}$ H_2O_2 solution, 1 mL of pH 3.6 acetate buffer, 1 mL of 400 mg L^{-1} Fe_3O_4 NPs solution, 1 mL of antioxidant solution (for reference solution, depending on the solvent of the sample, EtOH or 1:1 (v:v) EtOH: water was used), and 1 mL of $2 \times 10^{-3} \text{ mol L}^{-1}$ DMPD solution were mixed in this order. Finally, at the end of the 30 min, the absorbances of all solutions (including the reference solution) were measured against water at 553 nm. Antioxidant activity was expressed as the decrease of absorbance in the presence of scavengers.

2.6. Statistical analysis

CFA and RT were analyzed by the thiobarbituric acid reactive substances (TBARS) method, based on colored product formation of TBA with the peroxidation end product malondialdehyde [20]. CFA and RT solutions were performed with the proposed method at 4×10^{-6} and 2×10^{-4} mol L⁻¹, and with TBARS method at 4×10^{-6} and 2×10^{-5} mol L⁻¹, respectively. Rutin results, which were obtained by TBARS method, were multiplied by the dilution factor (10) for correction of concentration. The obtained results from both methods were used for statistical comparison using Student t-test and F tests. Also, Spearman's rank correlation (using Equation 1) was used for statistical comparison of ROS scavenging activity of selected antioxidants [33].

$$r_s = 1 - \frac{6 \sum d_i^2}{n(n^2 - 1)} \quad (1)$$

r_s = Spearman's rank correlation,

d_i = Difference between two ranks,

n = Number of pairs.

3. Results

When scanning electron microscopy (SEM) analysis was performed to observe the shape and size of synthesized magnetite nanoparticles, they were found to be spherical and in between 20 and 30 nm in diameter (Figure 1). It is known from the literature that smaller size magnetite nanoparticles show higher catalytic activity [26]. Magnetic properties of magnetite nanoparticles were also observed by applying a magnetic field with a magnet.

The proposed method is based on the production of ROS (i.e. mainly hydroxyl and perhydroxyl radicals) from H₂O₂ with the aid of Fe₃O₄ NPs at pH 3.6, scavenging of produced ROS by antioxidants, and finally oxidation by remaining ROS of the DMPD probe to the pink colored DMPD^{•+} radical cation (Scheme). Spectrophotometric measurements were performed with this obtained colored product.

3.1. Optimization of reaction parameters

For optimization of the developed method, H₂O₂, Fe₃O₄ NPs, and DMPD concentrations were investigated. As shown in Figures 2–4, for increased concentrations of substances, increasing absorbance values were obtained. For selecting the appropriate concentrations of reagents, attention was paid to the absorbance value of the reference solution which should lie in the range of 1.1–1.2. According to this precaution, H₂O₂ concentration was determined as 4×10^{-3} mol L⁻¹, Fe₃O₄ NPs





Figure 1. SEM image of 20–30 nm sized Fe₃O₄ particles.



Scheme. Schematic presentation of ROS scavenging activity of antioxidants with the proposed method.




Figure 2. Effect of H_2O_2 concentration on DMPD⁺⁺ absorbance. Working conditions: 1 mL of H_2O_2 + 1 mL of pH 3.6 acetic acid buffer + 1 mL of 400 mg L^{-1} Fe_3O_4 NPs + 1 mL of ethanol (wait for 5 min) + 1 mL of $5 \times 10^{-3} \text{ mol L}^{-1}$ DMPD (wait for 30 min) + centrifuge (for 5 min at 5000 rpm). All measurements were made against water at 553 nm.

concentration as 400 mg L^{-1} , and DMPD concentration as $2 \times 10^{-3} \text{ mol L}^{-1}$. For preserving the solubility and stability of magnetite nanoparticles, the selection of an appropriate working pH is important. In this work, Fe_3O_4 nanoparticles were used as a catalyst in conjunction with DMPD sensing. Without deactivating the catalyst and without allowing ferric iron to hydrolyze, an optimal pH of 3.6 was selected, as in the literature [34,35].

3.2. Determination of hydroxyl radical scavenging activity of antioxidants with the aid of Fe_3O_4 NPs and DMPD probe
The developed method was applied to CYS, AA, TR, and different classes of antioxidants such as flavanol (QR, MR), phenolic acid (GA, FA, CFA, p-CUM), flavanol (CAT, EC), and flavon (RT). The calibration curves were drawn between antioxidant concentrations and absorbance decrements (ΔA). When the developed method was applied to caffeic acid, the obtained DMPD⁺⁺ spectra are shown in Figure 5, indicating absorbance decrements with increasing CFA concentration. Two peaks were observed at 553 and 515 nm, with quite close absorbance values. The 553 nm-peak was preferred for further measurements so that it would be less affected by plant pigments absorbing in the visible range.




Figure 3. Effect of Fe₃O₄ NPs concentration on DMPD⁺⁺ absorbance. Working conditions: 1 mL of 4×10^{-4} mol L⁻¹ H₂O₂ + 1 mL of pH 3.6 acetic acid buffer + 1 mL of Fe₃O₄ NPs + 1 mL of ethanol (wait for 5 min) + 1 mL of 5×10^{-3} mol L⁻¹ DMPD (wait for 30 min) + centrifuge (for 5 min at 5000 rpm). All measurements were made against water at 553 nm.




Figure 4. Effect of DMPD concentration on absorbance. Working conditions: 1 mL of 4×10^{-4} mol L⁻¹ H₂O₂ + 1 mL of pH 3.6 acetic acid buffer + 1 mL of 400 mg L⁻¹ Fe₃O₄ NPs + 1 mL of ethanol (wait for 5 min) + 1 mL of DMPD (wait for 30 min) + centrifuge (for 5 min at 5000 rpm). All measurements were made against water at 553 nm.

The calibration equations, linear ranges of antioxidant concentration, trolox equivalent antioxidant capacity (TEAC) values, and correlation coefficients are shown in Table 1. According to the developed method, hydroxyl radical scavenging efficiency of tested antioxidants followed the order: EC > CFA > FA > GA > CAT > CYS > QR > MR > AA > p-CUM > RT. This finding complies with the hypothesis of Bors and Michel based on experimental results of pulse radiolysis transient spectra and decay rate constants of \cdot OH-absorbed species that catechol and/or pyrogallol groups were the active sites of radical attack, with subsequent conversion to quinoid compounds [36]. In this research, catechins, CFA, and GA with o-dihydroxy phenol (catechol) substitution were the leading \cdot OH scavengers, in accordance with the findings of Demirci Çekiç et al. [21] who demonstrated the priority of o-catechol and galloyl moieties of phenolic antioxidants in hydroxyl radical scavenging ability measured with a DMPD probe on a Nafion membrane.

3.3. Analysis of synthetic antioxidant mixtures and real samples

When the additivity of differential absorbances of antioxidants in their binary and ternary synthetic mixtures was tested with the proposed method, the percentage relative errors (calculated from the difference of expected and found values) of




Figure 5. The DMPD^{•+} spectra showing the [•]OH scavenging activity of different concentrations of caffeic acid solutions.

mixtures ranged between -7 and $+5\%$ (Table 2). This basically showed the absence of chemical deviations from Beer's law in the proposed method.

As real samples, orange juice and green tea were analyzed with the proposed method, and the results were calculated in trolox equivalents (TE) as defined in the literature [24,25]. The antioxidant capacity of orange juice, and green tea were found as $2.33 \text{ mmol TE L}^{-1}$, and $1.23 \text{ mmol TE g}^{-1}$, respectively. One juice sample was also spiked with an antioxidant compound to observe recovery and to confirm the absence of chemical deviations from Beer's law in a real sample. Standard addition of a certain concentration of $2 \times 10^{-5} \text{ mol L}^{-1}$ AA solution to diluted orange juice yielded an AA recovery of 97% with the proposed method.

3.4. Statistical evaluation of the developed method

For method validation, reference TBARS method was applied to CFA and RT solutions [20]. The CFA and RT solutions in ethanol at $1.0 \times 10^{-6} - 5.0 \times 10^{-6} \text{ mol L}^{-1}$ concentrations were measured and calibration curves were obtained. A CFA solution at $4 \times 10^{-6} \text{ mol L}^{-1}$ concentration was tested using both the developed and TBARS methods. For method validation with RT, a solution at $2 \times 10^{-4} \text{ mol L}^{-1}$ concentration was used for the developed method and $2 \times 10^{-5} \text{ mol L}^{-1}$ concentration was used for TBARS method, and then the calculated result was multiplied by the dilution factor. The measurements were carried out for $N = 5$ repetitive analyses. The compared methods showed no significant differences between the precision and accuracy of results found for CFA and RT (Table 3). The Student t-test and F tests were used for statistical comparison of the population means and variances, respectively, at 95% confidence levels for both tests (Table 3).

For comparing the proposed method with ABTS (radical scavenging) reference assay using the TEAC values of FA, GA, CAT, QR, MR, AA, p-CUM, and RT, Spearman's rank correlation was used (Table 4). The results showed a 'strong' (Spearman's rank correlation coefficient, $r_s = 0.78571$) relationship between the developed method and ABTS assay ($P = 0.02082$; two-tailed).

4. Discussion

A novel easy and low-cost DMPD-based ROS scavenging activity method was developed with the aid of Fe_3O_4 NPs having peroxidase-like activity. MNPs catalyzed the degradation of H_2O_2 to ROS (mainly hydroxyl radicals) which in turn converted DMPD to a colored radical cation, and antioxidants (when present) quenched [•]OH, thereby causing a decrement in the DMPD^{•+} absorbance. Although a Fenton-type reaction (catalyzed by soluble Fe^{2+} ions) in homogeneous solution can also generate [•]OH from H_2O_2 , the MNPs used in this work enable a more easily monitored reaction as the

Table 1. The linear concentration ranges, calibration equations, and TEAC coefficients of the tested antioxidants in the proposed method.

Antioxidant	Linear range (Final concentrations) (mol L ⁻¹)	Calibration equations and TEAC values
Trolox (TR)	$2 \times 10^{-5} - 1.2 \times 10^{-4}$	$\Delta A = 1.05 \times 10^4 C - 0.04$ ($r = 0.9966$)
Quercetin (QR)	$2 \times 10^{-5} - 1 \times 10^{-4}$	$\Delta A = 9.96 \times 10^3 C - 0.03$ ($r = 0.9981$) (TEAC Coeff. = 0.95)
Gallic acid (GA)	$2 \times 10^{-6} - 2 \times 10^{-5}$	$\Delta A = 2.07 \times 10^4 C + 0.14$ ($r = 0.9976$) (TEAC Coeff. = 1.97)
Ferulic acid (FA)	$2 \times 10^{-6} - 2 \times 10^{-5}$	$\Delta A = 2.54 \times 10^4 C + 0.03$ ($r = 0.9966$) (TEAC Coeff. = 2.42)
Caffeic acid (CFA)	$2 \times 10^{-6} - 2 \times 10^{-5}$	$\Delta A = 3.05 \times 10^4 C + 0.03$ ($r = 0.9977$) (TEAC Coeff. = 2.90)
Catechin (CAT)	$2 \times 10^{-6} - 2 \times 10^{-5}$	$\Delta A = 1.7 \times 10^4 C + 0.09$ ($r = 0.9992$) (TEAC Coeff. = 1.62)
Epicatechin (EC)	$2 \times 10^{-6} - 2 \times 10^{-5}$	$\Delta A = 3.12 \times 10^4 C + 0.04$ ($r = 0.9982$) (TEAC Coeff. = 2.97)
p-coumaric acid (p-CUM)	$1 \times 10^{-4} - 1 \times 10^{-3}$	$\Delta A = 6.56 \times 10^2 C + 0.28$ ($r = 0.9917$) (TEAC Coeff. = 0.06)
Ascorbic acid (AA)	$2 \times 10^{-5} - 2 \times 10^{-4}$	$\Delta A = 2.01 \times 10^3 C + 0.08$ ($r = 0.9996$) (TEAC Coeff. = 0.19)
L-cysteine (CYS)	$1 \times 10^{-5} - 4 \times 10^{-5}$	$\Delta A = 1.53 \times 10^4 C + 0.09$ ($r = 0.9948$) (TEAC Coeff. = 1.46)
Morin (MR)	$1.2 \times 10^{-5} - 2 \times 10^{-4}$	$\Delta A = 6.28 \times 10^3 C + 0.08$ ($r = 0.9951$) (TEAC Coeff. = 0.6)
Rutin (RT)	$1 \times 10^{-4} - 1 \times 10^{-3}$	$\Delta A = 3.29 \times 10^2 C + 0.16$ ($r = 0.9979$) (TEAC Coeff. = 0.03)

reaction can be stopped at any stage by magnetic separation of Fe₃O₄ NPs. ROS scavenging activity of different classes of antioxidants was determined with the developed method. Polyphenol compounds are attacked by hydroxyl radical predominantly at the o-dihydroxy site [37]; this can be explained by the formation of stable semiquinone radicals (i.e. stabilized by intramolecular hydrogen bonding) from the reaction of the o-dihydroxy group with hydroxyl radicals [38]. Inspection of the most effective hydroxyl radical scavengers of this work, i.e. EC > CFA > FA > GA > CAT, reveals that all these compounds have the 3,4-dihydroxyphenyl (FA having 4-hydroxy-3-methoxyphenyl) moiety that can be attacked by

Table 2. Relative errors of binary and ternary synthetic antioxidant mixtures.

Antioxidant mixtures (in final conc.)	Relative errors (%)
$5 \times 10^{-6} \text{ mol L}^{-1} \text{ CAT} + 6 \times 10^{-6} \text{ mol L}^{-1} \text{ FA}$	-2.00
$5 \times 10^{-6} \text{ mol L}^{-1} \text{ CAT} + 4 \times 10^{-5} \text{ mol L}^{-1} \text{ QR}$	-0.04
$5 \times 10^{-6} \text{ mol L}^{-1} \text{ CAT} + 4 \times 10^{-5} \text{ mol L}^{-1} \text{ Morin}$	-1.00
$5 \times 10^{-5} \text{ mol L}^{-1} \text{ AA} + 2 \times 10^{-5} \text{ mol L}^{-1} \text{ GA}$	-3.00
$5 \times 10^{-5} \text{ mol L}^{-1} \text{ AA} + 6 \times 10^{-6} \text{ mol L}^{-1} \text{ FA}$	-3.00
$5 \times 10^{-5} \text{ mol L}^{-1} \text{ AA} + 4 \times 10^{-5} \text{ mol L}^{-1} \text{ QR}$	-7.00
$1 \times 10^{-6} \text{ mol L}^{-1} \text{ CFA} + 1 \times 10^{-5} \text{ mol L}^{-1} \text{ QR} + 1 \times 10^{-6} \text{ mol L}^{-1} \text{ CAT}$	-6.00
$1 \times 10^{-5} \text{ mol L}^{-1} \text{ TR} + 1 \times 10^{-6} \text{ mol L}^{-1} \text{ CAT} + 1 \times 10^{-6} \text{ mol L}^{-1} \text{ GA}$	+5.00
$2 \times 10^{-5} \text{ mol L}^{-1} \text{ AA} + 2 \times 10^{-5} \text{ mol L}^{-1} \text{ TR} + 2 \times 10^{-6} \text{ mol L}^{-1} \text{ CFA}$	-6.00

Table 3. Statistical comparison of the developed method with reference TBARS method for CFA and RT solutions with the aid of Student t-test and F tests.

Analyte	Method	Mean conc. (mol L ⁻¹)	SD (σ)	S ^{a,b}	t ^{a,b}	t _{table} ^b	F ^b	F _{table} ^b
CFA	Developed method	4.30×10^{-6}	2.27×10^{-7}	-	-	-	-	-
	TBARS method	4.14×10^{-6}	2.27×10^{-7}	3.08×10^{-7}	0.846	2.306	0.371	6.39
RT	Developed Method	2.10×10^{-4}	3.18×10^{-5}	-	-	-	-	-
	TBARS Method	2.17×10^{-4}	2.22×10^{-5}	2.74×10^{-5}	0.415	2.306	2.045	6.39

^a $S^2 = \{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2\} / (n_1 + n_2 - 2)$ and $t = (\bar{a}_1 - \bar{a}_2) / \{S(1/n_1 + 1/n_2)^{1/2}\}$, where S is the pooled standard deviation, s_1 and s_2 are the standard deviations of the two populations with sample sizes of n_1 and n_2 , and sample means of \bar{a}_1 and \bar{a}_2 respectively (t has $(n_1 + n_2 - 2)$ degrees of freedom); here, $n_1 = n_2 = 5$.

^b Statistical comparison made on paired data produced with proposed and reference methods; the results given only on the row of the reference method.

Table 4. Spearman's rank correlation analysis according to TEAC values of antioxidants.

	1	2	3	4	5	6	7	8	r _s	Definition
Developed method	FA	GA	CAT	QR	MR	AA	p-CUM	RT	0.78571	Strong
ABTS/persulfate	GA	CAT	QR	FA	MR	p-CUM	RT	AA		

*OH to produce stable semiquinone radicals. Likewise, the cell protective activity of quercetin against oxidative hazard was attributed to certain structure-activity relationships including the o-dihydroxy structure in the B-ring [39]. Also, binary and ternary synthetic antioxidant mixtures were measured and the additivity of antioxidant concentrations (in trolox equivalents) was observed. In addition, the developed method was successfully applied to the real samples of green tea and orange juice. The proposed method was validated against the TBARS method through analysis of CFA and RT standard solutions. Spearman's rank correlation (tested on 8 antioxidants) was found 'strong' between the developed method and the ABTS (radical scavenging) assay.

Acknowledgment

This study was funded by İstanbul University-Cerrahpaşa with the project number 28963. We express our gratitude to İstanbul University-Cerrahpaşa Application and Research Center for the Measurement of Food Antioxidants (GAAM) for assay development.

References

1. Halliwell B. Free radicals and antioxidants—quo vadis? *Trends in Pharmacological Sciences* 2011; 32 (3): 125-130. doi: 10.1016/j.tips.2010.12
2. Winterbourn CC. Reconciling the chemistry and biology of reactive oxygen species. *Nature Chemical Biology* 2008; 4: 278-286. doi: 10.1038/nchembio.85
3. Krause KH. Aging: a revisited theory based on free radicals generated by NOX family NADPH oxidases. *Experimental Gerontology* 2007; 42 (4): 256-262. doi: 10.1016/j.exger.2006.10.011
4. Qin B, Cartier L, Dubois-Dauphin M, Li B, Serrander L et al. A key role for the microglial NADPH oxidase in APP-dependent killing of neurons. *Neurobiology of Aging* 2006; 27 (11): 1577-1587. doi: 10.1016/j.neurobiolaging.2005.09.036
5. Zhang Y, Dawson VL, Dawson TM. Oxidative stress and genetics in the pathogenesis of Parkinson's disease. *Neurobiology of Disease* 2000; 7 (4): 240-250. doi: 10.1006/nbdi.2000.0319
6. Dhalla NS, Temsah RM, Netticadan T. Role of oxidative stress in cardiovascular diseases. *Journal of Hypertension* 2000; 18 (6): 655-673.
7. Phaniendra A, Jestadi DB, Periyasamy L. Free radicals: properties, sources, targets, and their implication in various diseases. *Indian Journal of Clinical Biochemistry* 2015; 30 (1): 11-26. doi: 10.1007/s12291-014-0446-0
8. Pellegrini N, Re R, Yang M, Rice-Evans C. Screening of dietary carotenoids and carotenoid-rich fruit extracts for antioxidant activities applying 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) radical cation decolorization assay. *Methods Enzymology* 1999; 299: 379-389. doi: 10.1016/S0076-6879(99)99037-7
9. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M et al. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology and Medicine* 1999; 26 (9-10): 1231-1237. doi: 10.1016/S0891-5849(98)00315-3
10. Blois MS. Antioxidant determinations by the use of a stable free radical. *Nature* 1958; 181: 1199-1200. doi: 10.1038/1811199a0
11. Sanchez-Moreno C, Larrauri JA, Saura-Calixto FA. A procedure to measure the antiradical efficiency of polyphenols. *Journal of the Science of Food and Agriculture* 1998; 76 (2): 270-276. doi: 10.1002/(SICI)1097-0010(199802)76:2<270::AID-JSFA945>3.0.CO;2-9
12. Cao G, Alessio HM, Cutler RG. Oxygen-radical absorbance capacity assay for antioxidants. *Free Radical Biology and Medicine* 1993; 14 (3): 303-311. doi: 10.1016/0891-5849(93)90027-r
13. Ou B, Hampsch-Woodill M, Prior RL. Development and validation of an improved oxygen radical absorbance capacity assay using fluorescein as the fluorescent probe. *Journal of Agricultural and Food Chemistry* 2001; 49 (10): 4619-4626. doi: 10.1021/jf010586o
14. Apak R, Güçlü K, Özyürek M, Karademir SE. Novel total antioxidant capacity index for dietary polyphenols and vitamins C and E, using their cupric ion reducing capability in the presence of neocuproine: CUPRAC method. *Journal of Agricultural and Food Chemistry* 2004; 52 (26): 7970-7981. doi: 10.1021/jf048741x
15. Apak R, Güçlü K, Özyürek M, Karademir SE, Altun M. Total antioxidant capacity assay of human serum using copper (II)-neocuproine as chromogenic oxidant: the CUPRAC method. *Free Radical Research* 2005; 39 (9): 949-961. doi: 10.1080/10715760500210145
16. Benzie IFF, Strain JJ. The ferric reducing ability of plasma (FRAP) as a measure of 'antioxidant power': the FRAP assay. *Analytical Biochemistry* 1996; 239 (1): 70-76. doi: 10.1006/abio.1996.0292
17. Benzie IFF, Strain JJ. Ferric reducing/antioxidant power assay: direct measure of total antioxidant activity of biological fluids and modified version for simultaneous measurement of total antioxidant power and ascorbic acid concentration. *Methods Enzymology* 1999; 299: 15-27. doi: 10.1016/S0076-6879(99)99005-5
18. Benzie IFF, Szeto YT. Total antioxidant capacity of teas by the ferric reducing/antioxidant power assay. *Journal of Agricultural and Food Chemistry* 1999; 47 (2): 633-636. doi: 10.1021/jf9807768
19. Pulido R, Bravo L, Saura-Calixto F. Antioxidant activity of dietary polyphenols as determined by a modified ferric reducing/antioxidant power assay. *Journal of Agricultural and Food Chemistry* 2000; 48 (8): 3396-3402. doi: 10.1021/jf9913458
20. Bektaşoğlu B, Özyürek M, Güçlü K, Apak R. Hydroxyl radical detection with salicylate probe using modified CUPRAC spectrophotometry and HPLC. *Talanta* 2008; 77 (1): 90-97. doi: 10.1016/j.talanta.2008.05.043
21. Demirci Çekiç S, Avan AN, Uzunboş S, Apak R. A colorimetric sensor for the simultaneous determination of oxidative status and antioxidant activity on the same membrane: N,N-Dimethyl-p-phenylene diamine (DMPD) on Nafion. *Analytica Chimica Acta* 2015; 865: 60-70. doi: 10.1016/j.aca.2015.01.041
22. Kamer G, Demirata B, Bayraktar R, Özyurt D, Apak R. Nanoceria-based reactive species scavenging activity of antioxidants using N,N-dimethyl-p-phenylenediamine (DMPD) probe. *Analytical Methods* 2019; 11 (14): 1908-1915. doi: 10.1039/C8AY02561D
23. Can K, Üzer A, Apak R. A manganese oxide (MnO₂)-Based colorimetric nanosensor for indirect measurement of lipophilic and hydrophilic antioxidant capacity. *Analytical Methods* 2020; 12 (4): 448-455. doi: 10.1039/c9ay02027f

24. Tufan AN, Baki S, Güçlü K, Özyürek M, Apak R. A novel differential pulse voltammetric (DPV) method for measuring the antioxidant capacity of polyphenols-reducing cupric neocuproine complex. *Journal of Agricultural and Food Chemistry* 2014; 62 (29): 7111-7117. doi: 10.1021/jf5017797
25. Arman A, Üzer A, Sağlam Ş, Erçağ E, Apak R. Indirect electrochemical determination of antioxidant capacity with hexacyanoferrate(III) reduction using a gold nanoparticle-coated o-phenylenediamine-aniline copolymer electrode. *Analytical Letters* 2019; 52 (8): 1282-1297. doi: 10.1080/00032719.2018.1536137
26. Gao L, Zhuang J, Nie L, Zhang J, Zhan Y et al. Intrinsic peroxidase-like activity of ferromagnetic nanoparticles. *Nature Nanotechnology* 2007; 2: 577-583. doi: 10.1038/nnano.2007.260
27. Wang N, Zhu L, Wang D, Wang M, Lin Z et al. Sono-assisted preparation of highly-efficient peroxidase-like Fe_3O_4 magnetic nanoparticles for catalytic removal of organic pollutants with H_2O_2 . *Ultrasonics Sonochemistry* 2010; 17 (3): 526-533. doi: 10.1016/j.ultsonch.2009.11.001
28. Zhang S, Zhao X, Niu H, Shi Y, Cai Y et al. Superparamagnetic Fe_3O_4 nanoparticles as catalysts for the catalytic oxidation of phenolic and aniline compounds. *Journal of Hazardous Materials* 2009; 167 (1-3): 560-566. doi: 10.1016/j.jhazmat.2009.01.024
29. Wei H, Wang E. Fe_3O_4 Magnetic nanoparticles as peroxidase mimetics and their applications in H_2O_2 and glucose detection. *Analytical Chemistry* 2008; 80 (6): 2250-2254. doi: 10.1021/ac702203f
30. Chang Q, Deng KJ, Zhu LH, Jiang GD, Yu C et al. Determination of hydrogen peroxide with the aid of peroxidase-like Fe_3O_4 magnetic nanoparticles as the catalyst. *Microchimica Acta* 2009; 165: 299-305. doi: 10.1007/s00604-008-0133-z
31. Zhuang J, Zhang JB, Gao LZ, Zhang Y, Gu N et al. A novel application of iron oxide nanoparticles for detection of hydrogen peroxide in acid rain. *Materials Letters* 2008; 62 (24): 3972-3974. doi: 10.1016/j.matlet.2008.05.025
32. Can Z, Üzer A, Türkekul K, Erçağ E, Apak R. Determination of triacetone triperoxide with a N,N-Dimethyl-p-phenylenediamine sensor on nafion using Fe_3O_4 magnetic nanoparticles. *Analytical Chemistry* 2015; 87 (19): 9589-9594. doi: 10.1021/acs.analchem.5b01775
33. Miller J, Miller JC. *Statistics and Chemometrics for Analytical Chemistry*. Gosport, United Kingdom: Pearson, 2010.
34. Tombacz E, Illes E, Majzik A, Hajdu A, Rideg N et al. Ageing in the inorganic nanoworld: example of magnetite nanoparticles in aqueous medium. *Croatica Chemica Acta* 2007; 80 (3-4): 503-515.
35. Sun ZX, Su FW, Forsling W, Samskog POJ. Surface characteristics of magnetite in aqueous suspension. *Journal of Colloid and Interface Science* 1998; 197 (1): 151-159. doi: 10.1006/jcis.1997.5239
36. Bors W, Michel C. Antioxidant capacity of flavanols and gallate esters: pulse radiolysis studies. *Free Radical Biology and Medicine* 1999; 27 (11-12): 1413-1426. doi: 10.1016/S0891-5849(99)00187-2
37. Bors W, Heller W, Michel C, Saran M. Flavonoids as antioxidants: determination of radical-scavenging efficiencies. *Methods in Enzymology* 1990; 186: 344-355. doi: 10.1016/0076-6879(90)86128-i
38. Wang P, Kang J, Zheng R, Yang Z, Lu J et al. Scavenging effects of phenylpropanoid glycosides from *Pedicularis* on superoxide anion and hydroxyl radical by the spin trapping method (95)02255-4. *Biochemical Pharmacology* 1996; 51 (5):687-691. doi: 10.1016/s0006-2952(95)02255-4.
39. Metodiewa D, Jaiswal AK, Cenas N, Dickanaité E, Segura-Aguilar J. Quercetin may act as a cytotoxic prooxidant after its metabolic activation to semiquinone and quinoidal product. *Free Radical Biology and Medicine* 1999; 26 (1-2): 107-116. doi: 10.1016/S0891-5849(98)00167-1