

Volatile constituents of three *Thymus sipyleus* Boiss. subspecies from different sites in Turkey

Hale Gamze AĞALAR^{1*}, Mine KÜRKÇÜOĞLU¹, Kemal Hüsnü Can BAŞER², Kenan TURGUT³

¹Department of Pharmacognosy, Faculty of Pharmacy, Anadolu University, Eskişehir, Turkey

²Department of Pharmacognosy, Faculty of Pharmacy, Near East University, Lefkoşa, North Cyprus

³Department of Field Crops, Faculty of Agriculture, Akdeniz University, Antalya, Turkey

Received: 04.03.2021 • Accepted/Published Online: 20.09.2021 • Final Version: 20.12.2021

Abstract: This study was designed to reveal the chemical diversity of some *Thymus sipyleus* subspecies growing wild in Turkey and to compare the volatile compound profiles by using micro(hydro)distillation technique. For this purpose, volatile compounds isolated by microdistillation from nine samples (three plant samples collected from different regions in Antalya) of *Thymus sipyleus* Boiss. subsp. *sipyleus* var. *sipyleus*, *T. sipyleus* Boiss. subsp. *sipyleus* var. *davisanus* Ronniger, and *T. sipyleus* Boiss. subsp. *rosulans* (Borbas) Jalas were analyzed by GC and GC-MS systems. 1,8-Cineole, *p*-cymene, α -terpineol and carvacrol were identified as major compounds in *T. sipyleus* subsp. *sipyleus* var. *sipyleus* samples. Geranial, neral, 1,8-cineole and β -caryophyllene, and α -terpineol and geranial were the main compounds in *T. sipyleus* subsp. *sipyleus* var. *davisanus* samples. β -Caryophyllene, intermedeol, 1,8-cineole and α -terpineol, α -pinene were the major compounds in *T. sipyleus* subsp. *rosulans* samples. As known, thymol is the main compound in most *Thymus* species in Turkey, but, according to our study, chemical polymorphism has been found among the *T. sipyleus* subspecies.

Key words: *Thymus sipyleus* subsp. *sipyleus* var. *sipyleus*, *T. sipyleus* subsp. *sipyleus* var. *davisanus*, *T. sipyleus* subsp. *rosulans*, microdistillation, GC and GC-MS analysis, chemical polymorphism, terpenes

1. Introduction

The genus *Thymus* is note-worthy among the numerous species and varieties of wild-growing aromatic plants belonging to the family Lamiaceae. Many of these species are typical for the Mediterranean area. The genus *Thymus* is represented by 42 species and 47 taxa, 20 of which are endemic in Turkey [1]. All of them produce essential oils, and only a few are important herbs used in all parts of the world. Most of the terpenoid volatiles detected in *Thymus* oils belong to the monoterpene group. Sesquiterpenes are always present, but with only a few exceptions in minor percentages [2].

Most of these taxa growing in Turkey are aromatic plants which are generally used as herbal tea, condiments and in folk medicine. Carvacrol and thymol are abundant monoterpenes in the essential oils of this genus. However, there are *Thymus* species poor in phenolic compounds and some do not contain phenolic compounds at all. Phenol-rich *Thymus* species are used in diabetes, stomach and intestinal diseases, for cough as herbal tea and also as a condiment; whereas, phenol-poor or phenol-less *Thymus* species are used, due to their pleasant aroma, as herbal tea in Turkey [3].

Thymus L. is known in the world as 'thyme' and in Anatolia as 'kekik' or 'kaya kekiği'. Volatile oils of thyme are used as antiseptics, antispasmodics and fungicidal [4, 5]. The antiseptic, antioxidative, insecticidal, preservative and anaesthetic properties of thyme are due to their biologically active substances, such as thymol, carvacrol, linalool, geraniol and other volatiles in the essential oil [6]. In addition to the plant applications, thyme oils are also used in flavour and food industries, mainly in the manufacture of perfumes and cosmetics, or for flavouring chocolates, toothpaste, mouthwashes [7].

Due to the high economic value of *Thymus* species, a high number of studies on several aspects of this genus are available as well as the existing monographs on *Thymus* in Pharmacopoeias [8–10].

Thymus sipyleus Boiss. is endemic in Turkey, and known with local Turkish names as "kekik, limon kokulu kekik, keklik otu, yayla kekiği, nemamul otu, sater" [11]. According to ethnobotanical records, *T. sipyleus* and its subspecies are used for different purposes. In Adana, infusion of branches and leaves are consumed before meals for the treatment of stomach aches [12]. The aerial parts of *T. sipyleus* subsp. *sipyleus* var. *sipyleus* and *T. sipyleus* subsp. *sipyleus* var. *rosulans* are used as spice and tea (dried and grounded), in the treatment of haemorrhoids, atherosclerosis, and stomach disorders in Osmaniye, Bilecik [13]. The leaves of both subspecies also boiled with lemon as tea are taken against common cold and

* Correspondence: ecz.halegamze@gmail.com

coughs in Sivas and Yozgat [14]. In Ulukışla, Niğde, an infusion of the aerial parts of *T. sipyleus* subsp. *sipyleus* var. *sipyleus* is consumed three times a day for colds and stomach aches [15]. *T. sipyleus* subsp. *rosulans* known as “catri” in the Eastern part of Turkey is used for diabetes, colds, abdominal ailments as an infusion and decoction [16].

The present study is focused on determining the variation of volatile compounds from different populations of *Thymus sipyleus* Boiss. subsp. *sipyleus* var. *sipyleus*, *T. sipyleus* Boiss. subsp. *sipyleus* var. *davisianus* Ronniger, and *T. sipyleus* Boiss. subsp. *rosulans* (Borbas) Jalas. To date, the oil composition, biological activities of these subspecies of *Thymus sipyleus* have been reported [3, 11, 17–19]. In the present study, microdistilled aerial parts of *T. sipyleus* subsp. *sipyleus* var. *sipyleus*, *T. sipyleus* subsp. *sipyleus* var. *davisianus*, and *T. sipyleus* subsp. *rosulans* collected from different regions of Antalya, Turkey were analyzed by GC and GC-MS systems, simultaneously. Each microdistilled sample was characterized with major and minor volatile constituents by using in house and commercial libraries.

2. Materials and methods

2.1. Plant material

Air dried aerial parts of *T. sipyleus* subsp. *sipyleus* var. *sipyleus* (KT:190, 191, 192), *T. sipyleus* subsp. *sipyleus* var. *davisianus* (KT:196, 197, 198) and *T. sipyleus* subsp. *rosulans* (KT:199, 200, 201) were collected from three regions in Elmalı, Saklıkent, Gazipaşa (Antalya), respectively (Table 1). Identification of plant samples was done by one of us (KT). All herbarium samples coded as KT were kept at the Department of Field Crops, Faculty of Agriculture, Akdeniz University, Antalya, Turkey.

2.2. Isolation of the volatiles

Each sample was obtained by microdistillation of the dried, ground plant material (50 mg) using an Eppendorf MicroDistiller with 10 mL distilled water per sample vial. The sample vial was heated to 108 °C at a rate of 20 °C/min and kept at this temperature for 90 min, then heated to 112 °C at a rate of 20 °C/min and kept at this temperature for 30 min. The sample was subjected to a final postrun for 2 min under the same conditions. The collecting vial, containing a solution of NaCl (2.5 g, Sigma-Aldrich) and water (0.5 mL, ultrapure) plus 350 µL of *n*-hexane [Sigma-Aldrich, ≥99% (GC)] to trap volatile components, was cooled to –5 °C during distillation. After the distillation was completed, the organic layer in the collection vial was separated and analyzed by gas chromatography (GC) and gas chromatography-mass spectrometry (GC-MS) systems, simultaneously.

2.3. GC analysis

The GC analysis was carried out using an Agilent 6890N GC system. FID detector temperature was 300 °C. To obtain the same elution order with GC-MS, simultaneous autoinjection was done on a duplicate of the same column applying the same operational conditions. Relative percentage amounts of the separated compounds were calculated from FID chromatograms. The results of the analysis are shown in Table 2.

2.4. GC-MS analysis

The GC-MS analysis was carried out with an Agilent 5975 GC-MSD system. Innowax FSC column (60 m × 0.25 mm, 0.25 mm film thickness) was used with helium as carrier gas (0.8 mL/min). GC oven temperature was kept at 60 °C for 10 min and programmed to 220 °C at a rate of 4 °C/min, and kept constant at 220 °C for 10 min and then programmed to 240 °C at a rate of 1 °C/min. Split ratio was adjusted at 40:1. The injector temperature was set at 250 °C. Mass spectra were recorded at 70 eV. Mass range was from *m/z* 35 to 450.

2.5. Identification of components

Identification of volatile compounds was carried out by comparison of their relative retention times with those of authentic samples or by comparison of their relative retention indices (RRI) to series of *n*-alkanes (C₈ to C₂₅). Computer matching against commercial (Wiley GC-MS Library, Adams Library, MassFinder 3 Library) and in-house “Başer Library of Essential

Table 1. Data on GPS and locations of the plant materials

| Taxon | Location | Coordinate | Altitude |
|--|-----------|-----------------------|----------|
| <i>T. sipyleus</i> subsp. <i>sipyleus</i> var. <i>sipyleus</i> | Elmalı | N36 43.581 E29 43.531 | 1599 m |
| <i>T. sipyleus</i> subsp. <i>sipyleus</i> var. <i>davisianus</i> | Saklıkent | N36 49.921 E30 19.600 | 2023 m |
| <i>T. sipyleus</i> subsp. <i>sipyleus</i> var. <i>rosulans</i> | Gazipaşa | N36 25.167 E32 33.113 | 2005 m |

Table 2. Volatile compounds of *Thymus sipyleus* subspecies.

| RR ^{1a} | RRI ^b | Compound | <i>T. sipyleus</i> subsp. <i>rosulans</i> % | | | <i>T. sipyleus</i> subsp. <i>sipyleus</i> var. <i>davisiianus</i> % | | | <i>T. sipyleus</i> subsp. <i>sipyleus</i> var. <i>sipyleus</i> % | | | IM |
|------------------|--|---|---|-------------|-------------|---|-------------|-----|--|-----|-------------|---------------------|
| | | | A | A | A | B | B | B | C | C | C | |
| | | | 199 | 200 | 201 | 196 | 197 | 198 | 190 | 191 | 192 | |
| 1000 | 1000 ^c | Decane | 0.2 | - | - | 0.1 | - | - | - | - | - | t _R , MS |
| 1014 | 998-1029 ^d | Tricyclene | - | 0.1 | - | tr | - | 0.1 | 0.1 | 0.1 | 0.2 | MS |
| 1032 | 1008-1039 ^d | α-Pinene | 1.4 | 2.2 | 18.4 | 0.9 | 2.1 | 1.4 | 1.5 | 1.4 | 2.0 | t _R , MS |
| 1035 | 1012-1039 ^d | α-Thujene | - | - | - | - | 2.6 | - | 1.5 | 1.0 | 1.0 | MS |
| 1076 | 1043-1086 ^d | Camphene | 0.9 | 3.4 | 1.7 | 1.7 | - | 3.4 | 4.5 | 3.6 | 7.2 | t _R , MS |
| 1118 | 1085-1130 ^d | β-Pinene | 0.8 | 1.0 | 0.9 | 0.2 | 4.4 | 0.9 | 1.0 | 0.3 | 0.8 | t _R , MS |
| 1132 | 1098-1140 ^d | Sabinene | 0.4 | 0.9 | - | - | 2.8 | 0.5 | 0.3 | 0.2 | 0.3 | t _R , MS |
| 1136 | 1109-1137 ^d | Thuja-2,4(10)-diene | - | - | 1.5 | - | - | - | - | - | - | MS |
| 1174 | 1140-1175 ^d | Myrcene | 0.8 | 0.7 | 1.9 | 0.9 | 1.0 | 4.9 | 1.5 | 3.4 | 1.0 | t _R , MS |
| 1188 | 1154-1195 ^d | α-Terpinene | 0.3 | - | - | - | 0.9 | - | 1.6 | 0.8 | 0.3 | t _R , MS |
| 1195 | 1167-1197 ^d | Dehydro 1,8-cineole | - | 0.1 | - | - | - | - | - | 0.1 | - | t _R , MS |
| 1203 | 1178-1219 ^d | Limonene | 2.9 | 1.8 | 3.9 | tr | 1.5 | 1.7 | 0.7 | 0.8 | 0.7 | t _R , MS |
| 1213 | 1186-1231 ^d | 1,8-Cineole | 8.7 | 11.6 | 2.0 | 0.9 | 31.1 | 7.4 | 11.2 | - | 11.6 | t _R , MS |
| 1215 | 1215 ^e | <i>p</i> -Mentha-1,3,6-triene | - | - | 0.9 | 0.4 | - | - | - | - | - | MS |
| 1218 | 1188-1233 ^d | β-Phellandrene | - | - | - | - | - | - | - | 0.2 | - | t _R , MS |
| 1244 | 1213-1249 ^d | 2-Pentyl furan | - | - | - | tr | - | - | - | - | - | MS |
| 1246 | 1211-1251 ^d | (<i>Z</i>)-β-Ocimene | - | 0.3 | 0.5 | - | - | - | 0.1 | - | - | t _R , MS |
| 1255 | 1222-1266 ^d | γ-Terpinene | 0.7 | 0.2 | tr | - | 1.9 | 0.2 | 7.5 | 4.2 | 2.4 | t _R , MS |
| 1266 | 1232-1267 ^d | (<i>E</i>)-β-Ocimene | 1.2 | 3.1 | 1.8 | - | 1.8 | 0.6 | 2.2 | - | - | t _R , MS |
| 1267 | 1230-1280 ^d | 3-Octanone | 1.0 | 1.8 | 1.0 | 0.8 | - | 1.5 | - | 0.4 | 1.5 | t _R , MS |
| 1280 | 1246-1291 ^d | <i>p</i>-Cymene | 1.1 | 0.4 | 2.4 | 0.7 | 12.4 | 0.5 | 21.8 | 8.7 | 9.2 | t _R , MS |
| 1290 | 1261-1300 ^d | Terpinolene | - | 0.2 | - | - | tr | 0.2 | 0.3 | - | tr | t _R , MS |
| 1296 | 1267-1312 ^d | Octanal | - | - | - | tr | - | - | - | - | - | t _R , MS |
| 1348 | 1317-1357 ^d | 6-methyl-5-hepten-2-one | - | - | - | 0.9 | - | 0.3 | - | - | - | MS |
| 1382 | 1334-1394 ^d | <i>cis</i> -Alloocimene | - | - | - | - | - | - | 0.1 | - | - | MS |
| 1393 | 1372-1408 ^d | 3-Octanol | 0.4 | 0.7 | - | 0.4 | 0.5 | 1.1 | 0.1 | 0.8 | 5.8 | MS |
| 1400 | 1370-1414 ^d | Nonanal | 0.9 | 0.1 | - | 0.3 | 1.1 | 0.8 | tr | tr | - | MS |
| 1413 | 1413 ^e | Rosefuran | - | - | - | 1.1 | - | | - | - | - | - |
| 1429 | 1405-1431 ^d | Perillene | - | - | - | 0.4 | - | 0.2 | - | - | - | MS |
| 1449 | 1412-1457 ^d | <i>p</i> -Cymenene | - | - | - | tr | - | - | tr | - | - | MS |
| 1452 | 1411-1465 ^d | 1-Octen-3-ol | 1.0 | 1.2 | 1.7 | 2.8 | 0.8 | 1.3 | 1.9 | 0.5 | 0.7 | t _R , MS |
| 1460 | 1460 ^f | 2,6-Dimethyl-1,3(<i>E</i>), 5(<i>E</i>)-7-octatetraene | - | 0.2 | - | - | - | - | tr | - | - | MS |
| 1461 | 1463 ⁿ | (<i>E</i>)-2-hexenyl butyrate | - | - | - | - | - | - | tr | - | tr | MS |
| 1466 | 1438-1480 ^d | α-Cubebene | - | - | - | - | - | - | - | - | 0.4 | MS |
| 1474 | 1425-1478 ^d 1474 ^f | <i>trans</i> -Sabinene hydrate | 0.9 | tr | - | - | 1.1 | - | 1.5 | 0.5 | - | MS |
| 1478 | 1478 ^f 1479 ^h | <i>cis</i> -Linalool oxide (fur.) | - | - | - | - | - | - | tr | - | - | MS |
| 1493 | 1459-1500 ^d | α-Ylangene | - | - | - | - | - | - | tr | - | - | MS |

Table 2. (Continued).

| | | | | | | | | | | | | |
|------|--|---|-------------|-------------|-----|-------------|-------------|-------------|-----|-------------|-----|---------------------|
| 1495 | 1452-1513 ^d | 2-Ethyl hexanol | 0.3 | tr | - | - | 0.4 | tr | - | - | - | MS |
| 1496 | 1471-1495 ^d | Bicycloelemene | 0.5 | 0.2 | 0.5 | - | 0.9 | - | 0.4 | 0.8 | 1.3 | MS |
| 1496 | 1495 ^h | <i>cis, cis</i> - Photocitral | - | - | - | 0.7 | - | - | - | - | - | MS |
| 1497 | 1462-1522 ^d | α -Copaene | - | tr | 0.3 | - | - | - | 0.1 | - | - | MS |
| 1519 | 1519 ^h | <i>trans,trans</i> - Photocitral | - | - | - | 1.4 | - | 0.7 | - | - | - | MS |
| 1532 | 1481-1537 ^d | Camphor | 1.5 | 0.1 | 2.6 | 0.2 | - | - | 0.1 | tr | 8.3 | t _R , MS |
| 1535 | 1496-1546 ^d | β -Bourbonene | 1.4 | 0.3 | 4.5 | 0.6 | 0.3 | 0.3 | 0.1 | 0.1 | 0.3 | t _R , MS |
| 1549 | 1518-1560 ^d | β -Cubebene | - | - | 0.3 | - | - | - | - | - | - | MS |
| 1553 | 1507-1564 ^d | Linalool | 0.7 | 0.2 | 0.3 | 0.7 | 6.4 | 4.7 | 0.3 | 0.5 | 0.4 | t _R , MS |
| 1555 | 1557 ^g | 1-Nonen-3-ol | - | - | - | - | - | - | 0.2 | 0.1 | 0.1 | MS |
| 1556 | 1526-1565 ^d 1556 ^f | <i>cis</i> -Sabinene hydrate | - | - | - | - | - | - | 0.3 | 0.2 | 0.2 | MS |
| 1562 | 1519-1574 ^d | Octanol | - | - | - | - | 0.2 | - | tr | - | - | t _R , MS |
| 1571 | 1557-1625 ^d | <i>trans-p</i> -Menth-2-en-1-ol | - | - | - | - | - | - | 0.1 | - | - | MS |
| 1588 | 1588-1610 ^d | Bornyl formate | - | 0.1 | - | - | - | - | - | - | - | MS |
| 1589 | 1547-1589 ^d | β -Ylangene | 0.6 | - | - | - | - | - | - | - | - | MS |
| 1590 | 1549-1597 ^d | Bornyl acetate | - | 1.2 | - | 0.9 | - | 2.0 | 1.6 | 2.0 | 2.5 | t _R , MS |
| 1600 | 1565-1608 ^d | β -Elemene | 1.0 | tr | 0.7 | - | - | - | - | - | - | MS |
| 1611 | 1564-1630 ^d | Terpinen-4-ol | 1.3 | 0.3 | - | 0.5 | 3.3 | 0.3 | 2.5 | 0.8 | 0.9 | t _R , MS |
| 1612 | 1569-1632 ^d | β-Caryophyllene | 14.2 | 3.0 | 8.9 | 5.1 | 14.6 | 8.2 | 7.1 | 2.3 | 5.0 | t _R , MS |
| 1624 | 1600-1650 ^d | <i>trans</i> -Dihydrocarvone | - | - | - | - | - | - | - | 0.4 | 0.2 | t _R , MS |
| 1628 | 1583-1668 ^d | Aromadendrene | - | - | - | - | - | - | 0.4 | 0.2 | 0.5 | MS |
| 1645 | 1645 ^h | <i>cis</i> -Dihydrocarvone | - | - | - | - | - | - | - | 0.1 | - | t _R , MS |
| 1663 | 1647-1668 ^d | <i>cis</i> -Verbenol | - | - | 2.7 | - | - | - | - | - | - | MS |
| 1661 | 1624-1668 ^d | Alloaromadendrene | - | 0.1 | - | - | - | - | 0.1 | - | 0.1 | MS |
| 1668 | 1627-1668 ^d | (<i>Z</i>)- β -Farnesene | - | 0.5 | - | 0.7 | - | - | - | 0.1 | 0.4 | MS |
| 1683 | 1665-1691 ^d | <i>trans</i> -Verbenol | - | - | 2.8 | - | - | - | - | - | - | MS |
| 1687 | 1637-1689 ^d | α -Humulene | 3.7 | 0.6 | 0.9 | 0.3 | tr | 0.4 | 0.7 | 0.2 | tr | t _R , MS |
| 1694 | 1641-1706 ^d | Neral | 0.5 | - | - | 19.6 | - | 6.6 | - | 0.1 | - | MS |
| 1704 | 1655-1714 ^d | γ -Muuroolene | - | - | 0.5 | - | - | - | - | - | - | MS |
| 1706 | 1659-1724 ^d | α-Terpineol | 0.7 | 35.0 | - | 0.3 | 1.3 | 19.8 | 3.2 | 35.8 | 0.8 | t _R , MS |
| 1708 | 1708 ^{c,h} 1707 ^f | Ledene | - | - | - | - | - | - | 0.2 | 0.1 | 0.3 | MS |
| 1719 | 1653-1728 ^d | Borneol | - | 0.5 | - | 2.8 | 0.5 | 4.0 | 7.6 | 2.8 | 4.9 | t _R , MS |
| 1725 | 1696-1735 ^d | Verbenone | - | - | 0.7 | - | - | - | - | - | - | t _R , MS |
| 1726 | 1676-1726 ^d | Germacrene D | 0.8 | 0.4 | 4.8 | - | 0.3 | 0.3 | 0.3 | 0.3 | 0.5 | MS |
| 1730 | 1730 ^g | δ -Guaiene | 0.7 | - | - | - | - | - | - | - | - | MS |
| 1732 | 1732 ^m | Bicyclosquiphellandrene | - | - | 0.5 | - | - | - | - | - | tr | MS |
| 1740 | 1698-1748 ^d | β -Bisabolene | - | - | - | - | - | - | - | 0.8 | 2.1 | t _R , MS |
| 1740 | 1686-1753 ^d | α -Muuroolene | - | - | - | - | - | - | 0.1 | - | - | MS |
| 1741 | 1680-1750 ^d | Geranial | 0.8 | - | - | 30.3 | - | 11.1 | - | 0.4 | - | MS |
| 1744 | 1696-1748 ^d | α -Selinene | - | - | - | - | - | - | 0.1 | - | - | MS |
| 1746 | 1744 ^m | Selina-4(15),7(11)-diene | 0.8 | - | - | - | - | - | - | - | - | MS |

Table 2. (Continued).

| | | | | | | | | | | | | |
|------|---|-----------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|---------------------|
| 1755 | 1692-1757 ^d | Bicyclogermacrene | 0.6 | - | 0.3 | - | 0.1 | 0.4 | 0.5 | 1.0 | 1.5 | MS |
| 1772 | 1734-1789 ^d | Citronellol | - | - | - | 1.4 | - | 0.5 | - | - | - | t _R , MS |
| 1773 | 1722-1774 ^d | δ-Cadinene | - | - | 0.5 | - | - | - | 0.3 | 0.1 | 0.4 | t _R , MS |
| 1776 | 1735-1782 ^d | γ-Cadinene | - | 2.7 | 0.2 | - | - | - | 0.5 | 0.1 | 0.3 | MS |
| 1797 | 1787 ^m | Aromadendra-1(10), 4(15)-diene | - | - | - | - | - | - | - | - | tr | MS |
| 1808 | 1752-1832 ^d | Nerol | - | - | - | - | - | tr | - | - | - | t _R , MS |
| 1816 | 1734-1803 ^d | α-Cadinene | - | tr | - | - | - | - | - | - | - | MS |
| 1853 | 1800-1853 ^d | cis-Calamenene | - | 0.4 | - | - | - | - | - | tr | tr | MS |
| 1857 | 1795-1865 ^d | Geraniol | - | - | - | tr | - | - | - | - | - | t _R , MS |
| 1864 | 1813-1865 ^d | p-Cymen-8-ol | - | - | - | - | - | - | - | tr | - | MS |
| 1868 | 1868 ^{c, g, h} | (E)-Geranyl acetone | - | - | - | tr | - | - | - | - | - | t _R , MS |
| 1900 | 1900 ^p | Nonadecane | - | - | - | - | - | - | tr | - | - | t _R , MS |
| 2008 | 1936-2023 ^d | Caryophyllene oxide | 6.2 | 3.1 | 4.2 | 6.2 | - | 3.5 | 0.7 | 0.5 | 0.6 | t _R , MS |
| 2029 | 1963-2029 ^d | Perilla alcohol | - | - | - | 1.2 | - | - | - | - | - | MS |
| 2037 | 2016-2043 ^d | Salvial-4(14)-en-1-one | 0.3 | tr | 0.8 | - | - | - | - | - | - | MS |
| 2069 | 2000-2070 ^d | Germacrene D-4-β-ol | - | - | - | - | - | - | - | - | tr | MS |
| 2071 | 2003-2071 ^d | Humulene epoxide II | 1.4 | 0.5 | - | - | - | - | 0.1 | tr | - | MS |
| 2080 | 2019-2090 ^d | Cubenol | - | 1.1 | - | - | - | - | 0.1 | tr | tr | MS |
| 2096 | 2043-2103 ^d | Elemol | - | - | - | - | - | - | tr | - | - | MS |
| 2098 | 2049-2104 ^d | Globulol | - | - | - | - | - | 0.1 | - | - | tr | MS |
| 2123 | 2123 ^e 2130 ^k | Salviadienol | - | 0.2 | 1.1 | - | - | - | - | - | - | MS |
| 2144 | 2074-2150 ^d | Spathulenol | 7.0 | 4.4 | 6.3 | 0.3 | 0.7 | 2.7 | 0.8 | 1.2 | 2.0 | t _R , MS |
| 2170 | 2090-2189 ^d | b-Bisabolol | - | 0.2 | - | - | - | - | - | - | - | MS |
| 2187 | 2136-2200 ^d | T-Cadinol | - | 9.4 | - | 0.3 | - | - | 0.4 | 0.1 | 0.3 | MS |
| 2198 | 2100-2205 ^d | Thymol | tr | - | - | - | 0.1 | - | 0.8 | 0.9 | 0.4 | t _R , MS |
| 2219 | 2211 ^k | Clovenol | - | - | - | - | - | tr | - | - | - | t _R , MS |
| 2239 | 2140-2246 ^d | Carvacrol | 0.7 | tr | 0.2 | - | 0.9 | tr | 9.1 | 20.5 | 18.2 | t _R , MS |
| 2243 | 2243 ^e | Torilenol | 0.5 | 0.3 | 1.1 | - | - | - | - | - | - | MS |
| 2247 | 2247 ^{f, g} | trans-α-Bergamotol | 0.6 | 0.2 | 0.4 | - | - | 0.2 | - | tr | 0.1 | MS |
| 2255 | 2180-2255 ^d | α-Cadinol | 0.3 | 0.3 | 0.3 | 0.4 | - | - | - | tr | 0.1 | t _R , MS |
| 2264 | 2218-2264 ^d | Intermedeol | 13.2 | - | - | - | - | - | - | - | - | MS |
| 2316 | 2316 ^g | Caryophylladienol I | 0.7 | 0.5 | 0.8 | tr | - | 0.2 | tr | - | - | MS |
| 2324 | 2324 ^{c, f, g} | Caryophylladienol II | 1.4 | - | - | 1.6 | - | 1.3 | 0.1 | 0.1 | - | MS |
| 2369 | 2351-2402 ^d | Eudesma-4(15),7-diene- 1-β-ol | 1.1 | 0.3 | 1.7 | - | - | - | - | - | - | MS |
| 2389 | 2389 ^{g, h} | Caryophyllenol I | 1.3 | 0.6 | 0.6 | 1.5 | - | 1.0 | 0.1 | 0.1 | - | MS |
| 2392 | 2392 ^{c, d, f, g} -2396 ^d | Caryophyllenol II | 1.6 | 0.7 | 1.0 | 1.0 | - | 0.9 | 0.1 | 0.1 | - | MS |
| | | Total | 90.0 | 97.4 | 84.3 | 90.5 | 96.0 | 96.2 | 98.5 | 99.0 | 97.8 | |

RRI^a: relative retention indices calculated against *n*-alkanes (C₈ to C₂₅). %: calculated from the FID chromatograms. RRI^b: RRI from literature (c [28], d [29], e [30], f [31], g [32], h [33], k [34], m [35], n [36], p [37]) for polar column values. tr: trace (<0.1 %). IM: identification method. t_R: identification based on the retention times (t_R) of genuine compounds on the HP Innowax column. MS: identified on the basis of computer matching of the mass spectra with those of the in-house Baser Library of Essential Oil Constituents, Adams, MassFinder and Wiley libraries and comparison with literature data.

Oil Constituents” built up by genuine compounds and components of known oils, as well as MS literature data were used for the identification [20].

3. Results and discussion

GC and GC-MS analysis of the samples obtained by microdistillation resulted in a total of one hundred fifteen volatile compounds were identified in *Thymus sipyleus* subspecies by using in house and commercial libraries. The elution of the compounds in the microdistilled oils was done by using an HP-Innowax FSC column. Table 2 shows the list of detected and identified volatile constituents with their RRI and relative percentages in the samples.

Seventy seven total components of three *T. sipyleus* subsp. *rosulans* samples were identified by GC-MS. Forty-eight components of the KT199 sample were detected representing 90% of the oil. β -Caryophyllene (14.2%) and intermedeol (13.3%) were the major compounds of this sample. Twenty one volatiles are higher than 1% and other major compounds are 1,8-cineole (8.7%), caryophyllene oxide (6.2%), spathulenol (7.0%), α -humulene (3.7%), limonene (2.9%).

Fifty-four components of the KT200 sample were identified representing 97.4% of the oil. α -Terpineol (35%) and 1,8-cineole (11.6%) were the major compounds and sixteen volatiles are higher than 1%. Other major compounds are T-cadinol (9.4%), spathulenol (4.4%), camphene (3.4%), (*E*)- β -ocimene (3.1%), caryophyllene oxide (3.1%), β -caryophyllene (3.0%), γ -cadinene (2.7%), α -pinene (2.2%).

Forty-three compounds of the KT201 sample were detected representing 84.3% of the oil and α -pinene (18.4%) and β -caryophyllene (8.9%) were the major components. Spathulenol (6.3%), germacrene D (4.8%), β -bourbonene (4.5%), caryophyllene oxide (4.2%), limonene (3.9%), *trans*-verbenol (2.9%), *cis*-verbenol (2.7%) were the other major volatiles.

A previous study reported that the essential oil of aerial parts at the flowering stage of *T. sipyleus* subsp. *sipyleus* var. *rosulans* collected from İspir, Erzurum was characterized with carvacrol (30.0%), thymol (14.5%), *p*-cymene, α -terpinyl acetate and linalool as main components [21]. Akçin (2008) published that the volatile constituents of the oil of *T. sipyleus* subsp. *rosulans* samples collected from different regions showed significant differences. In the essential oil of Kastamonu sample, higher levels of myrcene (5.2%), 1,8-cineole (16.6%) were found while germacrene D-4-ol (8.2%), α -cadinol (6.4%), germacrene D (5.21%), (*Z*)- β -farnesene (4.4%) and bicyclogermacrene (4.0%) in the samples from Çorum. In general, β -caryophyllene (6.8-14.2%), linalool (0.1-22.5%), 1,8-cineole (0.1-16.6%), α -terpineol (2.2-7.0%), caryophyllene oxide (1.9-8.1%), germacrene D (1.4-5.2%) and spathulenol (2.1-4.8%) were detected as major compounds in the samples [22]. Tepe et al. (2005) reported that 47 constituents were identified representing 98.7% of the oil of *Thymus sipyleus* subsp. *sipyleus* var. *rosulans* at flowering stage collected from Kangal, Sivas. This oil is characterised by the high monoterpene fraction (94.0%) and carvacrol (58.1%), thymol (20.5%) and *p*-cymene (4.1%) and γ -terpinene (4.4%) as main constituents [23].

Sixty-four total components of three *T. sipyleus* subsp. *sipyleus* var. *davisianus* samples were identified. Forty-six components of the KT196 sample were detected representing 90.5% of the oil, geranial (30.3%) and neral (19.6%) were the major compounds. Fourteen volatiles are higher than 1% and other notable components are caryophyllene oxide (6.2%), β -caryophyllene (5.1%), borneol (2.8%), 1-octen-3-ol (2.8%).

Thirty-one components of the KT197 sample were identified representing 96.0% of the oil. 1,8-cineole (31.1%) and β -caryophyllene (14.6%) were the major components. Sixteen volatiles are higher than 1% and other major compounds are *p*-cymene (12.4%), β -pinene (4.4%), linalool (6.4%), terpinen-4-ol (3.3%), sabinene (2.8%), α -thujene (2.6%) and α -pinene (2.1%).

Forty five components of the KT198 sample were detected representing 96.2% of the oil. The major compounds are α -terpineol (19.8%) and geranial (11.1%). Other major volatiles are β -caryophyllene (8.2%), 1,8-cineole (7.4%), neral (6.6%), myrcene (4.9%), linalool (4.7%), borneol (4.0%) and caryophyllene oxide (3.5%). Contents of nineteen compounds are higher than 1%.

In a previous study, Meriçli and Tanker (1986) reported that the essential oil of *T. sipyleus* subsp. *sipyleus* var. *davisianus* collected from Tefenni was rich in geranial (32.1%) [24]. The essential oil of aerial parts at full flowering stage of *T. sipyleus* subsp. *sipyleus* var. *davisianus* collected from Uşak was characterized with thymol (38.3%) and carvacrol (37.9%) among identified fourteen constituents [25].

Totally eighty volatile compounds of three *T. sipyleus* subsp. *sipyleus* var. *sipyleus* samples were identified by GC and GC-MS systems. Sixty-one volatile compounds of the KT190 sample were identified representing 98.5% of the oil. The major compounds are *p*-cymene (21.8%) and 1,8-cineole (11.2%). Eighteen volatiles are higher than 1% and other major compounds are carvacrol (9.1%), γ -terpinene (7.5%), borneol (7.6%), β -caryophyllene (7.1%) and camphene (4.5%).

Fifty-six volatiles of the KT191 sample were detected representing 99.0% of the oil, α -terpineol (35.8%) and carvacrol (20.5%) were the main compounds. Thirteen volatiles are higher than 1% and other major compounds are *p*-cymene (8.7%), γ -terpinene (4.2%), camphene (3.6%), myrcene (3.4%).

Fifty-four volatiles of the KT192 sample were identified representing 97.8% of the oil and the major compounds are carvacrol (18.2%) and 1,8-cineole (11.6%). The contents of fifteen volatiles are higher than 1% and other major compounds are *p*-cymene (9.2%), camphor (8.3%), camphene (7.2%), 3-octanol (5.8%), β -caryophyllene (5.0%), borneol (4.9%).

Demirci et al. (2018) reported that the essential oil of air dried and crushed aerial parts of *T. sipyleus* subsp. *sipyleus* var. *sipyleus* collected from Ulaş, Sivas, was characterized by high amount of thymol (66.2%), followed by *p*-cymene (9.4%), and γ -terpinene (9.2%) [11]. In another study, the chemical composition of *T. sipyleus* subsp. *sipyleus* var. *sipyleus* essential oil which originated from different regions (Denizli, Afyon, Ankara, Muğla, Konya) contained geranial (8.4%–37.0%), neral (3.1%–25.6%), linalool (21.8%), and α -terpineol+isoborneol (25.5%) as main components [26]. In a study published by Tepe et al. (2005), the aerial parts of *T. sipyleus* subsp. *sipyleus* var. *sipyleus* collected from Düziçi, Osmaniye were subjected to water distillation. Seventy-one volatile compounds were identified representing 92.5% of the total oil. The major compounds were borneol (11.2%), α -muurolol (9.2%), β -caryophyllene (7.6%), geranial (7.3%) and neral (5.4%) [23]. Pekgözlü and Özcan (2018) found citronellol as major compound in the SDE sample of *T. sipyleus* var. *sipyleus* leaves collected from Büğdüz, Burdur [27].

To sum up, published studies and our present study have generally shown a great deal of variability and diversity. *Thymus sipyleus* subsp. *sipyleus* var. *sipyleus* samples collected from three regions of Elmalı were characterized with different major compounds such as 1,8-cineole, *p*-cymene, α -terpineol and carvacrol. The major volatile compounds in *T. sipyleus* subsp. *sipyleus* var. *davisanus* samples (three different sites of Saklıkent) were identified as 1,8-cineole, *p*-cymene, β -caryophyllene, geranial, and α -terpineol with different percentage amounts. *T. sipyleus* subsp. *rosulans* (three different sites of Gazipaşa) samples with major constituents as α -pinene, 1,8-cineole, β -caryophyllene, α -terpineol were identified.

4. Conclusion

Thymol is the major compound of most *Thymus* species. According to published data and our present study, chemical polymorphism has been found among the *Thymus sipyleus* subspecies even though the samples were collected from the same region. *Thymus* populations collected from Turkey have a greater variation of the major components in volatile oils. The variation of volatile oil composition has great importance due to its uses as food and in food processes. The results obtained here suggest that the growing conditions of thyme may alter the volatile oil content and composition.

References

1. Celep F, Dirmenci T. Systematic and biogeographic overview of Lamiaceae in Turkey. *Natural Volatiles and Essential oils* 2017; 4 (4): 14-27.
2. Stahl-Biskup E, Sáez F. *The Genus Thymus*, London and New York: Taylor & Francis, 2002.
3. Tümen G, Kirimer N, Başer KHC. Composition of the essential oils of *Thymus* species growing in Turkey. *Chemistry of Natural Compounds* 1995; 31 (1): 42-47. doi: 0009-3130/95/3101-0042\$12.50
4. Meriçli F. Volatile oils of *Thymus kotschyanus* var. *glabrescens* and *T. fedtschenkoi* var. *handeli*. *Journal of Natural Products* 1986; 49: 942-943. doi: 10.1021/np50047a039
5. Çingi I, Kirimer N, Sarıkardaşoğlu I, Baser KHC. Pharmacological effects of the essential oil of *Origanum onites* and *O. minutiflorum*, In: *Proceedings of the Symposium on Plant Drugs*; Eskisehir, Turkey; 1991, pp. 99-100.
6. Van-Den Broucke CO, Lemli, JA. Pharmacological and chemical investigation of Thyme liquid extracts. *Planta Medica* 1981; 41: 129-135. doi: 10.1055/s-2007-971689
7. Figueiredo AC, Barroso JG, Pedro LG, Salgueiro L, Miguel MG et al. Portuguese *Thymbra* and *Thymus* species volatiles: chemical composition and biological activities. *Current Pharmaceutical Design* 2008; 14: 3120-3140. doi: 10.2174/138161208786404218
8. European Pharmacopoeia 9.0. European Directorate for the Quality of Medicines & HealthCare (EDQM) of the Council of Europe. Volume 1. Strasbourg, France: pp.1538-1541.
9. European Medicine Agency (EMA). Assessment report on *Thymus vulgaris* L., *Thymus zygis* L., aetheroleum. Revision 1. Committee on Herbal Medicinal Products (HMPC), EMA/HMPC/52980/2017. Amsterdam, Netherlands: 8 July 2020.
10. WHO monographs on selected medicinal plants. Volume 1. World Health Organization, Geneva, Switzerland: 1999. pp 259-266.
11. Demirci F, Karaca N, Tekin M, Demirci B. Anti-inflammatory and antibacterial evaluation of *Thymus sipyleus* Boiss. subsp. *sipyleus* var. *sipyleus* essential oil against rhinosinusitis pathogens. *Microbial Pathogenesis* 2018; 122: 117-121. doi: 10.1016/j.micpath.2018.06.025
12. Güneş S, Savran A, Paksoy MY, Koşar M, Çakılciöğlü U. Ethnopharmacological survey of medicinal plants in Karaisalı and its surrounding (Adana-Turkey). *Journal of Herbal Medicine* 2017; 8: 68-75. doi: 10.1016/j.hermed.2017.04.002

13. Koyuncu O, Yaylacı ÖK, Öztürk D, Potoğlu Erkara İ, Savaroğlu F et al. Risk categories and ethnobotanical features of the Lamiaceae taxa growing naturally in Osmaniye (Bilecik/Turkey) and environs. *Biological Diversity and Conservation* 2010; 3 (3): 31-45.
14. Özüdoğru B, Akaydın G, Erik S, Yesilada E. Inferences from an ethnobotanical field expedition in the selected locations of Sivas and Yozgat provinces (Turkey). *Journal of Ethnopharmacology* 2011; 137: 85-98. doi: 10.1016/j.jep.2011.04.050
15. Paksoy MY, Selvi S, Savran A. Ethnopharmacological survey of medicinal plants in Ulukışla (Nigde-Turkey). *Journal of Herbal Medicine* 2016; 6 (1): 1-7. doi: 10.1016/j.hermed.2015.04.003
16. Altundag E, Ozturk M. Ethnomedicinal studies on the plant resources of east Anatolia, Turkey. *Procedia Social and Behavioral Sciences* 2011; 19: 756-777. doi: 10.1016/j.sbspro.2011.05.195
17. Özcan S, Toprak G, Torun C, Vural C. *Thymus sipyleus* Boiss subsp. *rosulans* (Borbas) J. Jalas'ın organik ekstrakt ve uçucu yağlarının antimikrobiyal aktivitesi. *Biyoloji Bilimleri Araştırma Dergisi* 2008; 1 (2): 17-22.
18. Kotan R, Dadasoglu F, Kordali S, Cakır A, Dikbas N et al. Antibacterial activity of essential oils extracted from some medicinal plants, carvacrol and thymol on *Xanthomonas axonopodis* pv. *vesicatoria* (Doidge) dye causes bacterial spot disease on pepper and tomato. *Journal of Agricultural Technology* 2007; 3 (2): 299-306.
19. Akçin TA. Numerical taxonomic studies on some species of the genus *Thymus* L. (Labiatae) in Turkey. *Asian Journal of Plant Sciences* 2006; 5 (5):782-788. doi: 10.3923/ajps.2006.782.788
20. Agalar HG, Temiz B, Demirci B, Baser KHC. Drying effects on the volatile compounds of kumquat, limequat and Mexican lime fruits. *Journal of Essential Oil Bearing Plants* 2020; 23 (6): 1395-1408. doi: 10.1080/0972060X.2020.1868347
21. Cetin B, Cakmaci S, Gurses M. Anti-probiotic effects of essential oils from some Turkish endemic thyme species. *Asian Journal of Chemistry* 2013; 25 (15): 8625-8628. doi: 10.14233/ajchem.2013.14868
22. Akçin TA. Volatile oil variability in *Thymus sipyleus* Boiss. subsp. *rosulans* (Borbas) Jalas populations collected in Turkey. *Journal of Essential Oil Bearing Plants* 2008; 11 (3): 214-223. doi: 10.1080/0972060X.2008.10643623
23. Tepe B, Sokmen M, Akpulat HA, Daferera D, Polissiou M et al. Antioxidative activity of the essential oils of *Thymus sipyleus* subsp. *sipyleus* var. *sipyleus* and *Thymus sipyleus* subsp. *sipyleus* var. *rosulans*. *Journal of Food Engineering* 2005; 66 (4): 447-454. doi: 10.1016/j.jfoodeng.2004.04.015
24. Meriçli İF, Tanker M. The volatile oils of some endemic *Thymus* species growing in Southern Anatolia. *Planta Medica* 1986; 52 (4): 340. doi: 10.1055/s-2007-969176
25. Ceylan A, Uğur A. Chemical composition and anti-biofilm activity of *Thymus sipyleus* Boiss. subsp. *sipyleus* Boiss. var. *davisanus* Ronniger essential oil. *Archives of Pharmacal Research* 2015; 38 (6): 957-965. doi: 10.1007/s12272-014-0516-0
26. Baser KHC, Kürkçüoğlu M, Özek T, Tümen G, Akgül A. Essential oil of *Thymus sipyleus* Boiss. subsp. *sipyleus* var. *sipyleus*. *Journal of Essential Oil Research* 1995; 7 (4): 411-413.
27. Pekgözlü Kılıç A, Özcan K. Effect of drying methods on the composition of volatile compounds of Thyme (*Thymus sipyleus* Boiss. var. *sipyleus*). *Journal of Bartın Faculty of Forestry* 2018; 20 (2): 210-215. doi: 10.24011/barofd.425039
28. Tabanca N, Demirci F, Ozek G, Tumen G, Baser KHC. Composition and antimicrobial activity of the essential oil of *Origanum × dolichosiphon* P.H. Davis. *Chemistry of Natural Compounds* 2001; 37 (3): 238-241. doi: 10.1023/A:1012513922871
29. Babushok VI, Linstrom PJ, Zenkevich IG. Retention indices for frequently reported compounds of plant essential oils. *Journal of Physical and Chemical Reference Data* 2011; 40 (4): 043101. doi: 10.1063/1.3653552
30. Kürkçüoğlu M, Işcan G, Demirci F, Baser KHC, Malyer H et al. Composition and antibacterial activity of the essential oil of *Ferulago confusa* Velen. *Journal of Essential Oil Research* 2010; 22 (6): 490-492. doi: 10.1080/10412905.2010.9700380
31. Başer KHC, Demirci B, Kirimer N, Satil F, Tümen G. The essential oils of *Thymus migricus* and *T. fedtschenkoi* var. *handelii* from Turkey. *Flavour and Fragrance Journal* 2002; 17 (1): 41-45. doi: 10.1002/ffj.1036
32. Baser KHC, Demirci B, Kirimer N. Compositions of the essential oils of four *Helichrysum* species from Madagascar. *Journal of Essential Oil Research* 2002; 14 (1): 53-55. doi: 10.1080/10412905.2002.9699762
33. Başer KHC, Tümen G, Tabanca N, Demirci F. Composition and antibacterial activity of the essential oils from *Satureja wiedemanniana* (Lallem.) Velen. *Zeitschrift für Naturforschung C* 2001; 56 (9-10): 731-738. doi: 10.1515/znc-2001-9-1011
34. Köse YB, Demirci B, Başer KHC, Yücel E. Composition of the essential oil of three endemic *Centaurea* species from Turkey. *Journal of Essential Oil Research* 2008; 20 (4): 335-338. doi: 10.1080/10412905.2008.9700025
35. Hulley IM, Sadgrove NJ, Tilney PM, Özek G, Yur S et al. Essential oil composition of *Pentzia incana* (Asteraceae), an important natural pasture plant in the Karoo region of South Africa. *African Journal of Range and Forage Science* 2018; 35(2): 137-145. doi:10.2989/10220119.2018.1495265

36. Demirci B, Demirci F, Başer KHC. Composition of the essential oil of *Cotinus coggygia* Scop. from Turkey, Flavour and Fragrance Journal 2003; 18 (1): 43-44. doi: 10.1002/ffj.1149
37. Kürkçüoğlu M, Abdel-Megeed A, Başer KHC. The composition of Taif rose oil. Journal of Essential Oil Research 2013; (25) 5: 364-367. doi: 10.1080/10412905.2013.809322