

G-frames as special frames

Abas ASKARIZADEH, Mohammad Ali DEGHAN*
Department of Mathematics, Vali-e-Asr University, Rafsanjan, Iran

Received: 18.03.2011 • Accepted: 18.07.2011 • Published Online: 17.12.2012 • Printed: 14.01.2013

Abstract: G-frames are generalizations of ordinary frames for Hilbert spaces. In the present paper we study frames, and operators on a special separable Hilbert C^* -module, $B(H, K)$, where H and K are Hilbert spaces, and we prove that every g-frame for H is a frame for $B(H, K)$ and vice versa. Also, we derive some relationships between g-Riesz bases for H and Riesz bases in $B(H, K)$. Similar results for orthogonal bases will be discussed.

Key words: Hilbert C^* -module, Frame, g-Frame, Riesz basis, g-Riesz basis, Orthogonal basis, g-Orthonormal basis

1. Introduction

Frames were first introduced in 1952 by Duffin and Schaeffer [5] in the study of nonharmonic Fourier series. More than thirty years later, Young [17] and Daubechies et al. [4] reintroduced frames and used them as bases in Hilbert spaces, especially $L^2(\mathbb{R})$. Recent research has shown that frame theory has applications in pure [2, 8] and applied mathematics [7], harmonic analysis [3] and even quantum communication [1].

Generalizations of frames have also been used in many applications. The best-known generalizations of frames, called g-frames, were defined by Sun [15]. The class of g-frames includes the class of ordinary frames. Also, frames in Hilbert C^* -modules were extended to unital C^* -algebras by Frank and Larson [6].

For Hilbert spaces H and K , the Banach space $B(H, K)$ of all bounded linear operators from H into K is a Hilbert $B(K)$ -module.

The goal of this paper is to show that a sequence of operators in $B(H, K)$ is a g-frame for H if and only if it is a frame for $B(H, K)$. We then conclude that g-frames are frames. Also, we illustrate some differences between g-orthonormal and g-Riesz bases. We show that the set of Riesz bases in $B(H, K)$ contains the set of g-Riesz bases, but that the sets are not equal. The same relation is true for orthogonal bases and g-orthonormal bases.

The rest of the paper is organized as follows. In Section 2, we review Hilbert C^* -modules and some properties of the operators on $B(H, K)$, which will be used in Section 3. In Section 3, we offer a necessary and sufficient condition for a sequence of operators in $B(H, K)$ to be a g-frame. Also, we study the relations between orthogonal and Riesz bases in $B(H, K)$ considered as a Hilbert C^* -module with g-orthogonal and g-Riesz bases for H .

Throughout the paper, I and \mathbb{C} denote the sets of all integers and all complex numbers, respectively.

*Correspondence: dehghan@vru.ac.ir

2000 AMS Mathematics Subject Classification: Primary 46L99; Secondary 47A05, 42C15, 46H25.

2. Frames in Hilbert C^* -modules and $B(H, K)$

2.1. Review of the Hilbert C^* -modules

Hilbert C^* -modules form a category between the category of Banach spaces and the category of Hilbert spaces. The basic idea was to study modules over C^* -algebras instead of linear spaces and to allow an inner product to take its values in a more general C^* -algebra than that of the complex numbers \mathbb{C} . The structure was used by Kaplansky [9] in 1952 and was investigated in detail by Rieffel [13] and Paschke [12] in 1972–73.

We shall give only a brief introduction to the theory of Hilbert C^* -modules to make our explanations self-contained. For a comprehensive account, readers are referred to the books by Lance [10] and Wegge-Olsen [16].

Let A be a C^* -algebra and H be a (left) A -module. Suppose that the linear structures given on A and H are compatible, i.e., $\lambda(ax) = a(\lambda x)$ for every $\lambda \in \mathbb{C}$, $a \in A$ and $x \in H$. If there exists a mapping $\langle \cdot, \cdot \rangle : H \times H \rightarrow A$ with the properties,

- (i) $\langle x, x \rangle \geq 0$ for every $x \in H$,
- (ii) $\langle x, x \rangle = 0$ if and only if $x = 0$,
- (iii) $\langle x, y \rangle = \langle y, x \rangle^*$ for every $x, y \in H$,
- (iv) $\langle ax, y \rangle = a \langle x, y \rangle$ for every $a \in A$ and $x, y \in H$,
- (v) $\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle$ for every $x, y, z \in H$,

then the pair $\{H, \langle \cdot, \cdot \rangle\}$ is called a (left) pre-Hilbert A -module. The map $\langle \cdot, \cdot \rangle$ is called an A -valued inner product. If the pre-Hilbert A -module $\{H, \langle \cdot, \cdot \rangle\}$ is complete with respect to the norm $\|x\| = \|\langle x, x \rangle\|^{\frac{1}{2}}$, then it is called a Hilbert C^* -module over A , or a Hilbert A -module. For example, the C^* -algebra A itself can be recognized to become a Hilbert A -module if the inner product is defined by $\langle a, b \rangle = ab^*$, for all $a, b \in A$.

Frames, orthogonal bases and Riesz bases for Hilbert C^* -modules were defined by Frank and Larson [6].

Let A be a unital C^* -algebra. A sequence $\{x_i\}_{i \in I}$ of elements in a Hilbert A -module H is called a frame for H if there exist two constants $C, D > 0$, such that

$$C \langle x, x \rangle \leq \sum_{i \in I} \langle x, x_i \rangle \langle x_i, x \rangle \leq D \langle x, x \rangle, \quad \forall x \in H,$$

where the sum converges weakly. The constants C and D are called the lower and upper frame bounds, respectively.

The frame $\{x_i\}_{i \in I}$ is called a tight frame if $C = D$, and is said to be a Parseval or normalized tight frame if $C = D = 1$. Likewise, $\{x_i\}_{i \in I}$ is called a Bessel sequence for H with positive bound D if

$$\sum_{i \in I} \langle x, x_i \rangle \langle x_i, x \rangle \leq D \langle x, x \rangle, \quad \forall x \in H.$$

A sequence $\{x_i\}_{i \in I}$ in a Hilbert A -module H is called an orthogonal basis for H if it is a generating set (i.e., the A -linear hull of $\{x_i\}_{i \in I}$ is weak-dense in H) such that

i) $\langle x_i, x_j \rangle = 0$ for each $i \neq j$,

ii) $\|x_i\| = 1$ for each $i \in I$,

iii) the A -linear combinations $\sum_{i \in S} a_i x_i$ with coefficients $\{a_i : i \in S\} \subseteq A$ and $S \subseteq I$ are equal to zero if and only if every summand $a_i x_i$ is equal to zero, $i \in S$.

A sequence $\{x_i\}_{i \in I}$ in a separable Hilbert A -module H is called a Riesz basis for H if it is a frame and a generating set with the additional property that A -linear combinations $\sum_{i \in I} a_i x_i$ with coefficients $\{a_i : i \in S\} \subseteq A$ and $S \subseteq I$ are equal to zero if and only if every summand $a_i x_i$ is equal to zero, $i \in S$.

2.2. Positive operators in $B(H, K)$

In the rest of this paper, let H and K be separable Hilbert spaces and let $B(H, K)$ be the set of all bounded linear operators from H into K . $B(H, K)$ is a Hilbert $B(K)$ -module with a $B(K)$ -valued inner product $\langle S, T \rangle = ST^*$ for all $S, T \in B(H, K)$, and with a linear operation of $B(K)$ on $B(H, K)$ by the composition of operators. On the other hand, $B(H, K)$ is also a Banach space with respect to the operator norm $\|T\|_o = \sup\{\|Tx\| : \|x\| \leq 1, x \in H\}$, for all $T \in B(H, K)$. The norm in $B(H, K)$ considered as a Hilbert $B(K)$ -module is defined by

$$\|T\|_{c^*} = \|\langle T, T \rangle\|_o^{\frac{1}{2}} = \|TT^*\|_o^{\frac{1}{2}} = \|T\|_o.$$

Therefore, the norms in $B(H, K)$ considered as a Hilbert A -module and as a Banach space are the same. However, $B(H, K)$ is not a Hilbert space, and some facts that are true for Hilbert spaces may not hold for $B(H, K)$.

In the study of frame operators on $B(H, K)$, we need to know some facts about operators and, especially, positive operators on $B(H, K)$.

Proposition 2.1 *Let S be an operator on $B(H, K)$, then $\langle SU, U \rangle = 0$ for all $U \in B(H, K)$ if and only if $S = 0$.*

Proof Clearly if $S = 0$, then $\langle SU, U \rangle = 0$ for all $U \in B(H, K)$. On the other hand, we have

$$\langle S(U + V), U + V \rangle = 0, \quad \forall U, V \in B(H, K),$$

so that $\langle SV, U \rangle + \langle SU, V \rangle = 0$. If V changes with iV , we have

$$i \langle SV, U \rangle - i \langle SU, V \rangle = 0,$$

that implies $2i \langle SU, V \rangle = 0$ or $\langle SU, V \rangle = 0$. By setting $V = SU$, we conclude that $SU = 0$ for all $U \in B(H, K)$ and so $S = 0$. □

A map S on $B(H, K)$ is said to be adjointable if there exists a map S^* on $B(H, K)$ such that

$$\langle SU, V \rangle = \langle U, S^*V \rangle, \quad \forall U, V \in B(H, K).$$

Such a map S^* is called the adjoint of S . It follows that S and S^* are bounded linear $B(K)$ -module maps. By $B(B(H, K))$ we denote the set of all adjointable linear $B(K)$ -module maps on $B(H, K)$, and $B_b(B(H, K))$

denotes the set of all bounded linear $B(K)$ -module maps on $B(H, K)$. An adjointable map S on $B(H, K)$ is said to be self-adjoint if $S = S^*$ [16].

Proposition 2.2 *Let S be a adjointable linear $B(K)$ -module map on $B(H, K)$. Then*

i) S is self adjoint if and only if $\langle SU, U \rangle$ is self adjoint for all $U \in B(H, K)$,

ii) S is self adjoint if and only if for all $U \in B(H, K)$, $\langle SU, U \rangle$ is normal and the spectrum of $\langle SU, U \rangle$ is a subset of the real line.

Proof i) If S is self adjoint, then

$$\langle SU, U \rangle = \langle U, SU \rangle = \langle SU, U \rangle^* \quad \forall U \in B(H, K).$$

Conversely, if $\langle SU, U \rangle = \langle SU, U \rangle^*$ for all $U \in B(H, K)$, then

$$\langle SU, U \rangle = \langle U, SU \rangle = \langle S^*U, U \rangle$$

or

$$\langle SU, U \rangle = \langle S^*U, U \rangle \quad \forall U \in B(H, K)$$

and this means $S = S^*$.

ii) If $S = S^*$, then

$$\langle SU, U \rangle = \langle U, S^*U \rangle = \langle U, SU \rangle = \langle SU, U \rangle^*$$

is a self adjoint operator on $B(K)$ for each $U \in B(H, K)$, and by [14] its spectrum is a subset of the real line. Conversely, let the spectrum of $\langle SU, U \rangle$ be a subset of the real line for all $U \in B(H, K)$. For $\alpha \in \mathbb{C}$ and $U, V \in B(H, K)$ we have

$$\langle S(U + \alpha V), U + \alpha V \rangle = \langle SU, U \rangle + \bar{\alpha} \langle SV, U \rangle + \alpha \langle SU, V \rangle + |\alpha|^2 \langle SV, V \rangle.$$

Since the spectrum of $\langle S(U + \alpha V), U + \alpha V \rangle$ is a subset of the real line,

$$\langle S(U + \alpha V), U + \alpha V \rangle = \langle S(U + \alpha V), U + \alpha V \rangle^*,$$

and

$$\begin{aligned} \alpha \langle SV, U \rangle + \bar{\alpha} \langle SU, V \rangle &= \bar{\alpha} \langle SV, U \rangle^* + \alpha \langle SU, V \rangle^* \\ &= \bar{\alpha} \langle U, SV \rangle + \alpha \langle V, SU \rangle \\ &= \bar{\alpha} \langle S^*U, V \rangle + \alpha \langle S^*V, U \rangle. \end{aligned}$$

By setting $\alpha = 1$ and $\alpha = i$,

$$\begin{aligned} \langle SV, U \rangle + \langle SU, V \rangle &= \langle S^*U, V \rangle + \langle S^*V, U \rangle, \\ i \langle SV, U \rangle - i \langle SU, V \rangle &= -i \langle S^*U, V \rangle + i \langle S^*V, U \rangle. \end{aligned}$$

Now by product i in the second equality, we obtain $\langle SU, V \rangle = \langle S^*U, V \rangle$. Therefore, $S = S^*$. □

Remark 2.3 An element $S \in B(B(H, K))$ is said to be positive if $S = S^*$ and the spectrum of S is contained in the positive real line [11]. Wegge-Olsen [16] has shown that $S \geq 0$ if and only if the spectrum of $\langle ST, T \rangle$ is a subset of $[0, \infty)$ for all $T \in B(H, K)$.

Proposition 2.4 Let S be a positive operator in $B(B(H, K))$. Then

$$\|S\| = \sup_{\|T\| \leq 1} \|\langle ST, T \rangle\|.$$

Proof Since $B(B(H, K))$ is a C^* -algebra [16] and S is positive, $S^{\frac{1}{2}}$ exists. Then, we have

$$\begin{aligned} \sup_{\|T\| \leq 1} \|\langle ST, T \rangle\| &= \sup_{\|T\| \leq 1} \left\| \left\langle S^{\frac{1}{2}} S^{\frac{1}{2}} T, T \right\rangle \right\| \\ &= \sup_{\|T\| \leq 1} \left\| \left\langle S^{\frac{1}{2}} T, S^{\frac{1}{2}} T \right\rangle \right\| \\ &= \sup_{\|T\| \leq 1} \left\| S^{\frac{1}{2}} T \right\|^2 \\ &= \left\| S^{\frac{1}{2}} \right\|^2 = \left\| S^{\frac{1}{2}} (S^{\frac{1}{2}})^* \right\| \\ &= \left\| S^{\frac{1}{2}} S^{\frac{1}{2}} \right\| = \|S\|. \end{aligned}$$

□

Lemma 2.5 Let $\Lambda \in B(H)$. Then Λ is positive if and only if $T\Lambda T^* \in B(K)$ is positive for all $T \in B(H, K)$.

Proof Let Λ be positive. Since $B(H)$ is a C^* - algebra, there is $\Gamma \in B(H)$ such that $\Lambda = \Gamma\Gamma^*$, and so

$$T\Lambda T^* = T\Gamma\Gamma^* T^* = T\Gamma(T\Gamma)^*.$$

On the other hand, for all $f \in H$, we have

$$\langle T\Gamma(T\Gamma)^* f, f \rangle = \langle (T\Gamma)^* f, (T\Gamma)^* f \rangle = \|(T\Gamma)^* f\|^2 \geq 0.$$

Hence $T\Lambda T^*$ is positive.

Conversely, let $f \in H$ be arbitrary. We can find $g \in K$ and $T \in B(H, K)$ such that $T^*g = f$. Then by the positivity of $T\Lambda T^*$,

$$\langle \Lambda f, f \rangle = \langle T\Lambda T^* g, g \rangle \geq 0.$$

Therefore, Λ is positive. □

3. Operator sequences, g-sequences and their relations

3.1. Frames

A sequence $\{T_i \in B(H, K) : i \in I\}$ is said to be a frame for $B(H, K)$ if there exist $0 < A, B < \infty$ such that

$$A \langle T, T \rangle \leq \sum_{i \in I} \langle T, T_i \rangle \langle T_i, T \rangle \leq B \langle T, T \rangle, \quad \forall T \in B(H, K), \quad (3.1)$$

where the series converges in the strong operator topology. The frame operator on $B(H,K)$ is defined by

$$S: B(H, K) \longrightarrow B(H, K),$$

$$ST = \sum_{i \in I} \langle T, T_i \rangle T_i = \sum_{i \in I} TT_i^*T_i.$$

Proposition 2.2, Remark 2.3 and (3.1) assert that S is a positive, self adjoint and invertible operator, and

$$\langle ST, T \rangle = \sum_{i \in I} TT_i^*T_iT^* = \sum_{i \in I} \langle T, T_i \rangle \langle T_i, T \rangle.$$

Therefore, we have

$$A \langle T, T \rangle \leq \langle ST, T \rangle \leq B \langle T, T \rangle.$$

Convergence in the definition of frames, Bessel sequences, orthogonal and Riesz bases in $B(H,K)$ as a Hilbert $B(K)$ -module is in the strong operator topology.

Various generalizations of frames have been studied by many authors. Sun [15] introduced a type of frames called g-frames, and showed that most generalizations of frames can be regarded as special cases of g-frames. Here we point out that g-frames can be regarded as frames in $B(H, K)$ with the same bounds.

A sequence $\{\Lambda_i \in B(H, K_i) : i \in I\}$ is called a generalized frame, or simply a g-frame for H with respect to a sequence of Hilbert spaces $\{K_i\}_{i \in I}$ if there exist two positive constants A and B such that

$$B\|f\|^2 \leq \sum_{i \in I} \|\Lambda_i f\|^2 \leq A\|f\|^2, \quad \forall f \in H.$$

A and B are called the lower and upper g-frame bounds, respectively. A g-frame is called tight if $A = B$ and Parseval g-frame if $A = 1$. In simple terms, $\{\Lambda_i\}_{i \in I}$ is called a g-frame for H whenever the space sequence $\{K_i : i \in I\}$ is clear, and also a g-frame for H with respect to K whenever $K_i = K$ for each $i \in I$. A sequence $\Lambda_i \in B(H, K_i) : i \in I$ is called a g-Bessel sequence with bound B if we have only an upper bound in the definition of g-frames. The space $(\sum_{i \in I} \oplus K_i)_{l_2}$ is defined by

$$\left(\sum_{i \in I} \oplus K_i \right)_{l_2} = \left\{ \{f_i\}_{i \in I} : f_i \in K_i, \quad i \in I \quad \text{and} \quad \sum_{i \in I} \|f_i\|^2 \leq \infty \right\}$$

and has the inner product,

$$\langle \{f_i\}, \{g_i\} \rangle = \sum_{i \in I} \langle f_i, g_i \rangle.$$

It is clear that $(\sum_{i \in I} \oplus K_i)_{l_2}$ is a Hilbert space and contains K_i as a subspace, $i \in I$.

Remark 3.1 Let $\{\Lambda_i\}_{i \in I}$ be a g-frame for H with respect to $\{K_i\}_{i \in I}$ and let $K = (\sum_{i \in I} \oplus K_i)_{l_2}$. For $i \in I$, define $\Lambda'_i : H \mapsto K$ by

$$\Lambda'_i f = (\dots, 0, 0, 0, \Lambda_i f, 0, 0, 0, \dots), \quad \forall f \in H.$$

Then

$$\|\Lambda_i' f\| = \|\Lambda_i f\|, \quad \forall i \in I, \forall f \in H.$$

Hence $\{\Lambda_i\}_{i \in I}$ is a g -frame for H with respect to $\{K_i\}_{i \in I}$ if and only if $\{\Lambda_i'\}_{i \in I}$ is a g -frame for H with respect to K . Therefore, without loss of generality, we may deal with g -frames for H with respect to K .

Now we shall show that a g -frame for H with respect to K is a frame for $B(H, K)$, and vice versa.

Theorem 3.2 Let $\{\Lambda_i \in I\}_{i \in I}$ be a sequence in $B(H, K)$. Then it is a frame for $B(H, K)$ considered as a Hilbert C^* -module if and only if it is a g -frame for H with respect to K .

Proof Let $\{\Lambda_i \in B(H, K) : i \in I\}$ be a g -frame for H with respect to K . Then there are positive constants A and B , such that

$$B \langle f, f \rangle \leq \sum_{i \in I} \langle \Lambda_i \Lambda_i^* f, f \rangle \leq A \langle f, f \rangle, \quad \forall f \in H.$$

Hence

$$BI_H \leq \sum_{i \in I} \Lambda_i^* \Lambda_i \leq AI_H.$$

Lemma 2.5 asserts that the inequality

$$BTT^* \leq \sum_{i \in I} T \Lambda_i^* \Lambda_i T^* \leq ATT^*,$$

satisfies for all $T \in B(H, K)$. Thus,

$$B \langle T, T \rangle \leq \sum_{i \in I} \langle T, \Lambda_i \rangle \langle \Lambda_i, T \rangle \leq A \langle T, T \rangle, \quad \forall T \in B(H, K),$$

and $\{\Lambda_i\}_{i \in I}$ is a frame for $B(H, K)$. Conversely, let $\{\Lambda_i \in B(H, K) : i \in I\}$ be a frame for $B(H, K)$ and $f \in H$. We can choose T in $B(H, K)$ and g in K such that $T^*g = f$. Therefore,

$$\begin{aligned} \left\langle \sum_{i \in I} \langle T, \Lambda_i \rangle \langle \Lambda_i, T \rangle g, g \right\rangle &= \left\langle \sum_{i \in I} T \Lambda_i^* \Lambda_i T^* g, g \right\rangle \\ &= \sum_{i \in I} \langle T \Lambda_i^* \Lambda_i T^* g, g \rangle \\ &= \sum_{i \in I} \langle \Lambda_i T^* g, \Lambda_i T^* g \rangle \\ &= \sum_{i \in I} \langle \Lambda_i f, \Lambda_i f \rangle \\ &= \sum_{i \in I} \|\Lambda_i f\|^2. \end{aligned}$$

Also we have

$$\langle B \langle T, T \rangle g, g \rangle = \langle BTT^* g, g \rangle = B \langle T^* g, T^* g \rangle = B \langle f, f \rangle = B \|f\|^2.$$

Thus

$$B \langle T, T \rangle \leq \sum_{i \in I} \langle T, \Lambda_i \rangle \langle \Lambda_i, T \rangle \leq A \langle T, T \rangle,$$

implies that

$$B \|f\|^2 \leq \sum_{i \in I} \|\Lambda_i f\|^2 \leq A \|f\|^2, \quad \forall f \in H,$$

as desired. □

The following are immediate consequences.

Corollary 3.3 *The sequence $\{\Lambda_i \in B(H, K) : i \in I\}$ is a tight frame for $B(H, K)$ if and only if it is a g -tight frame for H with respect to K .*

Corollary 3.4 *The sequence $\{\Lambda_i \in B(H, K) : i \in I\}$ is a Bessel sequence for $B(H, K)$ if and only if it is a g -Bessel sequence for H with respect to K .*

Remark 3.5 *Let $\{\Lambda_i \in B(H, K) : i \in I\}$ be a g -frame for H . The g -frame operator of $\{\Lambda_i\}_{i \in I}$ is defined by*

$$S_g : H \longrightarrow H, \quad f \longmapsto \sum_{i \in I} \Lambda_i^* \Lambda_i f.$$

Also, the frame operator of the frame $\{\Lambda_i\}_{i \in I}$ is defined by $ST = \sum_{i \in I} T \Lambda_i^ \Lambda_i$. Therefore, $ST = TS_g$, and from this equation, for any $T \in B(H, K)$ a reconstruction formula is derived by $T = S^{-1}TS_g$.*

3.2. Orthonormal bases and Riesz bases

Now, we study the relations between g -orthonormal bases and g -Riesz bases for H with respect to K with orthogonal bases and Riesz bases for $B(H, K)$ considered as a Hilbert C^* -module.

A sequence $\{\Lambda_i : i \in I\}$ is called a g -orthonormal basis for H with respect to K if it satisfies the following:

- 1) $\langle \Lambda_i^* f, \Lambda_j^* g \rangle = \delta_{i,j} \langle f, g \rangle, \quad \forall i, j \in I$ and $f, g \in H$,
- 2) $\sum_{i \in I} \|\Lambda_i f\|^2 = \|f\|^2, \quad \forall f \in H$.

If $\{\Lambda_i : i \in I\}$ is a sequence in $B(H, K)$ and $\{f : \Lambda_i f = 0, i \in I\} = \{0\}$, then $\{\Lambda_i : i \in I\}$ is called g -complete.

A sequence $\{\Lambda_i \in B(H, K) : i \in I\}$ is called a g -Riesz basis for H with respect to K if it is g -complete and there are positive constants A and B such that

$$A \sum_{i \in I_1} \|g_i\|^2 \leq \left\| \sum_{i \in I_1} \Lambda_i^* g_i \right\|^2 \leq B \sum_{i \in I_1} \|g_i\|^2,$$

for any finite subset I_1 of I and $\{g_i\}_{i \in I_1} \subseteq K$ [15].

The following theorem provide that every g -orthonormal basis for H is an orthogonal basis for $B(H, K)$. We will give an example to show that the converse of the theorem is not correct.

Theorem 3.6 *If $\{\Lambda_i \in B(H, K) : i \in I\}$ is a g-orthonormal basis for H with respect to K , then it is an orthogonal basis for $B(H, K)$ considered as a Hilbert C^* -module.*

Proof Since $\{\Lambda_i\}_{i \in I}$ is a g-orthonormal basis for H , for $i \neq j$, we have $\langle \Lambda_i, \Lambda_j \rangle = \Lambda_i \Lambda_j^* = 0$ and $\|\Lambda_i\|^2 = \|\langle \Lambda_i, \Lambda_i \rangle\| = \|\Lambda_i \Lambda_i^*\| = \|I_K\| = 1$, where I_K is the identity operator on K . Now suppose that $\sum_{i \in I} T_i \Lambda_i = 0$ where $T_i \in B(K), i \in I$. We have

$$\begin{aligned} 0 &= \left\langle \sum_{i \in I} T_i \Lambda_i, \Lambda_j \right\rangle = \sum_{i \in I} \langle T_i \Lambda_i, \Lambda_j \rangle \\ &= \sum_{i \in I} T_i \langle \Lambda_i, \Lambda_j \rangle \\ &= T_j \langle \Lambda_j, \Lambda_j \rangle \\ &= T_j I_H = T_j. \end{aligned}$$

Therefore, $T_j = 0$ and $T_j \Lambda_j = 0$, for each $j \in I$. It remains to show that every $T \in B(H, K)$ can be generated by $\{\Lambda_i\}_{i \in I}$. The second condition of g-orthonormal basis, $\sum_{i \in I} \|\Lambda_i f\|^2 = \|f\|^2$ for all $f \in H$, implies that $\sum_{i \in I} \Lambda_i^* \Lambda_i = I_H$. Then, for every $T \in B(H, K)$ we have

$$\begin{aligned} T &= T I_H = T \sum_{i \in I} \Lambda_i^* \Lambda_i \\ &= \sum_{i \in I} T \Lambda_i^* \Lambda_i \\ &= \sum_{i \in I} \langle T, \Lambda_i \rangle \Lambda_i \\ &= \sum_{i \in I} U_i \Lambda_i \end{aligned}$$

where $U_i = \langle T, \Lambda_i \rangle$ belongs to $B(K)$, for each $i \in I$. This completes the proof of the theorem. □

The relation between Riesz bases for $B(H, K)$ and g-Riesz bases for H with respect to K is similar to the above theorem.

Theorem 3.7 *If $\{\Lambda_i \in B(H, K) : i \in I\}$ is a g-Riesz basis for H with respect to K , then it is a Riesz basis for $B(H, K)$ considered as a Hilbert C^* -module.*

Proof Let $\{\Lambda_i \in B(H, K) : i \in I\}$ be a g-Riesz basis for H with respect to K . By ([15], Corollary 3.3) $\{\Lambda_i\}_{i \in I}$ is a g-frame and by Theorem 3.2 it is a frame for $B(H, K)$. It is clear that $\Lambda_i \neq 0$ for each $i \in I$. Now let $\sum_{i \in I} T_i \Lambda_i = 0$, where $T_i \in B(K)$. We have $\sum_{i \in I} \Lambda_i^* T_i^* = 0$, therefore, $\sum_{i \in I} \Lambda_i^* T_i^* g = 0$, for each $g \in K$. By the definition of g-Riesz basis, $\sum_{i \in I} \|T_i^* g\|^2 = 0$, then $\|T_i^* g\|^2 = 0$, for each $i \in I$ and $g \in K$. Therefore, $T_i = 0$ and hence $T_i \Lambda_i = 0$ for each $i \in I$. The invertibility of the frame operator S implies that

$$T = \sum_{i \in I} \langle S^{-1} T, \Lambda_i \rangle \Lambda_i, \quad \forall T \in B(H, K).$$

Thus, $\{\Lambda_i\}_{i \in I}$ is a generating set for $B(H, K)$ and the proof is complete. □

By an example we show that the converse of Theorem 3.5 and Theorem 3.6 is not true.

Example 3.8 Let H be a Hilbert space and $\{\varphi_i\}_{i \in I}$ be an orthonormal basis for H . For $i \in I$, define Λ_i and Λ_i^* by

$$\begin{aligned} \Lambda_i : H &\mapsto \mathbb{C}^2, & f &\mapsto (\langle f, \varphi_i \rangle, 0), \\ \Lambda_i^* : \mathbb{C}^2 &\mapsto H, & (c_1, c_2) &\mapsto c_1 \varphi_i. \end{aligned}$$

Since

$$\sum_{i \in I} T \Lambda_i^* \Lambda_i T^* = TT^*, \quad \text{for all } T \in B(H, \mathbb{C}^2),$$

the sequence $\{\Lambda_i\}_{i \in I}$ is a Parseval frame for $B(H, \mathbb{C}^2)$. Now let $\{T_i\}_{i \in I}$ be a sequence in $B(\mathbb{C}^2)$ and $\sum_{i \in I} T_i \Lambda_i = 0$. Then, for each $f \in H$ we have

$$0 = \sum_{i \in I} T_i \Lambda_i f = \sum_{i \in I} T_i (\langle \varphi_i, f \rangle, 0) = \sum_{i \in I} \langle \varphi_i, f \rangle T_i (1, 0).$$

By the orthonormality of $\{\varphi_i\}_{i \in I}$, $T_i (1, 0) = 0$, hence, $T_i \Lambda_i = 0$ for all $i \in I$. Also, $\sum_{i \in I} \Lambda_i^* \Lambda_i = I_H$ implies that

$$T = \sum_{i \in I} T \Lambda_i^* \Lambda_i = \sum_{i \in I} \langle T, \Lambda_i \rangle \Lambda_i.$$

This shows that $\{\Lambda_i\}_{i \in I}$ generates $B(H, \mathbb{C}^2)$ as $B(\mathbb{C}^2)$ -module. Therefore, all conditions of a Riesz basis are satisfied and $\{\Lambda_i\}_{i \in I}$ is a Riesz basis for $B(H, \mathbb{C}^2)$. But $\{\Lambda_i\}_{i \in I}$ is not a g -Riesz basis since $\Lambda_i^*(0, 1) = 0$, which implies that $A = 0$ in the definition of a g -Riesz basis.

However, $\langle \Lambda_i, \Lambda_i \rangle (0, 1) = \Lambda_i \Lambda_i^* = (0, 0)$, $\Lambda_i \Lambda_i^* \neq I_{\mathbb{C}^2}$, $\|\Lambda_i\| = 1$ and $\langle \Lambda_i, \Lambda_j \rangle = 0$ for $i \neq j$. Therefore, $\{\Lambda_i\}_{i \in I}$ is an orthogonal basis for $B(H, \mathbb{C}^2)$. On the other hand, $\langle \Lambda_i^*(0, 1), \Lambda_i^*(0, 1) \rangle = 0$ and $\delta_{ii} \langle (0, 1), (0, 1) \rangle = 1$, imply that $\{\Lambda_i\}_{i \in I}$ is not a g -orthonormal basis for H with respect to K .

References

- [1] Bodmann, B.G., Kribs, D.W., Paulsen, V.I.: Decoherence-insensitive quantum communication by optimal C^* -encoding. IEEE Trans. Inform. Theory 53, 4738-4749 (2007).
- [2] Christensen, O.: An Introduction to Frames and Riesz Bases. Boston-Basel-Berlin. Birkhäuser 2002.
- [3] D'Attellis, C.E., Fernfedez-Berdaguer, E.M.: Wavelet Theory and Harmonic Analysis in Applied Sciences. Boston - Basel - Berlin. Birkhäuser 1997.
- [4] Daubechies, I., Grossmann, A., Meyer, Y.: Painless nonorthogonal expansions. J. Math. Phys. 27, 1271-1283 (1986).
- [5] Duffin, R., Schaeffer, A.: A class of nonharmonic Fourier series. Trans. Amer. Math. Soc. 72, 341-366 (1952).
- [6] Frank, M., Larson, D.: Frames in Hilbert C^* -modules and C^* -algebra. J. Operator Theory 48, 273-314 (2002).
- [7] Gröchenig, K.: Foundations of Time-Frequency Analysis. Appl. Numer. Harmon. Anal. Boston. MA, Birkhäuser 2001.
- [8] Hemmat, A. A., Gabardo, J. P.: Properties of oblique dual frames in shift-invariant systems. J. Math. Anal. Appl. 356, 346-354 (2009).
- [9] Kaplansky, I.: Algebra of type I. Ann. Math. 56, 460-472 (1952).
- [10] Lance, E.: Hilbert C^* -modules - a Toolkit for Operator Algebraists. London Mathematical Society Lecture Note Series v. 210. Cambridge, England. Cambridge University Press 1995.

- [11] Murphy, G. J.: C^* -algebra and Operator Theory. London. Academic Press 1990.
- [12] Paschke, W.: Inner product modules over B^* -algebra. Trans. Amer. Math. Soc. 182, 443-468 (1973).
- [13] Rieffel, M.: Morita equivalence for C^* -algebra. J. Pure Applied Algebra 5, 51-96 (1974).
- [14] Rudin, W.: Functional Analysis. New York. McGraw-Hill Book Company 1973.
- [15] Sun, W.: G-frames and g-Riesz bases. J. Math. Anal. 322, 437-452 (2006).
- [16] Wegga-Olsen, N.: K-theory and C^* -algebra - a Friendly Approach. Oxford, England. Oxford University Press 1993.
- [17] Young, R.: An Introduction to Nonharmonic Fourier Series. New York. Academic Press 1980.