

## Almost analytic forms with respect to a quadratic endomorphism and their cohomology

Mircea CRASMAREANU, Cristian IDA\*

Faculty of Mathematics Department of Mathematics and Informatics University,  
“Al. I. Cuza” University Transilvania of Braşov, Iaşi Braşov Romania, Romania

Received: 09.04.2014

Accepted/Published Online: 14.01.2015

Printed: 29.05.2015

**Abstract:** The goal of this paper is to consider the notion of almost analytic form in a unifying setting for both almost complex and almost paracomplex geometries. We use a global formalism, which yields, in addition to generalizations of the main results of the previously known almost complex case, a relationship with the Frölicher–Nijenhuis theory. A cohomology of almost analytic forms is also introduced and studied as well as deformations of almost analytic forms with pairs of almost analytic functions.

**Key words:** Quadratic endomorphism, almost  $F$ -analytic form,  $F$ -symmetric form, almost (para)complex structure, cohomology

### 1. Introduction

The notion of almost analytic form was introduced a long time ago in the almost complex geometry and hence it was treated in local coordinates, especially by Japanese geometers [15, 16, 17, 18]. A global approach appeared in [14], unfortunately only in Romanian. Some of these global techniques were used in [9] and [13]; for example, in the former paper a differential is introduced in the algebra of pairs of almost analytic forms and a corresponding Poincaré type lemma is proved.

The present work aims to consider almost analytic forms in a unifying setting, which adds the almost paracomplex geometry. This type of even dimensional geometry is now in the mainstream of research as the surveys [1] and [4] and their several citations prove. In this way, we reveal the common parts of these geometries with respect to differential forms and present the techniques of [14] to a larger audience. An important feature of the global approach is that it yields a relationship with the Frölicher–Nijenhuis theory, widely used now for several important topics. Namely, we prove that for an almost  $F$ -analytic form its closeness with respect to the Frölicher–Nijenhuis derivative  $d_F$  is characterized by the usual (i.e. exterior derivative) closeness.

The content of the paper is as follows. In the first subsection of Section 2 we consider only 1-forms in order to offer a detailed picture of the techniques used herein. In the next subsection we consider the general case of  $r$ -forms with  $r$  less than or equal to  $n =$  half of the dimension of the underlying manifold. A  $d_F$ -cohomology of almost analytic forms is introduced and studied and also some deformations of almost analytic forms with pairs of almost analytic functions are considered. In Section 3 we restrict ourselves to the Hermitian and para-Norden framework and reobtain the characterization of almost analyticity for  $n$ -forms in terms of

\*Correspondence: cristian.ida@unitbv.ro

2010 AMS Mathematics Subject Classification: 53C15, 58A10, 58A12.

harmonicity. Considering again the case of 1-forms, a local computation in the case of integrability of given endomorphism  $F$  gives an usual characterization of coefficients in terms of (para)Cauchy–Riemann equations.

## 2. Almost analytic forms with respect to a quadratic endomorphism

### 2.1. Almost analytic 1-forms

Fix a triple  $(M, F, \omega)$  with  $M$  a smooth  $m$ -dimensional manifold,  $F$  a tensor field of  $(1, 1)$ -type on  $M$ , and  $\omega$  a differentiable 1-form, i.e.  $\omega \in \Omega^1(M)$ .

**Definition 2.1** *i)  $F$  is a quadratic endomorphism if there exists  $\varepsilon \in \mathbb{R}^*$  such that:*

$$F^2 = \varepsilon I. \tag{2.1}$$

*ii) The  $F$ -conjugate of  $\omega$  is the 1-form:*

$$\bar{\omega} = \omega_F := \omega \circ F^{-1} = \frac{1}{\varepsilon} \omega \circ F. \tag{2.2}$$

It follows that:

$$\bar{\bar{\omega}} = \frac{1}{\varepsilon} \bar{\omega} \circ F = \frac{1}{\varepsilon} \omega. \tag{2.3}$$

To the pair  $(F, \omega)$  we associate a 2-form defined by:

$$\Omega_{F,\omega}(X, Y) := d\omega(FX, Y) - \varepsilon d\bar{\omega}(X, Y), \tag{2.4}$$

which yields the main notion of this subsection:

**Definition 2.2** *The 1-form  $\omega$  is called almost  $F$ -analytic if  $\Omega_{F,\omega} = 0$ . Let  $\Omega^1(M, F)$  be the set of almost  $F$ -analytic 1-forms.*

In the following we use the identity:

$$2d\omega(X, Y) = X(\omega(Y)) - Y(\omega(X)) - \omega([X, Y]). \tag{2.5}$$

A first result shows that this property is invariant under  $F$ -conjugation:

**Proposition 2.1** *The 1-form  $\omega$  is almost  $F$ -analytic if and only if its  $F$ -conjugate  $\bar{\omega}$  is almost  $F$ -analytic. If  $\omega$  is almost  $F$ -analytic then  $\omega$  is closed if and only if  $\bar{\omega}$  is closed.*

**Proof** Using (2.1), (2.3), and (2.4) we get:

$$\Omega_{F,\bar{\omega}}(X, Y) = -\frac{1}{\varepsilon} \Omega_{F,\omega}(FX, Y), \quad \Omega_{F,\omega}(X, Y) = -\Omega_{F,\bar{\omega}}(FX, Y) \tag{2.6}$$

and the conclusion follows directly from (2.6). □

Recall now the Nijenhuis tensor field of  $F$ :

$$N_F(X, Y) := [FX, FY] - F[FX, Y] - F[X, FY] + F^2[X, Y], \tag{2.7}$$

which for our case (2.1) becomes  $N_F(X, Y) = [FX, FY] - F[FX, Y] - F[X, FY] + \varepsilon[X, Y]$ . We have the following skew-symmetries:

$$N_F(FX, Y) = -FN_F(X, Y) = N_F(X, FY), \quad N_F(FX, FY) = \varepsilon N_F(X, Y) \tag{2.8}$$

which yields a second property of almost  $F$ -analytic forms:

**Proposition 2.2** *If  $\omega$  is almost  $F$ -analytic then:*

$$\omega \circ N_F = \bar{\omega} \circ N_F = 0. \tag{2.9}$$

**Proof** Let  $\omega$  be almost  $F$ -analytic. Using (2.5),  $\omega \circ F = \varepsilon\bar{\omega}$ , and  $\bar{\omega} \circ F = \omega$ , from  $\Omega_{F,\omega}(X, Y) = d\omega(FX, Y) - \varepsilon d\bar{\omega}(X, Y) = 0$  we easily obtain:

$$\omega(F[X, Y]) = \varepsilon X\bar{\omega}(Y) - FX\omega(Y) + \omega([FX, Y]). \tag{2.10}$$

Putting  $X \mapsto FX$  and  $Y \mapsto FY$  in (2.10), by direct calculus we obtain:

$$\begin{aligned} (\omega \circ N_F)(X, Y) &= \omega([FX, FY]) - \omega(F[FX, Y]) - \omega(F[X, FY]) + \varepsilon\omega([X, Y]) \\ &= \omega([FX, FY]) - \varepsilon FX(\bar{\omega}(Y)) + \varepsilon X\omega(Y) - \varepsilon\omega([X, Y]) \\ &\quad - \varepsilon X\omega(Y) + \varepsilon FX\bar{\omega}(Y) - \omega([FX, FY]) + \varepsilon\omega([X, Y]) = 0. \end{aligned}$$

By Proposition 2.1  $\bar{\omega}$  is also almost  $F$ -analytic and the relation  $(\bar{\omega} \circ N_F)(X, Y) = 0$  follows in a similar manner starting from  $\Omega_{F,\bar{\omega}}(X, Y) = d\bar{\omega}(FX, Y) - d\omega(X, Y) = 0$ . □

Another tool in our study is provided by the Obata operators associated to  $F$ , namely the maps  $O_F, O_F^* : \Omega^2(M) \rightarrow \Omega^2(M)$ :

$$\begin{cases} O_F(\rho)(X, Y) := \frac{1}{2}[\rho(X, Y) - \rho(FX, FY)] \\ O_F^*(\rho)(X, Y) := \frac{1}{2}[\rho(X, Y) + \rho(FX, FY)], \end{cases} \tag{2.11}$$

which give a classification of 2-forms with respect to  $F$ :

**Definition 2.3** *The 2-form  $\rho$  is called  $F$ -pure if  $O_F^*(\rho) = 0$  and respectively  $F$ -hybrid if  $O_F(\rho) = 0$ .*

**Proposition 2.3** *i) If  $F$  is an almost complex structure ( $\varepsilon = -1$ ) and  $\omega$  is almost  $F$ -analytic form then the 2-forms  $d\omega, d\bar{\omega}$  are  $F$ -pure.*

*ii) If  $F$  is an almost product structure ( $\varepsilon = 1$ ) and  $\omega$  is almost  $F$ -analytic form then the 2-forms  $d\omega, d\bar{\omega}$  are  $F$ -hybrid.*

**Proof** i) Let  $\varepsilon = -1$ . From the characterization of almost  $F$ -analyticity, setting  $X \mapsto FX$  in (2.10) we have:

$$X(\omega(Y)) + FX(\omega(FY)) = \omega([X, Y]) + \omega(F[FX, Y]), \tag{2.12}$$

and now  $X \rightarrow Y$  in (2.12):

$$Y(\omega(X)) + FY(\omega(FX)) = -\omega([X, Y]) - \omega(F[X, FY]). \tag{2.13}$$

From (2.13) minus (2.12) we get:

$$2d\omega + \omega([X, Y]) + 2d\bar{\omega}(FX, FY) + \omega([FX, FY]) = 2\omega([X, Y]) + \omega \circ F([FX, Y] + [X, FY]),$$

which means:

$$4O_F^*(d\omega) = -\omega \circ N_F = 0.$$

By analogy:

$$4O_F^*(d\bar{\omega}) = -\bar{\omega} \circ N_F = 0.$$

ii) Let  $\varepsilon = 1$ . Again, with  $X \rightarrow FX$  in relation (2.10) we have:

$$X(\omega(Y)) - FX(\omega(FY)) = \omega([X, Y]) - \omega(F[FX, Y]) \tag{2.14}$$

and  $X \leftrightarrow Y$  in this equality gives:

$$Y(\omega(X)) - FY(\omega(FX)) = -\omega([X, Y]) + \omega(F[X, FY]). \tag{2.15}$$

With (2.14) minus (2.15) we obtain:

$$2d\omega(X, Y) + \omega([X, Y]) - 2d\bar{\omega}([FX, FY]) - \omega([FX, FY]) = 2\omega([X, Y]) - \omega \circ F([FX, Y] + [X, FY]),$$

which means:  $4O_F(d\omega) = \omega \circ N_F = 0$ . Also:  $4O_F(d\bar{\omega}) = \bar{\omega} \circ N_F = 0$  and the assertion is proved. □

An important consequence of this result is the following:

**Corollary 2.1** *If  $\varepsilon \in \{-1, +1\}$  then definition (2.4) and hence the definition of almost  $F$ -analyticity do not depend on the place of  $F$ .*

**Proof** From Proposition 2.3 we have that the almost  $F$ -analyticity implies:

$$d\omega(X, Y) = \varepsilon d\omega(FX, FY), \tag{2.16}$$

and then the right-hand side of (2.4) is:

$$d\omega(FX, Y) - \varepsilon d\bar{\omega}(X, Y) = \varepsilon d\omega(F^2X, FY) - \varepsilon d\bar{\omega}(X, Y) = \varepsilon^2 d\omega(X, FY) - \varepsilon d\bar{\omega}(X, Y)$$

and since  $\varepsilon^2 = 1$  we get the conclusion. □

We finish this subsection with a relationship of this formalism with the Frölicher–Nijenhuis theory. Recall that given a tensor field  $F$  of  $(1, 1)$ -type it defines the following:

i) an interior product  $i_F$ ; for an  $r$ -form  $\omega$  we have that  $i_F\omega$  is again an  $r$ -form given by:

$$i_F\omega(X_1, \dots, X_r) := \sum_{i=1}^r \omega(X_1, \dots, FX_i, \dots, X_r), \quad r \geq 1 \text{ and } i_F f = 0, \quad \forall f \in C^\infty(M); \tag{2.17}$$

ii) an exterior  $F$ -derivative  $d_F$  with:

$$d_F := i_F \circ d - d \circ i_F. \tag{2.18}$$

**Proposition 2.4** *If  $\varepsilon = \pm 1$  and  $\omega$  is almost  $F$ -analytic then the exterior  $F$ -derivatives of  $\omega$  and  $\bar{\omega}$  are:*

$$d_F\omega = \frac{1}{2}i_F \circ d\omega = \varepsilon d\bar{\omega}, \quad d_F\bar{\omega} = d\omega. \tag{2.19}$$

**Proof** For  $r = 1$  we have:

$$i_F\omega = \varepsilon\bar{\omega} \tag{2.20}$$

and then:

$$\begin{aligned} (d_F\omega)(X, Y) &= i_F(d\omega)(X, Y) - d(\varepsilon\bar{\omega})(X, Y) \\ &= d\omega(FX, Y) + d\omega(X, FY) - \varepsilon d\bar{\omega}(X, Y) = \Omega_{F,\omega}(X, Y) + d\omega(X, FY), \end{aligned}$$

which means that  $d_F\omega(\cdot, \cdot) = d\omega(\cdot, F\cdot)$ . We apply the previous Corollary 2.1 to get the first part of (2.19). The second part of the required formula follows by duality.  $\square$

Similarly to [6, 10, 16], a smooth function  $f$  on  $M$  is called *almost  $F$ -analytic* if there exists a smooth function  $g$  on  $M$  such that:

$$df \circ F = dg, \tag{2.21}$$

and in this case  $g$  is called *the corresponding function* of  $f$ . In this case  $g$  is also almost  $F$ -analytic with corresponding function  $\varepsilon f$ . Let us denote by  $C^\infty(M, F)$  the set of all almost  $F$ -analytic functions on  $M$ . If  $f \in C^\infty(M, F)$ , then by (2.20) we have:

$$d_Ff = i_F \circ df = \varepsilon d\bar{f} = df \circ F = dg. \tag{2.22}$$

**Proposition 2.5** *If  $f \in C^\infty(M, F)$  then  $df$  and  $d_Ff$  are both almost  $F$ -analytic.*

**Proof** Let  $f \in C^\infty(M, F)$ . Then:

$$\Omega_{F,df}(X, Y) = (d(df))(FX, Y) - \varepsilon(d(\bar{df}))(X, Y) = -(d(dg))(X, Y) = 0,$$

which says that  $df$  is almost  $F$ -analytic. The second assertion follows by setting  $X \mapsto FX$  in the above relation.  $\square$

## 2.2. Almost $F$ -analytic $r$ -forms and $d_F$ -cohomology

In this subsection we give a generalization of previous results to  $r$ -forms for  $r \geq 2$  with  $\varepsilon$  restricted to  $\{-1, +1\}$  and we study the  $d_F$ -cohomology of almost analytic  $r$ -forms.

Firstly, inspired by Proposition 2.3, we introduce a class of  $r$ -forms adapted to  $F$ :

**Definition 2.4** *The  $r$ -form  $\omega$  is called  $F$ -symmetric if for all vector fields  $X_1, \dots, X_r$ :*

$$\omega(FX_1, \dots, X_r) = \omega(X_1, \dots, FX_i, \dots, X_r), \quad 2 \leq i \leq r. \tag{2.23}$$

**Example 2.1** *i) If  $\theta \in \Omega^1(M, F)$  then the 2-forms  $\omega = d\theta$  and  $\bar{\omega} = d\bar{\theta}$  are  $F$ -symmetric. Indeed, equation (2.16) means:*

$$d\theta(X, Y) = \varepsilon d\theta(FX, FY), \quad d\bar{\theta}(X, Y) = \varepsilon d\bar{\theta}(FX, FY)$$

*and with  $X \rightarrow FX$  we get the conclusion.*

ii) More generally than i) if  $\varepsilon = +1$  then a  $F$ -hybrid 2-form is  $F$ -symmetric and for  $\varepsilon = -1$  an  $F$ -pure 2-form is  $F$ -symmetric.  $\square$

Secondly, we associate a conjugate form and an  $(r + 1)$ -form:

**Definition 2.5** If  $\omega \in \Omega^r(M)$  is  $F$ -symmetric then its  $F$ -conjugate is  $\bar{\omega} = \omega_F \in \Omega^r(M)$  given by:

$$\bar{\omega}(X_1, \dots, X_r) := \frac{1}{\varepsilon} \omega(FX_1, \dots, X_r). \tag{2.24}$$

We associate  $\Omega_{F,\omega} \in \Omega^{r+1}(M)$  given by:

$$\Omega_{F,\omega}(X_1, \dots, X_{r+1}) := d\omega(FX_1, \dots, X_{r+1}) - \varepsilon d\bar{\omega}(X_1, \dots, X_{r+1}). \tag{2.25}$$

Thirdly, we define the natural generalization of the previous subsection:

**Definition 2.6** The  $F$ -symmetric form  $\omega \in \Omega^r(M)$  is called almost  $F$ -analytic if:

$$\Omega_{F,\omega} = 0. \tag{2.26}$$

In order to unify the property that says when an  $F$ -symmetric  $r$ -form is almost  $F$ -analytic for both almost complex and paracomplex cases, we present:

**Proposition 2.6** An  $F$ -symmetric  $r$ -form  $\omega$  ( $r \geq 1$ ) is almost  $F$ -analytic iff

$$\begin{aligned} &FX_1(\omega(X_2, \dots, X_{r+1})) - X_1(\omega(FX_2, \dots, X_{r+1})) = \\ &= \sum_{j=2}^{r+1} (-1)^{1+j} \omega(F[X_1, X_j] - [FX_1, X_j], X_2, \dots, \widehat{X_j}, \dots, X_{r+1}). \end{aligned} \tag{2.27}$$

**Proof** It follows by a direct calculation involving the definition of the exterior derivative.  $\square$

**Remark 2.1** In a more general case of  $(0, r)$ -tensor fields we can consider the operator  $\Phi_F : \mathcal{T}_r^0(M) \rightarrow \mathcal{T}_{r+1}^0(M)$ ; see [18]:

$$\begin{aligned} \Phi_F \omega(X, Y_1, \dots, Y_r) &= FX(\omega(Y_1, \dots, Y_r)) - X(\omega(FY_1, Y_2, \dots, Y_r)) \\ &+ \omega((L_{Y_1} F)X, Y_2, \dots, Y_r) + \dots + \omega(Y_1, Y_2, \dots, (L_{Y_r} F)X), \end{aligned} \tag{2.28}$$

for every vector field  $X, Y_1, \dots, Y_r$ , where  $L_X$  denotes the Lie derivative with respect to  $X$ . Then, similarly to [5, 6, 7, 10, 12, 15, 18], the tensor field  $\omega$  is called almost  $F$ -analytic if  $\Phi_F \omega = 0$  and for  $r$ -forms this condition is equivalent to (2.26).

Let  $\Omega^r(M, F)$  be the set of almost  $F$ -analytic  $r$ -forms.

The following result is a motivation for this notion and also a generalization of the first remark above:

**Proposition 2.7** If  $\omega \in \Omega^r(M, F)$  then its differential is  $F$ -symmetric and its exterior  $F$ -differential of  $\omega$  is:

$$d_F \omega = \frac{1}{r + 1} i_F \circ d\omega = \varepsilon d\bar{\omega}. \tag{2.29}$$

**Proof** The first part follows directly from the skew-symmetry of  $d\bar{\omega}$  and the relation:

$$d\omega(FX_1, \dots, X_{r+1}) = \varepsilon d\bar{\omega}(X_1, \dots, X_{r+1}), \tag{2.30}$$

provided by the definition. For the second part we get that  $i_F\omega = \varepsilon r\bar{\omega}$  and with a similar calculus as in Proposition 2.4 we derive:

$$d_F\omega(X_1, \dots, X_{r+1}) = d\omega(FX_1, \dots, X_{r+1}),$$

by using the first part. Equation (2.29) follows then directly. □

**Proposition 2.8** *The  $F$ -symmetric  $r$ -form  $\omega$  is almost  $F$ -analytic if and only if  $\bar{\omega}$  is almost  $F$ -analytic. If  $\omega$  is almost  $F$ -analytic then  $\omega$  is closed if and only if  $\bar{\omega}$  is closed, and equivalently  $\omega$  and  $\bar{\omega}$  are  $d_F$ -closed.*

**Proof** It is sufficient to prove the implication that  $\omega$  is almost  $F$ -analytic  $\Rightarrow$   $\bar{\omega}$  is almost  $F$ -analytic since:

$$\bar{\bar{\omega}}(X_1, \dots, X_r) = \frac{1}{\varepsilon}\bar{\omega}(FX_1, \dots, X_r) = \omega(F^2X_1, \dots, X_r) = \varepsilon\omega(X_1, \dots, X_r) = \frac{1}{\varepsilon}\omega(X_1, \dots, X_r) \tag{2.31}$$

and remark that almost  $F$ -analyticity is invariant with respect to scalings  $\omega \rightarrow \lambda\omega$ .

Firstly we must prove that  $\bar{\omega}$  is  $F$ -symmetric. We have:

$$\bar{\omega}(FX_1, \dots, X_r) = \frac{1}{\varepsilon}\omega(F^2X_1, \dots, X_r) = \omega(X_1, \dots, X_r). \tag{2.32}$$

Also:

$$\begin{aligned} \bar{\omega}(X_1, \dots, FX_i, \dots, X_r) &= \frac{1}{\varepsilon}\omega(FX_1, \dots, FX_i, \dots, X_r) \\ &= \frac{1}{\varepsilon}\omega(X_1, \dots, F^2X_i, \dots, X_r) = \omega(X_1, \dots, X_r), \end{aligned}$$

which is what we claim.

Secondly, we must verify Definition 2.6. A straightforward calculation gives the generalization of (2.6):

$$\Omega_{F,\bar{\omega}}(X_1, \dots, X_{r+1}) = -\frac{1}{\varepsilon}\Omega_{F,\omega}(FX_1, \dots, X_{r+1}) \tag{2.33}$$

and the conclusion follows. □

**Proposition 2.9** *If  $\omega \in \Omega^r(M, F)$  then:*

$$\omega(N_F(X_1, X_2), \dots, X_{r+1}) = \bar{\omega}(N_F(X_1, X_2), \dots, X_{r+1}) = 0. \tag{2.34}$$

**Proof** Using the characterization of almost  $F$ -analyticity of  $\omega$  from (2.27) but with  $X_2 \mapsto FX_2$ , we have

$$\begin{aligned} &FX_1(\omega(FX_2, \dots, X_{r+1})) - \varepsilon X_1(\omega(X_2, \dots, X_{r+1})) = \\ &= -\omega(F[X_1, FX_2] - [FX_1, FX_2], X_3, \dots, X_{r+1}) + \\ &+ \sum_{j=3}^{r+1} (-1)^{1+j}\omega(F[X_1, X_j] - [FX_1, X_j], FX_2, X_3, \dots, \widehat{X_j}, \dots, X_{r+1}). \end{aligned} \tag{2.35}$$

On the other hand,  $\bar{\omega} \in \Omega^r(M, F)$ , too, and using again (2.27) for  $\bar{\omega}$ , we have

$$\begin{aligned} & FX_1(\omega(FX_2, \dots, X_{r+1})) - \varepsilon X_1(\omega(X_2, \dots, X_{r+1})) = \\ & = -\omega(\varepsilon[X_1, X_2] - F[FX_1, X_2], X_3, \dots, X_{r+1}) + \\ & + \sum_{j=3}^{r+1} (-1)^{1+j} \omega(F[X_1, X_j] - [FX_1, X_j], FX_2, X_3, \dots, \widehat{X_j}, \dots, X_{r+1}). \end{aligned} \tag{2.36}$$

Now, by (2.35) and (2.36), the first equality follows easily. The second equality follows in a similar manner.  $\square$

Inspired by the  $\varepsilon = -1$  case, we suppose now that  $m = 2n$  and for the  $\varepsilon = +1$  we suppose that  $F$  is an almost paracomplex structure, i.e. the dimensions of  $(+1)$ -eigenspace and  $(-1)$ -eigenspaces are both equal to  $n$ . It follows for both cases of  $\varepsilon$  the existence of local basis of vector fields of type  $B = \{e_1, \dots, e_n, Fe_1, \dots, Fe_n\}$ , where for the case  $\varepsilon = 1$  we must have  $F \neq \text{Id}$ , and then there exist nontrivial  $F$ -symmetric  $r$  forms only for  $r \leq n$ . An important result for this choice of dimension is:

**Proposition 2.10** *An  $F$ -symmetric  $n$ -form  $\omega$  is almost  $F$ -analytic if and only if  $\omega$  and  $\bar{\omega}$  are both closed.*

**Proof** Suppose firstly that  $\omega$  is almost  $F$ -analytic. When its differential is applied on data  $\{FX_1, X_1, \dots, X_n\}$  of elements of  $B$  we have  $d\omega(FX_1, X_1, \dots, X_n) = \varepsilon d\bar{\omega}(X_1, X_1, \dots, X_n) = 0$  and deduce that  $\omega$  (and consequently  $\bar{\omega}$ ) is closed. The proof of the converse part is directly from Definition 2.25  $\square$

We introduce now an exterior product adapted to our setting:

**Definition 2.7** *The exterior  $F$ -product is the map  $\wedge_F : \Omega^r(M) \times \Omega^s(M) \rightarrow \Omega^{r+s}(M)$  given by:*

$$\theta \wedge_F \omega := \theta \wedge \omega + \varepsilon \bar{\theta} \wedge \bar{\omega} \tag{2.37}$$

where  $\wedge$  is the usual exterior product of  $M$ .

A long but straightforward computation in the basis  $B$  gives:

**Proposition 2.11** *Let  $\theta$  and  $\omega$  be  $F$ -symmetric forms.*

- i) *The  $(r + s)$ -form  $\theta \wedge_F \omega$  is also  $F$ -symmetric.*
- ii) *The  $F$ -conjugate of the  $(r + s)$ -form above is:*

$$(\theta \wedge_F \omega)_F = \theta \wedge \bar{\omega} + \bar{\theta} \wedge \omega. \tag{2.38}$$

As a consequence, if  $\theta$  and  $\omega$  are almost  $F$ -analytic forms then  $\theta \wedge_F \omega$  is also an almost  $F$ -analytic form.

**Proposition 2.12** *Let  $\omega \in \Omega^r(M, F)$  and  $\theta \in \Omega^s(M, F)$ ,  $r, s \geq 0$ , where  $\Omega^0(M, F) = C^\infty(M, F)$ . Then:*

- i)  $d_F \omega \in \Omega^{r+1}(M, F)$ ;
- ii)  $d_F^2 \omega = 0$ ;
- iii)  $d_F(\omega \wedge_F \theta) = d_F \omega \wedge_F \theta + (-1)^r \omega \wedge_F d_F \theta$ .



**Proof** i) If  $\omega \in \Omega^r(M, F)$  then by (2.24) and (2.30) we have:

$$d\bar{\omega} = \overline{d\omega}. \tag{2.39}$$

Now, using (2.29) and (2.39), we have:

$$\Omega_{F, d_F\omega}(X_1, \dots, X_{r+2}) = (d(\varepsilon d\bar{\omega}))(FX_1, \dots, X_{r+2}) - \varepsilon(d(d\omega))(X_1, \dots, X_{r+2}) = 0,$$

which says that  $d_F\omega \in \Omega^{r+1}(M, F)$ .

ii) Using (2.29), (2.31), and (2.39) we have:

$$d_F(d_F\omega) = d_F(\varepsilon d\bar{\omega}) = \varepsilon^2 d(\overline{d\omega}) = \varepsilon^2 d(d\omega) = \varepsilon d(d\omega) = 0.$$

iii) Follows using (2.29), (2.37), and (2.38). □

We notice that  $(\Omega^r(M, F), \wedge_F)$  is a graded  $C^\infty(M, F)$ -algebra. Also, by ii) Proposition 2.12 we have the differential complex  $(\Omega^\bullet(M, F), d_F)$  and its cohomology  $H^\bullet(M, F)$  is called the  $d_F$ -cohomology of almost  $F$ -analytic forms on  $M$ .

Another important property of the operator  $d_F$  is the following Poincaré type lemma:

**Theorem 2.1** *Let  $\omega \in \Omega^r(U, F)$ ,  $r \geq 1$ , where  $U \subset M$  such that  $d_F\omega = 0$  on  $U$ . Then there exists  $\theta \in \Omega^{r-1}(U', F)$  where  $U' \subset U$  such that  $\omega = d_F\theta$  on  $U'$ .*

**Proof** Let  $\omega$  as in the hypothesis. Taking into account that  $d_F\omega = 0$  is equivalent with  $d\bar{\omega} = 0$  and by applying the classical Poincaré lemma for the operator  $d$  it follows that there exists  $\theta \in \Omega^r(U')$  where  $U' \subset U$  and such that  $\bar{\omega} = d\theta$  on  $U'$ . From  $0 = \overline{d\bar{\omega}} = \overline{d\bar{\omega}} = \frac{1}{\varepsilon}d\omega$  it follows also by Poincaré lemma for the operator  $d$  that there exists  $\theta_1 \in \Omega^r(U')$  where  $U' \subset U$  and such that  $\omega = d\theta_1$  on  $U'$ .

Similar arguments as in the proof of Theorem 1 from [9] show that both  $\theta$  and  $\theta_1$  are almost  $F$ -analytic and  $\theta = \overline{\theta_1}$ . Now, the proof follows easily since  $\omega = d\theta_1 = \varepsilon d\bar{\theta} = d_F\theta$ . □

We notice that  $\ker\{d_F : \Omega^0(U, F) \rightarrow \Omega^1(U, F)\} \cong \tilde{\mathbb{R}}$  where  $\tilde{\mathbb{R}}$  is the sheaf of germs associated to the constant pre-sheaf  $\mathbb{R}$ . Also, consider  $\Phi^r(M, F)$  the sheaf of germs of almost  $F$ -analytic  $r$ -forms on  $M$  and  $i : \tilde{\mathbb{R}} \rightarrow \Phi^0(M, F)$  the natural inclusion. The sheaves  $\Phi^r(M, F)$  are fine and taking into account Theorem 2.1 it follows that the following sequence of sheaves:

$$0 \longrightarrow \tilde{\mathbb{R}} \xrightarrow{i} \Phi^0(M, F) \xrightarrow{d_F} \Phi^1(M, F) \xrightarrow{d_F} \dots \xrightarrow{d_F} \Phi^n(M, F) \xrightarrow{d_F} 0$$

is a fine resolution of  $\tilde{\mathbb{R}}$  and we denote by  $H^r(M, F; \tilde{\mathbb{R}})$  the cohomology groups of  $M$  with coefficients in the sheaf  $\tilde{\mathbb{R}}$ . Thus, we obtain a de Rham theorem for the  $d_F$ -cohomology of almost  $F$ -analytic forms, namely:

**Theorem 2.2** *Then  $d_F$ -cohomology groups of almost  $F$ -analytic forms on  $M$  are given by:*

- i)  $H^0(M, F; \tilde{\mathbb{R}}) \cong \mathbb{R}$ ,
- ii)  $H^r(M, F; \tilde{\mathbb{R}}) \cong H^r(M, F)$ ,  $1 \leq r \leq n - 1$ ,
- iii)  $H^n(M, F; \tilde{\mathbb{R}}) \cong \Omega^n(M, F)/d_F(\Omega^{n-1}(M, F))$ ,

iv)  $H^r(M, F; \tilde{\mathbb{R}}) = 0, n + 1 \leq r \leq 2n.$

We consider now a deformation of almost  $F$ -analytic forms with pairs of almost  $F$ -analytic functions:

**Definition 2.8** Fix  $\omega \in \Omega^r(M)$  an almost  $F$ -analytic form and  $\alpha, \beta \in C^\infty(M)$ . The  $(\alpha, \beta)$ -deformation of  $\omega$  is the  $r$ -form:

$$\omega_{\alpha, \beta} := \alpha\omega + \beta\bar{\omega}. \tag{2.40}$$

Since  $\omega_{\alpha, \beta}$  is an  $F$ -symmetric form it is natural to ask in what conditions regarding these functions the new  $r$ -form is also an almost  $F$ -analytic one:

**Proposition 2.13** The  $F$ -symmetric  $r$ -form  $\omega_{\alpha, \beta}$  is almost  $F$ -analytic if and only if  $\alpha$  is almost  $F$ -analytic with corresponding function  $\beta$ .

**Proof** We have:

$$(\omega_{\alpha, \beta})_F = \alpha\bar{\omega} + \frac{\beta}{\varepsilon}\omega = (\omega_F)_{\alpha, \beta}. \tag{2.41}$$

The proof is easy to see in the case  $r = 1$  where the almost  $F$ -analyticity of  $\omega_{\alpha, \beta}$  means:

$$d\alpha(FX)\omega(Y) + \frac{1}{\varepsilon}d\beta(FX)\omega(FY) = d\alpha(X)\omega(FY) + d\beta(X)\omega(Y) \tag{2.42}$$

for all vector fields  $X, Y$ . A detailed proof for  $r \geq 2$  can be found in [3] for the case of almost (para) complex Lie algebroids. □

This results yields the introduction of the set:

$$\tilde{C}^\infty(M, F) = \{(\alpha, \beta) \in C^\infty(M, F) \times C^\infty(M, F); \quad d\beta = d\alpha \circ F\}. \tag{2.43}$$

A straightforward computation gives that  $\tilde{C}^\infty(M, F)$  is a commutative algebra with respect to the product:

$$(\alpha_1, \beta_1) \cdot (\alpha_2, \beta_2) := (\alpha_1\alpha_2 + \varepsilon\beta_1\beta_2, \alpha_1\beta_2 + \alpha_2\beta_1), \tag{2.44}$$

having as a unit the pair of the constant functions  $(1, 0) \in \tilde{C}^\infty(M, F)$ . The inverse of the element  $(\alpha, \beta) \in \tilde{C}^\infty(M, F)$  different from  $(0, 0)$  is the pair  $\left(\frac{\alpha}{\alpha^2 - \varepsilon\beta^2}, \frac{-\beta}{\alpha^2 - \varepsilon\beta^2}\right)$ ; for the case  $\varepsilon = +1$  we also exclude the cases  $(\alpha, \pm\alpha)$ .

Let us introduce the set of pairs of forms:

$$\tilde{\Omega}^r(M, F) = \{(\omega, \bar{\omega}); \omega \in \Omega^r(M, F)\}. \tag{2.45}$$

Proposition 2.13 says that  $\tilde{\Omega}^r(M, F)$  is a  $\tilde{C}^\infty(M, F)$ -module for all  $1 \leq r \leq n$  and hence the set

$$\tilde{\Omega}(M, F) = \sum_{r=1}^n \tilde{\Omega}^r(M, F)$$

is a graded  $\tilde{C}^\infty(M, F)$ -algebra. We consider the wedge product

$$(\omega, \bar{\omega})\tilde{\wedge}(\theta, \bar{\theta}) = (\omega \wedge_F \theta, (\omega \wedge_F \theta)_F) \tag{2.46}$$

and the operator

$$D : \tilde{\Omega}^r(M, F) \rightarrow \tilde{\Omega}^{r+1}(M, F), \quad D(\omega, \bar{\omega}) = (d\omega, d\bar{\omega}). \tag{2.47}$$

It follows that:

- i)  $D$  is a local operator and  $\mathbb{R}$ -linear;
- ii) for every  $(\omega, \bar{\omega}) \in \tilde{\Omega}^r(M, F)$  and  $(\theta, \bar{\theta}) \in \tilde{\Omega}^s(M, F)$  we have

$$D [(\omega, \bar{\omega}) \tilde{\wedge} (\theta, \bar{\theta})] = D(\omega, \bar{\omega}) \tilde{\wedge} (\theta, \bar{\theta}) + (-1)^r (\omega, \bar{\omega}) \tilde{\wedge} D(\theta, \bar{\theta});$$

- iii)  $D^2 = (0, 0)$ ;

and an associated cohomology of the differential complex  $(\tilde{\Omega}(M, F), D)$  can be considered exactly as in [9].

### 3. Almost analytic forms on almost para-Norden manifolds and examples

We continue with the setting of Subsection 2.2, namely  $\varepsilon = \pm 1$ , but we add a Riemannian metric  $g$  to our framework, which satisfies

$$g(FX, Y) = \varepsilon g(X, FY). \tag{3.1}$$

Then:

- a) for  $\varepsilon = -1$  the triple  $(M, F, g)$  is an usual almost Hermitian manifold,
- b) for  $\varepsilon = +1$  the triple  $(M, F, g)$  is an almost para-Norden manifold; see, for instance, [11].

In order to unify these cases we get the following formula:

$$g(FX, FY) = g(X, Y), \quad \forall X, Y \in \mathcal{X}(M). \tag{3.2}$$

The fundamental 2-form of an almost Hermitian manifold is  $\omega(X, Y) := g(X, FY)$ , which is not  $F$ -symmetric, since  $\omega(FX, Y) = -\omega(X, FY)$ , while the symmetric bilinear form  $\omega(X, Y) := g(X, FY)$  associated to an almost para-Norden manifold is  $F$ -symmetric.

The characterization of almost analyticity of differential forms on almost Hermitian manifolds in terms of their harmonicity was studied in [13]. In order to unify these results for both cases presented above, in this section we extend some similar results for the case of almost para-Norden manifolds.

The metric  $g$  yields the Hodge star operator  $\star$  and the orthonormal basis  $B$  of the type discussed above. Hence, similar to the almost Hermitian case, see Proposition 2.3 in [13, p. 77], a direct computation yields:

**Proposition 3.1** *If the  $n$ -form  $\omega$  is  $F$ -symmetric on the almost para-Norden manifold  $(M^{2n}, F, g)$  then  $\star\omega$  is also  $F$ -symmetric.*

The important consequence of this result is:

**Proposition 3.2** *If  $\omega$  is an almost  $F$ -analytic  $n$ -form on the almost para-Norden manifold  $(M^{2n}, F, g)$  then  $\star\omega$  is also almost  $F$ -analytic.*

We arrive now to the main result of this section, which provides a large class of almost  $F$ -analytic forms:

**Proposition 3.3** *An  $F$ -symmetric  $n$ -form on the almost para-Norden manifold  $(M, g, F)$  is almost  $F$ -analytic if and only if  $\omega$  and  $\bar{\omega}$  are both harmonic.*

**Proof** It is a direct consequence of  $d\omega = d(\star\omega) = 0$ . □

Suppose  $n = 2$  and  $\varepsilon = -1$ . By using the corollary 18 of [8, p. 208] it results that on a compact, oriented surface  $M^2$  with positive Ricci (equivalently Gaussian, if  $M$  is embedded in  $\mathbb{R}^3$ ) curvature at one point we have  $\Omega^1(M, F) = 0$ .

We end this section with some examples of (almost)  $F$ -analytic forms. In order to find large classes of almost  $F$ -analytic forms we suppose now that  $F$  is integrable. Then we call  $F$ -analytic forms the differential forms studied until now.

The integrability of  $F$  yields the local coordinates  $\{x^i, y^i; 1 \leq i \leq n\}$  such that the expression of  $F$  is:

$$F\left(\frac{\partial}{\partial x^i}\right) = \frac{\partial}{\partial y^i}, \quad F\left(\frac{\partial}{\partial y^i}\right) = \varepsilon \frac{\partial}{\partial x^i}. \tag{3.3}$$

Let  $\omega = a_i dx^i + b_i dy^i$  be a 1-form on  $M$ ; hence,  $\bar{\omega} = \varepsilon b_i dx^i + a_i dy^i$ . The  $F$ -analyticity of  $\omega$  means:

$$FX(\omega(Y)) - \omega([FX, Y]) = X(\omega(FY)) - \omega(F[X, Y]), \tag{3.4}$$

and the choice of  $X, Y$  in the basis  $\{\frac{\partial}{\partial x^i}, \frac{\partial}{\partial y^i}; 1 \leq i \leq n\}$  gives the following characterization:

**Theorem 3.1** *The 1-form  $\omega$  is an  $F$ -analytic form if and only if its coefficients satisfy the  $\varepsilon$ -Cauchy–Riemann equations:*

$$\frac{\partial a_j}{\partial y^i} = \frac{\partial b_j}{\partial x^i}, \quad \frac{\partial a_j}{\partial x^i} = \varepsilon \frac{\partial b_j}{\partial y^i}. \tag{3.5}$$

Similarly, the pair of smooth functions  $(\alpha, \beta)$  belongs to  $\tilde{C}^\infty(M, F)$  if and only if  $\alpha$  and  $\beta$  satisfies the  $\varepsilon$ -Cauchy–Riemann equations (3.5).

A natural framework where quadratic endomorphisms are involved is provided by  $\varepsilon$ -contact structures, namely triples  $(\varphi, \xi, \eta)$  consisting of an endomorphism, a vector field, and a 1-form on  $M^{2n+1}$  satisfying:

$$\varphi^2 = \varepsilon(I_M - \eta \otimes \xi), \quad \eta(\xi) = 1. \tag{3.6}$$

For  $\varepsilon = -1$  we get the almost contact geometry [2], while for  $\varepsilon = +1$  we have the almost paracontact geometry [19]. On the product manifold  $M \times \mathbb{R}$  we consider:

$$J(X, a \frac{d}{dt}) = (\varphi X + \varepsilon a \xi, \eta(X) \frac{d}{dt}), \tag{3.7}$$

and a straightforward computation yields that  $J^2 = \varepsilon I_{M \times \mathbb{R}}$ . For the 1-form  $\omega_b = \eta + bdt$  with  $b \in \mathbb{R}$ , its conjugate with respect to  $J$  is:

$$(\omega_b)_J = \varepsilon b \eta + dt, \tag{3.8}$$

and then  $\omega_b$  is almost  $J$ -analytic form if and only if:

$$d\eta(\varphi X, Y) = b\varepsilon d\eta(X, Y) \tag{3.9}$$

for all vector fields  $X, Y$ . In particular, if  $(M, \varphi, \xi, \eta)$  is  $\varepsilon$ -cosymplectic, i.e.  $\eta$  is closed, then all  $\omega_b$  are almost  $J$ -analytic.

## Acknowledgment

We are extremely indebted to two anonymous referees who improved the initial submission. The second author was supported by the Sectorial Operational Program Human Resources Development (SOP HRD), financed by the European Social Fund and by the Romanian Government under project number POSDRU/159/1.5/S/134378.

## References

- [1] Alekseevsky DA, Medori C, Tomassini A. Maximally homogeneous para-CR manifolds of semisimple type. In: Cortes V, editor. *Handbook of Pseudo-Riemannian Geometry and Supersymmetry*. Zurich, Switzerland: European Mathematical Society, 2010, pp. 559–577.
- [2] Blair DE. *Riemannian Geometry of Contact and Symplectic Manifolds*. Progress in Mathematics, Vol. 203. 2nd ed. Boston, MA, USA: Birkhäuser, 2010.
- [3] Crasmareanu M, Ida C. Almost analyticity on almost (para) complex Lie algebroids. *Results in Math* 2015 (in press).
- [4] Cruceanu V, Fortuny P, Gadea PM. A survey on paracomplex geometry. *Rocky Mountain J Math* 1996; 26: 83–115.
- [5] Gezer A, Cengiz N, Salimov AA. On integrability of golden Riemannian structures. *Turk J Math* 2013; 37: 693–703.
- [6] Iscan M, Salimov AA. On Kähler-Norden manifolds. *P Indian AS-Math Sci* 2009; 119: 71–80.
- [7] Kruchkovich GI. Hypercomplex structure on a manifold. *Tr Sem Vect Tens Anal Moscow Univ* 1972; 16: 174–201.
- [8] Petersen P. *Riemannian Geometry*. Graduate Texts in Mathematics, Vol. 171. 2nd ed. New York, NY, USA: Springer, 2006.
- [9] Pitis G, Smaranda D. Formes presque analytiques sur une variété presque complexe. *Zb Rad Prirod-Mat Fak Ser Mat* 1991; 21: 2329 (in French).
- [10] Salimov AA, Gezer A. Norden structures of Hessian type. *Turk J Math* 2014; 38: 462–469.
- [11] Salimov AA, Iscan M, Akbulut K. Notes on para-Norden-Walker 4-manifolds. *Int J Geom Methods Mod Phys* 2010; 7: 1331–1347.
- [12] Salimov AA, Iscan M, Etayo F. Paraholomorphic  $B$ -manifold and its properties. *Topol Appl* 2007; 154: 925–933.
- [13] Simionescu C, Pitis G. Some properties of almost analytic forms. *Bull Univ Brasov Ser C* 1989; 31: 75–80.
- [14] Smaranda D. The algebra of almost analytic forms and criteria of almost analyticity. In: *Special Chapters in Differential Geometry*. Bucharest, Romania: University of Bucharest, 1981, pp. 67–87 (in Romanian).
- [15] Tachibana S. Analytic tensor and its generalization. *Tôhoku Math J* 1960; 12: 208–221.
- [16] Tachibana S, Kotô S. On almost-analytic functions, tensors and invariant subspaces. *Tôhoku Math J* 1962; 14: 177–186.
- [17] Yano K. *Differential Geometry on Complex and Almost Complex Spaces*. International Series of Monographs in Pure and Applied Mathematics, Vol. 49. New York, NY, USA: Pergamon Press, 1965.
- [18] Yano K, Ako M. On certain operators associated with tensor fields. *Kodai Math Sem Rep* 1968; 20: 414–436.
- [19] Zamkovoy S. Canonical connections on paracontact manifolds. *Ann Glob Anal Geom* 2008; 36: 37–60.