

A further extension of the extended Riemann–Liouville fractional derivative operator

Martin BOHNER¹, Gauhar RAHMAN², Shahid MUBEEN³, Kottakkaran Sooppy NISAR^{4,*}

¹Department of Mathematics, Missouri University of Science and Technology, Rolla, MO, USA

²Department of Mathematics, International Islamic University, Islamabad, Pakistan

³Department of Mathematics, University of Sargodha, Sargodha, Pakistan

⁴Department of Mathematics, College of Arts and Science-Wadi Aldawaser, Prince Sattam bin Abdulaziz University, Alkharj, Saudi Arabia

Received: 29.05.2018

Accepted/Published Online: 28.07.2018

Final Version: 27.09.2018

Abstract: The main objective of this paper is to establish the extension of an extended fractional derivative operator by using an extended beta function recently defined by Parmar et al. by considering the Bessel functions in its kernel. We also give some results related to the newly defined fractional operator, such as Mellin transform and relations to extended hypergeometric and Appell's function via generating functions.

Key words: Hypergeometric function, extended hypergeometric function, Mellin transform, fractional derivative, Appell's function

1. Introduction

Recently, the applications and importance of fractional calculus have received more attention. In the field of mathematical analysis, fractional calculus is a powerful tool. Various extensions and generalizations of fractional derivative operators were recently investigated in [7, 8, 10, 16].

Euler's beta function is defined by

$$B(\sigma_1, \sigma_2) = \int_0^1 t^{\sigma_1-1} (1-t)^{\sigma_2-1} dt, \quad \operatorname{Re}(\sigma_1) > 0, \quad \operatorname{Re}(\sigma_2) > 0, \quad (1.1)$$

and its relation with the gamma function is given by

$$B(\sigma_1, \sigma_2) = \frac{\Gamma(\sigma_1)\Gamma(\sigma_2)}{\Gamma(\sigma_1 + \sigma_2)}.$$

The Gauss hypergeometric and the confluent hypergeometric functions are defined (see [15]) by

$${}_2F_1(\sigma_1, \sigma_2; \sigma_3; z) = \sum_{n=0}^{\infty} \frac{(\sigma_1)_n (\sigma_2)_n}{(\sigma_3)_n} \frac{z^n}{n!}, \quad |z| < 1, \quad \sigma_1, \sigma_2, \sigma_3 \in \mathbb{C}, \quad -\sigma_3 \notin \mathbb{N}_0 \quad (1.2)$$

*Correspondence: n.sooppy@psau.edu.sa

2010 AMS Mathematics Subject Classification: 33C15, 33C05

and

$${}_1\Phi_1(\sigma_2; \sigma_3; z) = \sum_{n=0}^{\infty} \frac{(\sigma_2)_n z^n}{(\sigma_3)_n n!}, \quad |z| < 1, \quad \sigma_2, \sigma_3 \in \mathbb{C}, -\sigma_3 \notin \mathbb{N}_0. \quad (1.3)$$

The Appell series or bivariate hypergeometric series is defined by

$$F_1(\sigma_1, \sigma_2, \sigma_3; \sigma_4; x, y) = \sum_{m,n=0}^{\infty} \frac{(\sigma_1)_{m+n} (\sigma_2)_m (\sigma_3)_n x^m y^n}{(\sigma_4)_{m+n} m! n!}, \quad (1.4)$$

for all $\sigma_1, \sigma_2, \sigma_3, \sigma_4 \in \mathbb{C}$, $-\sigma_4 \notin \mathbb{N}_0$, $|x| < 1$, $|y| < 1$. The integral representations of (1.2), (1.3), and (1.4) are respectively given by

$${}_2F_1(\sigma_1, \sigma_2; \sigma_3; z) = \frac{\Gamma(\sigma_3)}{\Gamma(\sigma_2)\Gamma(\sigma_3 - \sigma_2)} \int_0^1 t^{\sigma_2-1} (1-t)^{\sigma_3-\sigma_2-1} (1-zt)^{-\sigma_1} dt, \quad (1.5)$$

$\operatorname{Re}(\sigma_3) > \operatorname{Re}(\sigma_2) > 0$, $|\arg(1-z)| < \pi$, and

$${}_1\Phi_1(\sigma_2; \sigma_3; z) = \frac{\Gamma(\sigma_3)}{\Gamma(\sigma_2)\Gamma(\sigma_3 - \sigma_2)} \int_0^1 t^{\sigma_2-1} (1-t)^{\sigma_3-\sigma_2-1} e^{zt} dt, \quad (1.6)$$

$\operatorname{Re}(\sigma_3) > \operatorname{Re}(\sigma_2) > 0$,

$$F_1(\sigma_1, \sigma_2, \sigma_3; \sigma_4; x, y) = \frac{\Gamma(\sigma_4)}{\Gamma(\sigma_1)\Gamma(\sigma_4 - \sigma_1)} \int_0^1 t^{\sigma_1-1} (1-t)^{\sigma_4-\sigma_1-1} (1-xt)^{-\sigma_2} (1-yt)^{-\sigma_3} dt. \quad (1.7)$$

Chaudhry et al. [2] introduced the extended beta function as

$$B(\sigma_1, \sigma_2; p) = B_p(\sigma_1, \sigma_2) = \int_0^1 t^{\sigma_1-1} (1-t)^{\sigma_2-1} e^{-\frac{p}{t(1-t)}} dt, \quad (1.8)$$

where $\operatorname{Re}(p) > 0$, $\operatorname{Re}(\sigma_1) > 0$, $\operatorname{Re}(\sigma_2) > 0$. If $p = 0$, then $B(\sigma_1, \sigma_2; 0) = B(\sigma_1, \sigma_2)$. The extended hypergeometric and confluent hypergeometric functions are defined in [3] as

$$F_p(\sigma_1, \sigma_2; \sigma_3; z) = \sum_{n=0}^{\infty} \frac{B_p(\sigma_2 + n, \sigma_3 - \sigma_2)}{B(\sigma_2, \sigma_3 - \sigma_2)} (\sigma_1)_n \frac{z^n}{n!} \quad (1.9)$$

and

$$\Phi_p(\sigma_2; \sigma_3; z) = \sum_{n=0}^{\infty} \frac{B_p(\sigma_2 + n, \sigma_3 - \sigma_2)}{B(\sigma_2, \sigma_3 - \sigma_2)} \frac{z^n}{n!}, \quad (1.10)$$

where $p \geq 0$. Also, in [3], the authors gave the integral representations of the extended hypergeometric and confluent hypergeometric functions as

$$F_p(\sigma_1, \sigma_2; \sigma_3; z) = \frac{1}{B(\sigma_2, \sigma_3 - \sigma_2)} \int_0^1 t^{\sigma_2-1} (1-t)^{\sigma_3-\sigma_2-1} (1-zt)^{-\sigma_1} \exp\left(\frac{-p}{t(1-t)}\right) dt, \quad (1.11)$$

$$p \geq 0, \quad \operatorname{Re}(\sigma_3) > \operatorname{Re}(\sigma_2) > 0, \quad |\arg(1-z)| < \pi$$

and

$$\Phi_p(\sigma_2, \sigma_3; \gamma; z) = \frac{1}{\beta(\sigma_2, \sigma_3 - \sigma_2)} \int_0^1 t^{\sigma_2-1} (1-t)^{\sigma_3-\sigma_2-1} \exp\left(zt - \frac{p}{t(1-t)}\right) dt, \quad (1.12)$$

$$p \geq 0, \quad \operatorname{Re}(\sigma_3) > \operatorname{Re}(\sigma_2) > 0.$$

The extended Appell function is defined (see [11]) by

$$F_1(\sigma_1, \sigma_2, \sigma_3; \sigma_4; x, y; p) = \sum_{n=0}^{\infty} \frac{B_p(\sigma_1 + m + n, \sigma_4 - \sigma_1)}{B(\sigma_1, \sigma_4 - \sigma_1)} (\sigma_2)_m (\sigma_3)_n \frac{x^m y^n}{m! n!}, \quad (1.13)$$

where $p \geq 0$, and its integral representation is given by

$$F_1(\sigma_1, \sigma_2, \sigma_3; \sigma_4; x, y; p) = \frac{1}{B(\sigma_1, \sigma_4 - \sigma_1)} \int_0^1 t^{\sigma_1-1} (1-t)^{\sigma_4-\sigma_1-1} (1-xt)^{-\sigma_2} (1-yt)^{-\sigma_3} \exp\left(\frac{-p}{t(1-t)}\right) dt, \quad (1.14)$$

$$p \geq 0, \quad \operatorname{Re}(\sigma_4) > \operatorname{Re}(\sigma_1) > 0, \quad |\arg(1-x)| < \pi, \quad |\arg(1-y)| < \pi.$$

It is clear that if $p = 0$, then (1.9)–(1.14) reduce to the well-known hypergeometric, confluent hypergeometric, and Appell series and their integral representations, respectively.

For various extensions and generalizations, the readers may follow the recent work [1, 4, 9]. Parmar et al. [13] introduced the extended beta function as

$$\beta_v(\sigma_1, \sigma_2; p) = \sqrt{\frac{2p}{\pi}} \int_0^1 t^{\sigma_1-\frac{3}{2}} (1-t)^{\sigma_2-\frac{3}{2}} K_{v+\frac{1}{2}}\left(\frac{p}{t(1-t)}\right) dt, \quad (1.15)$$

where $K_{v+\frac{1}{2}}(\cdot)$ is the modified Bessel function of order $v + \frac{1}{2}$. Clearly, when $v = 0$, (1.15) reduces to (1.8) by using the fact that $K_{\frac{1}{2}}(z) = \sqrt{\frac{\pi}{2z}} e^{-z}$. Also, the extended hypergeometric and confluent hypergeometric functions, and their integral representations, are as given in [13]:

$$F_{p,v}(\sigma_1, \sigma_2; \sigma_3; z) = \sum_{n=0}^{\infty} (\sigma_1)_n \frac{\beta_v(\sigma_2 + n, \sigma_3 - \sigma_2; p)}{\beta(\sigma_2, \sigma_3 - \sigma_2)} \frac{z^n}{n!}, \quad (1.16)$$

$$p, v \geq 0, \quad \operatorname{Re}(\sigma_3) > \operatorname{Re}(\sigma_2) > 0, \quad |z| < 1,$$

$$\Phi_{p,v}(\sigma_2; \sigma_3; z) = \sum_{n=0}^{\infty} \frac{\beta_v(\sigma_2 + n, \sigma_3 - \sigma_2; p)}{\beta(\sigma_2, \sigma_3 - \sigma_2)} \frac{z^n}{n!}, \quad (1.17)$$

$$p, v \geq 0, \quad \operatorname{Re}(\sigma_3) > \operatorname{Re}(\sigma_2) > 0,$$

$$F_{p,v}(\sigma_1, \sigma_2; \sigma_3; z) = \sqrt{\frac{2p}{\pi}} \frac{1}{\beta(\sigma_2, \sigma_3 - \sigma_2)} \int_0^1 t^{\sigma_2-\frac{3}{2}} (1-t)^{\sigma_3-\sigma_2-\frac{3}{2}} (1-zt)^{-\sigma_1} K_{v+\frac{1}{2}}\left(\frac{p}{t(1-t)}\right) dt, \quad (1.18)$$

$$p, v \geq 0, \quad \operatorname{Re}(\sigma_3) > \operatorname{Re}(\sigma_2) > 0, \quad |\arg(1-z)| < \pi,$$

and

$$\Phi_{p,v}(\sigma_2; \sigma_3; z) = \sqrt{\frac{2p}{\pi}} \frac{1}{\beta(\sigma_2, \sigma_3 - \sigma_2)} \int_0^1 t^{\sigma_2 - \frac{3}{2}} (1-t)^{\sigma_3 - \sigma_2 - \frac{3}{2}} \exp(zt) K_{v+\frac{1}{2}} \left(\frac{p}{t(1-t)} \right) dt, \quad (1.19)$$

$$p, v \geq 0, \quad \operatorname{Re}(\sigma_3) > \operatorname{Re}(\sigma_2) > 0.$$

They also obtained the transformation formula for extended confluent hypergeometric functions as

$$\Phi_{p,v}(\sigma_2, \sigma_3; \gamma; z) = e^z \Phi_{p,v}(\sigma_3 - \sigma_2; \sigma_3; -z). \quad (1.20)$$

It is clear that, when $v = 0$, (1.16)–(1.19) reduce to the extended hypergeometric and confluent hypergeometric functions, and their integral representations are given in (1.9)–(1.12) by using the fact that $K_{\frac{1}{2}}(z) = \sqrt{\frac{\pi}{2z}} e^{-z}$.

Recently, Dar and Paris [5] introduced Appell's hypergeometric function by

$$\begin{aligned} F_{1,p,v}(\sigma_1, \sigma_2, \sigma_3; \sigma_4; x, y) &= F_{1,v}(\sigma_1, \sigma_2, \sigma_3; \sigma_4; x, y; p) \\ &= \sum_{m,n=0}^{\infty} (\sigma_2)_m (\sigma_3)_n \frac{\beta(\sigma_1 + m + n, \sigma_4 - \sigma_1)}{\beta(\sigma_1, \sigma_4 - \sigma_1)} \frac{x^m y^n}{n! m!}, \end{aligned} \quad (1.21)$$

where $|x| < 1$, $|y| < 1$, $\sigma_1, \sigma_2, \sigma_3, \sigma_4 \in \mathbb{C}$, $-\sigma_4 \notin \mathbb{N}_0$.

In the same paper [5], they gave its integral representation as

$$\begin{aligned} &F_{1,p,v}(\sigma_1, \sigma_2, \sigma_3; \sigma_4; x, y) \\ &= \sqrt{\frac{2p}{\pi}} \frac{1}{\beta(\sigma_1, \sigma_4 - \sigma_1)} \int_0^1 t^{\sigma_1 - \frac{3}{2}} (1-t)^{\sigma_4 - \sigma_1 - \frac{3}{2}} (1-tx)^{-\sigma_2} (1-ty)^{-\sigma_3} K_{v+\frac{1}{2}} \left(\frac{p}{t(1-t)} \right) dt, \end{aligned} \quad (1.22)$$

where $\operatorname{Re}(p) \geq 0$, $v \geq 0$, $\operatorname{Re}(\sigma_4) > \operatorname{Re}(\sigma_1) > 0$, $|\arg(1-x)| < \pi$, and $|\arg(1-y)| < \pi$. Obviously, when $v = 0$ in (1.21) and (1.22), we get the extended Appell function and its integral representation (see (1.13) and (1.14)) by using the fact that $K_{\frac{1}{2}}(z) = \sqrt{\frac{\pi}{2z}} e^{-z}$. Similarly, when $v = 0$ and $p = 0$, (1.21) and (1.22) reduce to the well-known classical Appell function and its integral representation.

2. Extension of the fractional derivative operator

In this section, we define a further extension of the extended Riemann–Liouville fractional derivative.

Definition 2.1 *The Riemann–Liouville fractional derivative of order μ is defined by*

$$\mathfrak{D}_x^\mu \{f(x)\} = \frac{1}{\Gamma(-\mu)} \int_0^x f(t) (x-t)^{-\mu-1} dt, \quad \operatorname{Re}(\mu) > 0. \quad (2.1)$$

For the case $m-1 < \operatorname{Re}(\mu) < m$, where $m \in \mathbb{N}$, we have

$$\mathfrak{D}_x^\mu \{f(z)\} = \frac{d^m}{dx^m} \mathfrak{D}_x^{\mu-m} \{f(x)\} = \frac{d^m}{dx^m} \left\{ \frac{1}{\Gamma(-\mu+m)} \int_0^x f(t) (x-t)^{-\mu+m-1} dt \right\}, \quad \operatorname{Re}(\mu) > 0. \quad (2.2)$$

Definition 2.2 (See [11]) *The extended Riemann–Liouville fractional derivative of order μ is defined by*

$$\mathfrak{D}_x^\mu \{f(x); p\} = \frac{1}{\Gamma(-\mu)} \int_0^x f(t)(x-t)^{-\mu-1} \exp\left(-\frac{px^2}{t(x-t)}\right) dt, \quad \operatorname{Re}(\mu) > 0. \quad (2.3)$$

For the case $m-1 < \operatorname{Re}(\mu) < m$, where $m \in \mathbb{N}$, we have

$$\begin{aligned} \mathfrak{D}_x^\mu \{f(z); p\} &= \frac{d^m}{dx^m} \mathfrak{D}_x^{\mu-m} \{f(x); p\} \\ &= \frac{d^m}{dx^m} \left\{ \frac{1}{\Gamma(-\mu+m)} \int_0^x f(t)(x-t)^{-\mu+m-1} \exp\left(-\frac{px^2}{t(x-t)}\right) dt \right\}, \quad \operatorname{Re}(\mu) > 0. \end{aligned} \quad (2.4)$$

Definition 2.3 (See [1]) *We define*

$$\mathfrak{D}_x^\mu \{f(x); p, q\} = \frac{1}{\Gamma(-\mu)} \int_0^x f(t)(x-t)^{-\mu-1} \exp\left(-\frac{px}{t} - \frac{qx}{(x-t)}\right) dt, \quad \operatorname{Re}(\mu) > 0. \quad (2.5)$$

For the case $m-1 < \operatorname{Re}(\mu) < m$, where $m \in \mathbb{N}$, we have

$$\begin{aligned} \mathfrak{D}_x^\mu \{f(z); p, q\} &= \frac{d^m}{dx^m} \mathfrak{D}_x^{\mu-m} \{f(x); p, q\} \\ &= \frac{d^m}{dx^m} \left\{ \frac{1}{\Gamma(-\mu+m)} \int_0^x f(t)(x-t)^{-\mu+m-1} \exp\left(-\frac{px}{t} - \frac{qx}{(x-t)}\right) dt \right\}, \quad \operatorname{Re}(\mu) > 0. \end{aligned} \quad (2.6)$$

A new extension of the Riemann–Liouville fractional derivative of order μ as follows.

Definition 2.4 *The extension of the extended Riemann–Liouville fractional derivative of order μ is defined as*

$$\mathfrak{D}_x^\mu \{f(x); p, q, \lambda, \rho\} = \frac{1}{\Gamma(-\mu)} \int_0^x f(t)(x-t)^{-\mu-1} {}_1F_1\left[\lambda; \rho; -\frac{px}{t}\right] {}_1F_1\left[\lambda; \rho; -\frac{qx}{(x-t)}\right] dt, \quad \operatorname{Re}(\mu) > 0. \quad (2.7)$$

For the case $m-1 < \operatorname{Re}(\mu) < m$, where $m \in \mathbb{N}$, we have

$$\begin{aligned} \mathfrak{D}_x^\mu \{f(z); p, q, \lambda, \rho\} &= \frac{d^m}{dx^m} \mathfrak{D}_x^{\mu-m} \{f(x); p, q, \lambda, \rho\} \\ &= \frac{d^m}{dx^m} \left\{ \frac{1}{\Gamma(-\mu+m)} \int_0^x f(t)(x-t)^{-\mu+m-1} {}_1F_1\left[\lambda; \rho; -\frac{px}{t}\right] {}_1F_1\left[\lambda; \rho; -\frac{qx}{(x-t)}\right] dt \right\}, \end{aligned} \quad (2.8)$$

where $\operatorname{Re}(\mu) > 0$, $\operatorname{Re}(p) > 0$, and $\operatorname{Re}(q) > 0$.

Next, we give an extension of the extended Riemann–Liouville fractional derivative operator (2.2) of order μ as follows.

Definition 2.5 *The extension of the extended Riemann–Liouville fractional derivative of order μ is defined as*

$$\mathfrak{D}_x^\mu \{f(x); p, v\} = \sqrt{\frac{2px^2}{\pi}} \frac{1}{\Gamma(-\mu)} \int_0^x f(t)t^{-\frac{1}{2}}(x-t)^{-\mu-\frac{3}{2}} K_{v+\frac{1}{2}}\left(\frac{px^2}{t(x-t)}\right) dt, \quad \operatorname{Re}(\mu) > 0. \quad (2.9)$$

For the case $m - 1 < \operatorname{Re}(\mu) < m$, where $m \in \mathbb{N}$, it follows that

$$\begin{aligned} \mathfrak{D}_x^\mu \{f(x); p, v\} &= \frac{d^m}{dx^m} \mathfrak{D}_x^{\mu-m} \{f(x); p, v\} \\ &= \frac{d^m}{dx^m} \left\{ \sqrt{\frac{2px^2}{\pi}} \frac{1}{\Gamma(-\mu+m)} \int_0^x f(t) t^{-\frac{1}{2}} (x-t)^{-\mu+m-\frac{3}{2}} K_{v+\frac{1}{2}} \left(\frac{px^2}{t(x-t)} \right) dt \right\}, \end{aligned} \quad (2.10)$$

where $\operatorname{Re}(\mu) > 0$, $\operatorname{Re}(p) > 0$ and $v \geq 0$.

Remark 2.1 *The following observations are clear.*

(i) If $v = 0$, then Definition 2.5 reduces to the extended fractional derivative defined in Definition 2.2 by using the fact that $K_{\frac{1}{2}}(z) = \sqrt{\frac{\pi}{2z}} e^{-z}$.

(ii) If $v = 0$ and $p = 0$, then Definition 2.5 reduces to the Riemann–Liouville fractional derivative defined in Definition 2.1.

Now we prove a result concerning the extension of the fractional derivative.

Theorem 2.1 *We have*

$$\mathfrak{D}_z^\mu \{z^\eta; p, v\} = \frac{\beta_v(\eta+1, -\mu; p)}{\Gamma(-\mu)} z^{\eta-\mu}, \quad \operatorname{Re}(\mu) > 0. \quad (2.11)$$

Proof From (2.9), we have

$$\begin{aligned} \mathfrak{D}_z^\mu \{z^\eta; p, v\} &= \sqrt{\frac{2pz^2}{\pi}} \frac{1}{\Gamma(-\mu)} \int_0^z t^{\eta-\frac{1}{2}} (z-t)^{-\mu-\frac{3}{2}} K_{v+\frac{1}{2}} \left(\frac{pz^2}{t(z-t)} \right) dt \\ &\stackrel{(t=zu)}{=} \sqrt{\frac{2pz^2}{\pi}} \frac{1}{\Gamma(-\mu)} \int_0^1 (uz)^{\eta-\frac{1}{2}} (z-uz)^{-\mu-\frac{3}{2}} K_{v+\frac{1}{2}} \left(\frac{pz^2}{uz(z-uz)} \right) z du \\ &= \frac{z^{\eta-\mu} \sqrt{\frac{2p}{\pi}}}{\Gamma(-\mu)} \int_0^1 u^{\eta-\frac{1}{2}} (1-u)^{-\mu-\frac{3}{2}} K_{v+\frac{1}{2}} \left(\frac{p}{u(1-u)} \right) du, \end{aligned}$$

which shows (2.11). □

Theorem 2.2 *Let $\operatorname{Re}(\mu) > 0$ and suppose that f is analytic at the origin with its Maclaurin expansion given by $f(z) = \sum_{n=0}^{\infty} a_n z^n$, where $|z| < \delta$ for some $\delta > 0$. Then*

$$\mathfrak{D}_z^\mu \{f(z); p, v\} = \sum_{n=0}^{\infty} a_n \mathfrak{D}_z^\mu \{z^n; p, v\}. \quad (2.12)$$

Proof Using the series expansion of f in (2.7) gives

$$\mathfrak{D}_z^\mu \{f(z); p, v\} = \sqrt{\frac{2pz^2}{\pi}} \frac{1}{\Gamma(-\mu)} \int_0^z \sum_{n=0}^{\infty} a_n t^{n-\frac{1}{2}} (z-t)^{-\mu-\frac{3}{2}} K_{v+\frac{1}{2}} \left(\frac{pz^2}{t(z-t)} \right) dt.$$

As the series is uniformly convergent on any closed disk centered at the origin with its radius smaller than δ , the series is on the line segment from 0 to a fixed z for $|z| < \delta$. Thus, we may integrate term by term to find

$$\begin{aligned}\mathfrak{D}_z^\mu\{f(z); p, v\} &= \sum_{n=0}^{\infty} a_n \left\{ \sqrt{\frac{2pz^2}{\pi}} \frac{1}{\Gamma(-\mu)} \int_0^z t^{n-\frac{1}{2}} (z-t)^{-\mu-\frac{3}{2}} K_{v+\frac{1}{2}} \left(\frac{pz^2}{t(z-t)} \right) dt \right\} \\ &= \sum_{n=0}^{\infty} a_n \mathfrak{D}_z^\mu\{z^n; p, v\},\end{aligned}$$

which shows (2.12). □

Theorem 2.3 *We have*

$$\mathfrak{D}_z^{\eta-\mu}\{z^{\eta-1}(1-z)^{-\beta}; p, v\} = \frac{\Gamma(\eta)}{\Gamma(\mu)} z^{\mu-1} {}_2F_{1; p, v}^{\lambda, \rho}(\beta, \eta; \mu; z), \quad (2.13)$$

where $\operatorname{Re}(\mu) > \operatorname{Re}(\eta) > 0$ and $|z| < 1$.

Proof By direct calculation, we have

$$\begin{aligned}\mathfrak{D}_z^{\eta-\mu}\{z^{\eta-1}(1-z)^{-\beta}; p, v\} &= \sqrt{\frac{2pz^2}{\pi}} \frac{1}{\Gamma(\mu-\eta)} \int_0^z t^{\eta-\frac{3}{2}} (1-t)^{-\beta} (z-t)^{\mu-\eta-\frac{3}{2}} K_{v+\frac{1}{2}} \left(\frac{pz^2}{t(z-t)} \right) dt \\ &= \sqrt{\frac{2pz^2}{\pi}} \frac{z^{\mu-\eta-\frac{3}{2}}}{\Gamma(\mu-\eta)} \int_0^z t^{\eta-\frac{3}{2}} (1-t)^{-\beta} \left(1-\frac{t}{z}\right)^{\mu-\eta-\frac{3}{2}} K_{v+\frac{1}{2}} \left(\frac{pz^2}{t(z-t)} \right) dt. \\ &\stackrel{(t=zu)}{=} \frac{z^{\mu-1}}{\Gamma(\mu-\eta)} \sqrt{\frac{2p}{\pi}} \int_0^1 u^{\eta-\frac{3}{2}} (1-uz)^{-\beta} (1-u)^{\mu-\eta-\frac{3}{2}} K_{v+\frac{1}{2}} \left(\frac{p}{u(1-u)} \right) du.\end{aligned}$$

Using (1.16) and after simplification, we get the required proof. □

Theorem 2.4 *We have*

$$\mathfrak{D}_z^{\eta-\mu}\{z^{\eta-1}(1-az)^{-\alpha}(1-bz)^{-\beta}; p, v\} = \frac{\Gamma(\eta)}{\Gamma(\mu)} z^{\mu-1} F_1(\eta, \alpha, \beta; \mu; az, bz; p, v), \quad (2.14)$$

where $\operatorname{Re}(\mu) > \operatorname{Re}(\eta) > 0$, $\operatorname{Re}(\alpha) > 0$, $\operatorname{Re}(\beta) > 0$, $|az| < 1$, and $|bz| < 1$.

Proof To prove (2.14), we use the power series expansion

$$(1-az)^{-\alpha}(1-bz)^{-\beta} = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} (\alpha)_m (\beta)_n \frac{(az)^m}{m!} \frac{(bz)^n}{n!}.$$

Now, applying Theorem 2.3, we obtain

$$\begin{aligned}\mathfrak{D}_z^{\eta-\mu}\{z^{\eta-1}(1-az)^{-\alpha}(1-bz)^{-\beta}; p, v\} &= \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} (\alpha)_m (\beta)_n \frac{(a)^m}{m!} \frac{(b)^n}{n!} \mathfrak{D}_z^{\eta-\mu}\{z^{\eta+m+n-1}; p, v\} \\ &= \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} (\alpha)_m (\beta)_n \frac{(a)^m}{m!} \frac{(b)^n}{n!} \frac{\beta_{p, v}(\eta+m+n, \mu-\eta)}{\Gamma(\mu-\eta)} z^{\mu+m+n-1}.\end{aligned}$$

Now, applying (1.21), we get

$$\mathfrak{D}_z^{\eta-\mu}\{z^{\eta-1}(1-az)^{-\alpha}(1-bz)^{-\beta}; p, v\} = \frac{\Gamma(\eta)}{\Gamma(\mu)} z^{\mu-1} F_1(\eta, \alpha, \beta; \mu; az, bz; p, v),$$

which shows (2.14). \square

Theorem 2.5 *We have the Mellin transform formula*

$$M\{\mathfrak{D}_{z;p,v}^\mu(z^\eta); p \rightarrow r\} = \frac{z^{\eta-\mu} 2^{r-1} \Gamma(\frac{r-v}{2}) \Gamma(\frac{r+v+1}{2})}{\sqrt{\pi} \Gamma(-\mu)} \beta(\eta+r+1, r-\mu), \quad (2.15)$$

where $\operatorname{Re}(\eta) > -1$, $\operatorname{Re}(\mu) > 0$, $\operatorname{Re}(r) > 0$.

Proof Applying the Mellin transform to (2.9), we have

$$\begin{aligned} M\{\mathfrak{D}_{z;p,v}^\mu(z^\eta); p \rightarrow r\} &= \int_0^\infty p^{r-1} \mathfrak{D}_{z;p,v}^\mu(z^\eta) dp \\ &= \sqrt{\frac{2pz^2}{\pi}} \frac{1}{\Gamma(-\mu)} \int_0^\infty p^{r-1} \left\{ \int_0^z t^{-\frac{1}{2}} t^\eta (z-t)^{-\mu-\frac{3}{2}} K_{v+\frac{1}{2}}\left(\frac{pz^2}{t(z-t)}\right) dt \right\} dp \\ &= \frac{z^{-\mu-\frac{3}{2}} \sqrt{\frac{2pz^2}{\pi}}}{\Gamma(-\mu)} \int_0^\infty p^{r-1} \left\{ \int_0^z t^{\eta-\frac{1}{2}} \left(1-\frac{t}{z}\right)^{-\mu-\frac{3}{2}} K_{v+\frac{1}{2}}\left(\frac{pz^2}{t(z-t)}\right) dt \right\} dp \\ &= \frac{z^{\eta-\mu} \sqrt{\frac{2p}{\pi}}}{\Gamma(-\mu)} \int_0^\infty p^{r-1} \left\{ \int_0^1 u^{\eta-\frac{1}{2}} (1-u)^{-\mu-\frac{3}{2}} K_{v+\frac{1}{2}}\left(\frac{p}{u(1-u)}\right) du \right\} dp \\ &= \frac{z^{\eta-\mu} \sqrt{\frac{2}{\pi}}}{\Gamma(-\mu)} \int_0^1 u^{\eta-\frac{1}{2}} (1-u)^{-\mu-\frac{3}{2}} \left(\int_0^\infty p^{r-\frac{1}{2}} K_{v+\frac{1}{2}}\left(\frac{p}{u(1-u)}\right) dp \right) du. \end{aligned} \quad (2.16)$$

Next,

$$\begin{aligned} \int_0^\infty p^{r-\frac{1}{2}} K_{v+\frac{1}{2}}\left(\frac{p}{u(1-u)}\right) dp &= u^{r+\frac{1}{2}} (1-u)^{r+\frac{1}{2}} \int_0^\infty w^{r-\frac{1}{2}} K_{v+\frac{1}{2}}(w) dw \\ &= u^{r+\frac{1}{2}} (1-u)^{r+\frac{1}{2}} 2^{r-\frac{3}{2}} \Gamma\left(\frac{r-v}{2}\right) \Gamma\left(\frac{r+v+1}{2}\right), \end{aligned} \quad (2.17)$$

where $w = \frac{p}{u(1-u)}$, $\operatorname{Re}(r-v) > 0$, $\operatorname{Re}(r+v) > -1$ (see [2, 13]). Using (2.17) in (2.16), we obtain

$$\begin{aligned} M\{\mathfrak{D}_{z;p,v}^\mu(z^\eta); p \rightarrow r\} &= \frac{z^{\eta-\mu} 2^{r-1} \Gamma(\frac{r-v}{2}) \Gamma(\frac{r+v+1}{2})}{\sqrt{\pi} \Gamma(-\mu)} \int_0^1 u^{\eta+r} (1-u)^{r-\mu-1} du \\ &= \frac{z^{\eta-\mu} 2^{r-1} \Gamma(\frac{r-v}{2}) \Gamma(\frac{r+v+1}{2})}{\sqrt{\pi} \Gamma(-\mu)} \beta(\eta+r+1, r-\mu), \end{aligned}$$

which is (2.15). \square

Theorem 2.6 We have the Mellin transform formula

$$M \left\{ \mathfrak{D}_{z;p,v}^\mu ((1-z)^{-\alpha}); p \rightarrow r \right\} = \frac{2^{r-1} \Gamma(\frac{r-v}{2}) \Gamma(\frac{r+v+1}{2}) z^{-\mu}}{\sqrt{\pi} \Gamma(-\mu)} \beta(r+1, r-\mu) {}_2F_1(\alpha, r+1; r-\mu+1; z), \quad (2.18)$$

where $\operatorname{Re}(\alpha) > 0$, $\operatorname{Re}(\mu) < 0$, $\operatorname{Re}(r) > 0$, $\operatorname{Re}(r) > 0$, and $|z| < 1$.

Proof Using the power series for $(1-z)^{-\alpha}$ and applying Theorem 2.5 with $\eta = n$, we find

$$\begin{aligned} M \left\{ \mathfrak{D}_{z;p,v}^\mu ((1-z)^{-\alpha}); p \rightarrow r \right\} &= \sum_{n=0}^{\infty} \frac{(\alpha)_n}{n!} M \left\{ \mathfrak{D}_{z;p,v}^\mu (z^n); p \rightarrow r \right\} \\ &= \frac{2^{r-1} \Gamma(\frac{r-v}{2}) \Gamma(\frac{r+v+1}{2})}{\sqrt{\pi} \Gamma(-\mu)} \sum_{n=0}^{\infty} \frac{(\alpha)_n}{n!} \beta(n+r+1, r-\mu) z^{n-\mu} \\ &= \frac{2^{r-1} \Gamma(\frac{r-v}{2}) \Gamma(\frac{r+v+1}{2}) z^{-\mu}}{\sqrt{\pi} \Gamma(-\mu)} \sum_{n=0}^{\infty} \beta(n+r, r-\mu) \frac{(\alpha)_n z^n}{n!} \\ &= \frac{2^{r-1} \Gamma(\frac{r-v}{2}) \Gamma(\frac{r+v+1}{2}) z^{-\mu}}{\sqrt{\pi} \Gamma(-\mu)} \beta(r+1, r-\mu) {}_2F_1(\alpha, r+1; r-\mu+1; z), \end{aligned}$$

which shows (2.18). □

3. Generating relations and some further results

In this section, we derive some generating relations of linear and bilinear type for extended (p, v) -hypergeometric functions.

Theorem 3.1 We have the generating relation

$$\sum_{n=0}^{\infty} \frac{(\alpha)_n}{n!} {}_2F_{1;p,v}(\alpha+n, \beta; \gamma; z) t^n = (1-t)^{-\alpha} {}_2F_{1;p,v}\left(\alpha, \beta; \gamma; \frac{z}{1-t}\right), \quad (3.1)$$

where $|z| < \min(|1, 1-t|)$, $\operatorname{Re}(\alpha) > 0$, $\operatorname{Re}(\gamma) > \operatorname{Re}(\beta) > 0$.

Proof Consider the series identity

$$[(1-z)-t]^{-\alpha} = (1-t)^{-\alpha} \left[1 - \frac{z}{1-t} \right]^{-\alpha}.$$

Thus, the power series expansion yields

$$\sum_{n=0}^{\infty} \frac{(\alpha)_n}{n!} (1-z)^{-\alpha} \left(\frac{t}{1-z} \right)^n = (1-t)^{-\alpha} \left[1 - \frac{z}{1-t} \right]^{-\alpha}. \quad (3.2)$$

Multiplying both sides of (3.2) by $z^{\beta-1}$ and then applying the operator $\mathfrak{D}_{z;p,v}^{\beta-\gamma}$ on both sides, we get

$$\mathfrak{D}_{z;p,v}^{\beta-\gamma} \left[\sum_{n=0}^{\infty} \frac{(\alpha)_n}{n!} (1-z)^{-\alpha} \left(\frac{t}{1-z} \right)^n z^{\beta-1} \right] = (1-t)^{-\alpha} \mathfrak{D}_{z;p,v}^{\beta-\gamma} \left[z^{\beta-1} \left(1 - \frac{z}{1-t} \right)^{-\alpha} \right].$$

Interchanging the order of summation and the operator $\mathfrak{D}_{z;p,v}^{\beta-\gamma}$, we have

$$\sum_{n=0}^{\infty} \frac{(\alpha)_n}{n!} \mathfrak{D}_{z;p,v}^{\beta-\gamma} [z^{\beta-1}(1-z)^{-\alpha-n}] t^n = (1-t)^{-\alpha} \mathfrak{D}_{z;p,v}^{\beta-\gamma} \left[z^{\beta-1} \left(1 - \frac{z}{1-t} \right)^{-\alpha} \right].$$

Thus, by applying Theorem 2.3, we obtain (3.1). \square

Theorem 3.2 *We have the generating relation*

$$\sum_{n=0}^{\infty} \frac{(\beta)_n}{n!} {}_2F_{1;p,v}(\delta-n, \beta; \gamma; z) t^n = (1-t)^{-\beta} F_1 \left(\alpha, \delta, \beta; \gamma; -\frac{zt}{1-t}; p, v \right), \quad (3.3)$$

where $|t| < \frac{1}{1+|t|}$, $\operatorname{Re}(\delta) > 0$, $\operatorname{Re}(\beta) > 0$, $\operatorname{Re}(\gamma) > \operatorname{Re}(\alpha) > 0$.

Proof Consider the series identity

$$[1 - (1-z)t]^{-\beta} = (1-t)^{-\beta} \left[1 + \frac{zt}{1-t} \right]^{-\beta}.$$

Using power series expansion on the left-hand side, we have

$$\sum_{n=0}^{\infty} \frac{(\beta)_n}{n!} (1-z)^n t^n = (1-t)^{-\beta} \left[1 - \frac{zt}{1-t} \right]^{-\beta}. \quad (3.4)$$

Multiplying both sides of (3.4) by $z^{\alpha-1}(1-z)^{-\delta}$ and then applying the operator $\mathfrak{D}_{z;p,v}^{\alpha-\gamma}$ on both sides, we get

$$\mathfrak{D}_{z;p,v}^{\alpha-\gamma} \left[\sum_{n=0}^{\infty} \frac{(\beta)_n}{n!} z^{\alpha-1}(1-z)^{-\delta+n} t^n \right] = (1-t)^{-\beta} \mathfrak{D}_{z;p,v}^{\alpha-\gamma} \left[z^{\alpha-1}(1-z)^{-\delta} \left(1 - \frac{zt}{1-t} \right)^{-\beta} \right],$$

where $\operatorname{Re}(\alpha) > 0$ and $|zt| < |1-t|$. Thus, by Theorem 2.2, we obtain

$$\sum_{n=0}^{\infty} \frac{(\beta)_n}{n!} \mathfrak{D}_{z;p,v}^{\alpha-\gamma} [z^{\alpha-1}(1-z)^{-\delta+n}] t^n = (1-t)^{-\beta} \mathfrak{D}_{z;p,v}^{\alpha-\gamma} \left[z^{\alpha-1}(1-z)^{-\delta} \left(1 - \frac{zt}{1-t} \right)^{-\beta} \right].$$

Applying Theorem 2.4 on both sides, we get (3.3). \square

Theorem 3.3 *We have*

$$\mathfrak{D}_{z;p,v}^{\eta-\mu} \left[z^{\eta-1} E_{\gamma,\delta}^{\mu}(z) \right] = \frac{z^{\mu-1}}{\Gamma(\mu-\eta)} \sum_{n=0}^{\infty} \frac{(\mu)_n}{\Gamma(\gamma n + \delta)} \beta_{p,v}(\eta+n, \mu-\eta) \frac{z^n}{n!}, \quad (3.5)$$

where $\gamma, \delta, \mu \in \mathbb{C}$, $\operatorname{Re}(p) > 0$, $\operatorname{Re}(q) > 0$, $\operatorname{Re}(\mu) > \operatorname{Re}(\eta) > 0$, $\operatorname{Re}(\lambda) > 0$, $\operatorname{Re}(\rho) > 0$, and $E_{\gamma,\delta}^{\mu}(z)$ is the Mittag-Leffler function (see [14]) defined as

$$E_{\gamma,\delta}^{\mu}(z) = \sum_{n=0}^{\infty} \frac{(\mu)_n}{\Gamma(\gamma n + \delta)} \frac{z^n}{n!}. \quad (3.6)$$

Proof Using (3.6) in (3.5), we obtain

$$\mathfrak{D}_{z;p,v}^{\eta-\mu} \left[z^{\eta-1} E_{\gamma,\delta}^{\mu}(z) \right] = \mathfrak{D}_{z;p,v}^{\eta-\mu} \left[z^{\eta-1} \left\{ \sum_{n=0}^{\infty} \frac{(\mu)_n}{\Gamma(\gamma n + \delta)} \frac{z^n}{n!} \right\} \right].$$

By Theorem 2.2, we have

$$\mathfrak{D}_{z;p,v}^{\eta-\mu} \left[z^{\eta-1} E_{\gamma,\delta}^{\mu}(z) \right] = \sum_{n=0}^{\infty} \frac{(\mu)_n}{\Gamma(\gamma n + \delta)} \left\{ \mathfrak{D}_{z;p,v}^{\eta-\mu} \left[z^{\eta+n-1} \right] \right\}.$$

Applying Theorem 2.1, we get (3.5). \square

Theorem 3.4 We have

$$\begin{aligned} & \mathfrak{D}_{z;p,v}^{\eta-\mu} \left\{ z^{\eta-1} {}_m\Psi_n \left[\begin{matrix} (\alpha_i, A_i)_{1,m}; \\ (\beta_j, B_j)_{1,n}; \end{matrix} \middle| z \right] \right\} \\ &= \frac{z^{\mu-1}}{\Gamma(\mu-\eta)} \sum_{k=0}^{\infty} \frac{\prod_{i=1}^m \Gamma(\alpha_i + A_i k)}{\prod_{j=1}^n \Gamma(\beta_j + B_j k)} \beta_{p,v}(\eta+k, \mu-\eta) \frac{z^k}{k!}, \end{aligned} \quad (3.7)$$

where $\operatorname{Re}(p) > 0$, $\operatorname{Re}(q) > 0$, $\operatorname{Re}(\mu) > \operatorname{Re}(\eta) > 0$, $\operatorname{Re}(\lambda) > 0$, $\operatorname{Re}(\rho) > 0$, and ${}_m\Psi_n(z)$ denotes the Fox–Wright function (see [6, pp. 56–58] defined by

$${}_m\Psi_n(z) = {}_m\Psi_n \left[\begin{matrix} (\alpha_i, A_i)_{1,m}; \\ (\beta_j, B_j)_{1,n}; \end{matrix} \middle| z \right] = \sum_{k=0}^{\infty} \frac{\prod_{i=1}^m \Gamma(\alpha_i + A_i k)}{\prod_{j=1}^n \Gamma(\beta_j + B_j k)} \frac{z^k}{k!}. \quad (3.8)$$

Proof Applying Theorem 2.1 and following the same procedure used in the proof of Theorem 3.3, we obtain (3.7). \square

4. Concluding remarks

In this paper, we defined an extension of the extended fractional derivative operator. We conclude that when $v = 0$ and using the fact that $K_{\frac{1}{2}}(z) = \sqrt{\frac{\pi}{2z}} e^{-z}$, all results established in this paper reduce to the results related to the extended Riemann–Liouville fractional derivative operator defined in [12]. Also, when $v = 0$ and $p = 0$, we get the results related to the classical Riemann–Liouville fractional derivative operator.

References

- [1] Baleanu D, Agarwal P, Parmar RK, Alqurashi MM, Salahshour S. Extension of the fractional derivative operator of the Riemann–Liouville. J Nonlinear Sci Appl 2017; 10: 2914–2924.
- [2] Chaudhry MA, Qadir A, Rafique M, Zubair SM. Extension of Euler’s beta function. J Comput Appl Math 1997; 78: 19–32.
- [3] Chaudhry MA, Qadir A, Srivastava HM, Paris RB. Extended hypergeometric and confluent hypergeometric functions. Appl Math Comput 2004; 159: 589–602.

- [4] Choi J, Rathie AK, Parmar RK. Extension of extended beta, hypergeometric and confluent hypergeometric functions. *Honam Math J* 2014; 36: 357-385.
- [5] Dar SA, Paris RB.A (p, ν) -extension of the Apell function $F_1(\cdot)$ and its properties. arXiv 2017; 1711.07780.
- [6] Kilbas AA, Srivastava HM, Trujillo J. *Theory and Applications of Fractional Differential Equations*. Amsterdam, the Netherlands: Elsevier Science BV, 2006.
- [7] Kıymaz O, Çetinkaya A, Agarwal P. An extension of Caputo fractional derivative operator and its applications. *J Nonlinear Sci Appl* 2016; 9: 3611-3621.
- [8] Luo MJ, Milovanovic GV, Agarwal P. Some results on the extended beta and extended hypergeometric functions. *Appl Math Comput* 2014; 248: 631-651.
- [9] Mubeen S, Rahman G, Nisar KS, Choi J, Arshad M. An extended beta function and its properties. *Far East J Math Sci* 2017; 102: 1545-1557.
- [10] Olver FWJ, Lozier DW, Boisvert RF, Clark, Charles W. *NIST Handbook of Mathematical Functions*. Washington, DC, USA: National Institute of Standards and Technology, 2010.
- [11] Özarlan MA, Özergin E. Some generating relations for extended hypergeometric functions via generalized fractional derivative operator. *Math Comput Modelling* 2010; 52: 1825-1833.
- [12] Özergin E, Özarlan MA, Altın A. Extension of gamma, beta and hypergeometric functions. *J Comput Appl Math* 2011; 235: 4601-4610.
- [13] Parmar RK, Chopra P, Paris RB. On an extension of extended beta and hypergeometric functions. *J Class Anal* 2017; 11: 91-106.
- [14] Prabhakar T. A singular integral equation with a generalized Mittag Leffler function in the kernel. *Yokohama Math J* 1971; 19: 7-15.
- [15] Rainville ED. *Special Functions*. New York, NY, USA: Macmillan, 1960.
- [16] Srivastava HM, Parmar RK, Chopra P. A class of extended fractional derivative operators and associated generating relations involving hypergeometric functions. *Axioms* 2012; 1: 238-258.