



On some sums at the a -points of the k -th derivatives of the Dirichlet L -functions

Mohammed MEKKAOUI¹ , Abdallah DERBAL¹ , Kamel MAZHOUDA^{2,*} 

¹Department of Mathematics, Laboratory of Nonlinear Partial Differential Equations and History of Mathematics, École Normale Supérieure, Algiers, Algeria

²Department of Mathematics, Faculty of Science of Monastir, University of Monastir, Monastir, Tunisia

Received: 24.02.2020

Accepted/Published Online: 04.05.2020

Final Version: 21.09.2020

Abstract: Let $L^{(k)}(s, \chi)$ be the k -th derivative of the Dirichlet L -function associated with a primitive character χ mod q and a be a complex number. The solutions of $L^{(k)}(s, \chi) = a$ are called a -points. In this paper, we give an asymptotic formula for the sums

$$\sum_{\rho_{0,\chi}^{(k)}: 0 < \gamma_{0,\chi}^{(k)} < T} L^{(j)}(\rho_{0,\chi}^{(k)}, \chi) \quad \text{and} \quad \sum_{\rho_{a,\chi}^{(k)}: 1 < \gamma_{a,\chi}^{(k)} < T} L^{(j)}(\rho_{a,\chi}^{(k)}, \chi) \quad \text{as} \quad T \rightarrow \infty$$

where j and k are nonnegative integers and $\rho_{a,\chi}^{(k)}$ denotes an a -point of the k -th derivative $L^{(k)}(s, \chi)$ and $\gamma_{a,\chi}^{(k)} = \text{Im}(\rho_{a,\chi}^{(k)})$. This work continues the investigations of Kaptan, Karabulut, and Yildirim [7, 10] and Mazhouda and Onozuka [12].

Key words: Dirichlet L -function, a -points, value-distribution

1. Introduction

Let $L(s, \chi)$ be the Dirichlet L -function associated with a primitive character χ mod q and a be a complex number. The zeros of $L(s, \chi) - a$, which will be denoted by $\rho_{a,\chi} = \beta_{a,\chi} + i\gamma_{a,\chi}$ are called the a -points of $L(s, \chi)$. First, we note that there is an a -point near any trivial zero $s = -2n$ if $\chi(-1) = 1$ and $s = -2n - 1$ if $\chi(-1) = -1$ for sufficiently large n . Apart from these a -points, there are only finitely many other a -points in the half-plane $\text{Re}(s) = \sigma \leq 0$. The a -points with $\beta_{a,\chi} \leq 0$ are said to be trivial. All other a -points lie in a strip $0 < \text{Re}(s) < A$, where A is a constant depending on a ; these numbers are called the nontrivial a -points. The number of these a -points satisfies a Riemann-von Mangoldt type formula (we refer to [14, chapter 7.2] for the proof of this formula which is stated for functions in a subclass of the Selberg class including the Dirichlet L -functions $L(s, \chi)$), namely

$$N_{a,\chi}(T) = \sum_{\substack{\rho_{a,\chi} : 0 < \gamma_{a,\chi} \leq T \\ \beta_{a,\chi} > 0}} 1 = \frac{T}{2\pi} \log \left(\frac{qT}{2\pi c_a e} \right) + O(\log T), \quad (1.1)$$

*Correspondence: kamel.mazhouda@fsm.rnu.tn

2010 AMS Mathematics Subject Classification: 11M06, 11M26, 11M36

where $c_a = m$ if $a = 1$ and $c_a = 1$, otherwise, with $m = \min\{n \geq 2, \chi(n) \neq 0\}$. Here and in the sequel the error term depends on q ; however, the main term is essentially independent of a . Moreover, $N_{a,\chi}(T) \sim N_\chi(T)$ as $T \rightarrow \infty$, where $N_\chi(T) = N_{0,\chi}(T)$ denotes the number of nontrivial zeros $\rho_\chi = \beta_\chi + i\gamma_\chi$ of $L(s, \chi)$ satisfying $0 < \gamma_\chi < T$.

In [1], Conrey and Ghosh suggested the problem of estimating the average $\sum_{0 < \gamma^{(k)} < T} \zeta^{(j)}(\rho^{(k)})$ for nonnegative integers j and k , where $\rho^{(k)} = \beta^{(k)} + i\gamma^{(k)}$ denote a zero of the k -th derivative $\zeta^{(k)}(s)$. One of the first results on this topic was given by Fujii [3]. He gave an asymptotic formula of the sum $\sum_{0 < \gamma < T} \zeta'(\rho)X^\rho$ for a rational number $X > 0$. The $k = 0$ case was treated by Kaptan et al. [7]. Garunkštis and Steuding [4] gave a generalization of Fujii's asymptotic formula with $X = 1$ that if $T \rightarrow \infty$, we have

$$\sum_{\substack{\rho_a : 0 < \gamma_a \leq T \\ \beta_a > 0}} \zeta'(\rho_a) = \left(\frac{1}{2} - a\right) \frac{T}{2\pi} \log^2\left(\frac{T}{2\pi}\right) + (c_0 - 1 + 2a) \frac{T}{2\pi} \log\left(\frac{T}{2\pi}\right) + (1 - c_0 - c_0^2 + 3c_1 - 2a) \frac{T}{2\pi} + O\left(Te^{-C\sqrt{\log T}}\right), \tag{1.2}$$

where C is some positive constant and c_n are the Stieltjes constants given by the Laurent series expansion of $\zeta(s)$ at $s = 1$,

$$\zeta(s) = \frac{1}{s-1} + \sum_{n=0}^{\infty} c_n(s-1)^n \tag{1.3}$$

Recently, Mazhouda and Onozuka [12] proved that for $j, k \in \mathbb{Z}_{\geq 0}$ and large T ,

$$\sum_{1 < \gamma_a^{(k)} < T} \zeta^{(j)}(\rho_a^{(k)}) = (-1)^j (\delta_{j,0} + a\delta_{k,0} + B(j, k)) \frac{T}{2\pi} \left(\log \frac{T}{2\pi}\right)^{j+1} + O_{j,k}(T(\log T)^j), \tag{1.4}$$

where the implicit constant in the error terms may depend on a . To do so, they used the following result of Karabulut and Yildirim [10] for fixed $j, k \in \mathbb{Z}_{\geq 0}$ and large T , one has

$$\sum_{0 < \gamma^{(k)} < T} \zeta^{(j)}(\rho^{(k)}) = (-1)^j (\delta_{j,0} + B(j, k)) \frac{T}{2\pi} \left(\log \frac{T}{2\pi}\right)^{j+1} + O_{j,k}(T \log^j T), \tag{1.5}$$

where $\delta_{j,0} = 1$ if $j = 0$ and 0 otherwise,

$$B(j, k) = -\frac{k+1}{j+1} - j! \sum_{r=1}^k \frac{e^{-z_r}}{z_r^{j+1}} P_k(z_r) + j! \sum_{r=1}^k \frac{1}{z_r^{j+1}}, \tag{1.6}$$

the sum over r being void in the case $k = 0$ and z_r ($r = 1, \dots, k$) being the zeros of $P_k(z) = \sum_{j=0}^k \frac{z^j}{j!}$.

Let $\rho_{a,\chi}^{(k)} = \beta_{a,\chi}^{(k)} + i\gamma_{a,\chi}^{(k)}$ denote an a -point of $L^{(k)}(s, \chi)$. Similar to the a -points of $L(s, \chi)$, there is an a -point of $L^{(k)}(s, \chi)$ near any trivial zero $s = -2n - \left(\frac{1-\chi(-1)}{2}\right)$ for sufficiently large n and apart from these

a -points, there are only finitely many other a -points in the half-plane $\sigma \leq C$ for any $C < 0$ (see Lemma 2.1 below).

In this paper, first we give an asymptotic formula for the sum

$$\sum_{\rho_{0,\chi}^{(k)}; 0 < \gamma_{0,\chi}^{(k)} < T} L^{(j)}(\rho_{0,\chi}^{(k)}, \chi) \tag{1.7}$$

and as a consequence, we obtain an estimate for

$$\sum_{\rho_{a,\chi}^{(k)}; 1 < \gamma_{a,\chi}^{(k)} < T} L^{(j)}(\rho_{a,\chi}^{(k)}, \chi) \tag{1.8}$$

where a is a complex number. The first sum extend Karabulut and Yildirim’s result to the k -th derivative of the Dirichlet L -functions and is evaluated in the following theorem.

Theorem 1.1 *Let $k, j \in \mathbb{N}$ be fixed and χ be a primitive character modulo q . Then as $T \rightarrow \infty$, we have*

$$\sum_{\rho_{\chi}^{(k)}; 0 < \gamma_{\chi}^{(k)} \leq T} L^{(j)}(\rho_{\chi}^{(k)}, \chi) = (-1)^j (\delta_{j,0} + B(j, k)) \frac{T}{2\pi} \left(\log \frac{qT}{2\pi} \right)^{j+1} + O_{j,k} \left(T (\log qT)^j \right), \tag{1.9}$$

where $B(j, k)$ is defined by (1.6).

From Theorem 1.1, we get our main result

Theorem 1.2 *Let $k, j \in \mathbb{N}$ be fixed, a be a complex number and χ be a primitive character modulo q . Then as $T \rightarrow \infty$, we have*

$$\sum_{\rho_{a,\chi}^{(k)}; 1 < \gamma_{a,\chi}^{(k)} \leq T} L^{(j)}(\rho_{a,\chi}^{(k)}, \chi) = (-1)^j (\delta_{j,0} + a\delta_{k,0} + B(j, k)) \frac{T}{2\pi} \left(\log \frac{qT}{2\pi} \right)^{j+1} + O_{j,k} \left(T (\log qT)^j \right) \tag{1.10}$$

Here and in the sequel, the implicit constant in the error terms may depend on a .

Remark. By Theorem 1.2, we deduce the average value of $L^{(j)}(\rho_a^{(k)}, \chi)$ over the a -points $\rho_{a,\chi}^{(k)}$ of $L^{(k)}(s, \chi)$ with $1 < \text{Im}(\rho_{a,\chi}^{(k)}) < T$, i.e.

$$\frac{1}{N_{k,\chi}(a, T)} \sum_{1 < \gamma_{a,\chi}^{(k)} < T} L^{(j)}(\rho_{a,\chi}^{(k)}, \chi),$$

where $N_{k,\chi}(a, T)$ is the number of terms in the above sum. By the same argument as in [13], we have an asymptotic formula for $N_{k,\chi}(a, T)$ which is $\sim (T/2\pi) \log \frac{qT}{2\pi}$ (see [15] for the asymptotic formula of $N_{k,\chi}(0, T)$).

Hence, the average is $(-1)^j (\delta_{j,0} + a\delta_{k,0} + B(j, k)) \left(\log \frac{qT}{2\pi} \right)^j$. Thus, this tells us about the size of $L^{(j)}(s, \chi)$ at certain points (namely the a -points of $L^{(k)}(s, \chi)$).

2. Preliminary lemmas and equations

In this section, we give some lemmas and formulas useful for the proof of our Theorems. We start with well-known results on the Dirichlet L -function $L(s, \chi)$ (see Davenport book [2]) and its k -th derivative. If $\chi \pmod q$ is a primitive character, then

$$L(s, \chi) = \Lambda(s, \chi)L(1 - s, \bar{\chi}), \tag{2.1}$$

where

$$\Lambda(s, \chi) = \frac{2\tau(\chi)}{i^\kappa q} \left(\frac{2\pi}{q}\right)^{s-1} \Gamma(1 - s) \sin\left(\frac{\pi}{2}(s + \kappa)\right), \tag{2.2}$$

with $\tau(\chi) = \sum_{r=1}^q \chi(r)e^{\frac{2\pi ir}{q}}$ and $\kappa = \frac{1}{2}(1 - \chi(-1))$. From (2.2) and by Stirling's formula (see[9, page 13]), we get

$$\begin{aligned} & \Lambda(1 - s, \chi) \\ &= \frac{\tau(\chi)}{i^\kappa \sqrt{q}} \exp\left\{it \log\left(\frac{q|t|}{2\pi e}\right) - \operatorname{sgn}(t)\left(\frac{i\pi}{2}\right)\left(\frac{1}{2} - \kappa\right)\right\} \left(\frac{q|t|}{2\pi}\right)^{\sigma-\frac{1}{2}} \left(1 + O\left(\frac{1}{|t|}\right)\right) \end{aligned} \tag{2.3}$$

in any fixed halfstrip $\alpha \leq \sigma \leq \beta, |t| \geq 1$. Moreover, for any fixed $\sigma, j \geq 0$ and $|t| \geq 1$, we have

$$\frac{\Lambda'}{\Lambda}(s, \chi) = -\log\frac{q|t|}{2\pi} + O\left(\frac{1}{|t|}\right), \left(\frac{d}{ds}\right)^j \frac{\Lambda'}{\Lambda}(s, \chi) \ll |t|^{-j} \tag{2.4}$$

and

$$\Lambda^{(j)}(1 - s, \chi) = \Lambda(1 - s, \chi) \left(-\log\frac{q|t|}{2\pi}\right)^j + O\left(q^{\sigma-\frac{1}{2}}|t|^{\sigma-\frac{3}{2}}(\log q|t|)^{j-1}\right). \tag{2.5}$$

Using equations (2.3)–(2.5) with upon j -fold differentiation of the functional equation (2.1), we obtain

$$L^{(j)}(1 - s, \chi) = (-1)^j \Lambda(1 - s, \chi) \left(1 + O\left(\frac{1}{t}\right)\right) \sum_{m=0}^j \binom{j}{m} \ell^{j-m} L^{(m)}(s, \bar{\chi}), \tag{2.6}$$

where σ is fixed, $|t| \geq 1$ and $\ell = \log\left(\frac{q|t|}{2\pi}\right)$. Furthermore, for any fixed $\sigma, k \in \mathbb{Z}_{\geq 0}$ and $t \geq 1$, we have

$$\begin{aligned} \frac{L^{(k+1)}}{L^{(k)}}(1 - s, \chi) &= -\left(1 + O\left(\frac{1}{t}\right)\right) \left(\ell + \frac{\sum_{v=0}^k \binom{k}{v} \ell^{k-v} \frac{L^{(v+1)}}{L}(s, \bar{\chi})}{\sum_{w=0}^k \binom{k}{w} \ell^{k-w} \frac{L^{(w)}}{L}(s, \bar{\chi})}\right) \\ &= -\left(1 + O\left(\frac{1}{t}\right)\right) \left(\ell + \frac{\sum_{v=0}^k \binom{k}{v} \frac{1}{\ell^v} \frac{L^{(v+1)}}{L}(s, \bar{\chi})}{1 + \sum_{w=1}^k \binom{k}{w} \frac{1}{\ell^w} \frac{L^{(w)}}{L}(s, \bar{\chi})}\right) \\ &= -\left(1 + O\left(\frac{1}{t}\right)\right) \left(\ell + \frac{G'_k}{G_k}(s, \ell, \bar{\chi})\right), \end{aligned} \tag{2.7}$$

with the differentiation in G' is respect to s . Since $\frac{L^{(w)}}{L}(s, \bar{\chi}) \ll_w 1$ when $\sigma \geq 1 + \delta$, for sufficiently large t , we get

$$\sum_{w=1}^k \binom{k}{w} \frac{1}{\ell^w} \frac{L^{(w)}}{L}(s, \bar{\chi}) \ll_k \frac{1}{\log qt}. \tag{2.8}$$

By expanding the denominator of (2.7) as a power series, we obtain

$$\begin{aligned} \left(1 + \sum_{w=1}^k \binom{k}{w} \frac{1}{\ell^w} \frac{L^{(w)}}{L}(s, \bar{\chi})\right)^{-1} &= \sum_{u=0}^{\infty} (-1)^u \left(\sum_{w=1}^k \binom{k}{w} \frac{1}{\ell^w} \frac{L^{(w)}}{L}(s, \bar{\chi})\right)^u \\ &= \sum_{u \leq \frac{\log A}{\log \log A}}^{\infty} (-1)^u \left(\sum_{w=1}^k \binom{k}{w} \frac{1}{\ell^w} \frac{L^{(w)}}{L}(s, \bar{\chi})\right)^u + O\left(\frac{1}{A}\right) \end{aligned} \tag{2.9}$$

where $\sigma \geq 1 + \delta$ and $t \geq A$ for large A . By the functional equation (2.1) and the Phragmén–Lindelöf principle, we deduce that

$$L(s, \chi) \ll_{\epsilon} \begin{cases} |qt|^{\frac{1}{2}-\sigma+\epsilon} & \sigma < 0, \\ |qt|^{\frac{1}{2}(1-\sigma)+\epsilon} & 0 \leq \sigma \leq 1, \\ |qt|^{\epsilon} & \sigma > 1, \end{cases} \tag{2.10}$$

as $|t| \rightarrow \infty$ and where ϵ is an arbitrarily small positive number. Moreover, by Cauchy’s integral formula, we get

$$L^{(k)}(s, \chi) = \frac{k!}{2\pi i} \int_{\mathbf{C}} \frac{L(w, \chi)}{(w - s)^{k+1}} ds,$$

where \mathbf{C} is any arbitrarily small circle centered at s . Using the last bound of $L(s, \chi)$, it follows that

$$L^{(k)}(s, \chi) \ll_{\epsilon} \begin{cases} |qt|^{\frac{1}{2}-\sigma+\epsilon} & \sigma < 0, \\ |qt|^{\frac{1}{2}(1-\sigma)+\epsilon} & 0 \leq \sigma \leq 1, \\ |qt|^{\epsilon} & \sigma > 1. \end{cases} \tag{2.11}$$

Now, using the same argument as in [13, Lemma 2.6], we get easily

$$\frac{L^{(k+1)}(s, \chi)}{L^{(k)}(s, \chi) - a} = \sum_{|\gamma_{a, \chi}^{(k)} - t| < 1} \frac{1}{s - \rho_{a, \chi}^{(k)}} + O(\log qt), \tag{2.12}$$

for any constants α, β and $s \in \mathbb{C}$ with $\alpha \leq \sigma \leq \beta$ and large t .

Lemma 2.1 *Let k be a positive integer, χ be a primitive character modulo q and $a \in \mathbb{C}$. Then, there exist real numbers $E_1 = E_1(k, a, q) \leq 0$ and $E_2 = E_2(k, a, q) \geq 1$ such that there is no a -point of $L^{(k)}(s, \chi)$ for $\{s \in \mathbb{C}, \sigma \leq E_1, |t| \geq 1\}$ and $\{s \in \mathbb{C}, \sigma \geq E_2\}$.*

Proof The case $a = 0$ was treated by Yildirim in [16]. Hence, we consider only the case $a \neq 0$. From equation (2.1) and by differentiating k times, we obtain

$$\begin{aligned} L^{(k)}(1-s, \chi) &= (-1)^k \frac{2\tau(\chi)}{i^\kappa q} \left(\frac{2\pi}{q}\right)^{-s} \sum_{j=0}^k \Gamma^{(j)}(s) R_{j,k}(s) \\ &= (-1)^k \frac{2\tau(\chi)}{i^\kappa q} \left(\frac{2\pi}{q}\right)^{-s} \left\{ \Gamma^{(k)}(s) \cos\left(\frac{\pi}{2}(s-\kappa)\right) L(s, \bar{\chi}) + \sum_{j=0}^{k-1} \Gamma^{(j)}(s) R_{j,k}(s) \right\}, \end{aligned} \tag{2.13}$$

where

$$R_{j,k}(s) = P_{j,k}(s) \cos\left(\frac{\pi}{2}(s-\kappa)\right) + Q_{j,k}(s) \sin\left(\frac{\pi}{2}(s-\kappa)\right), \tag{2.14}$$

$$P_{j,k}(s) = \sum_{n=0}^k a_{j,k,n} L^{(n)}(s, \bar{\chi}) \tag{2.15}$$

and

$$Q_{j,k}(s) = \sum_{n=0}^k b_{j,k,n} L^{(n)}(s, \bar{\chi}), \tag{2.16}$$

where $a_{j,k,n}$ and $b_{j,k,n}$ are constants that may depend on q . Using [16, Equation(13)], derivatives of the Gamma function can be estimated as follows:

$$\Gamma^{(j)}(s) = \Gamma(s) (\log s)^j \left(1 + O\left(\frac{1}{s \log s}\right)\right) \tag{2.17}$$

in the region $\{s \in \mathbb{C}, \sigma \geq 1 + \delta, |t| \geq 1\}$. Using the last estimate and the fact that in the same region $L(s, \chi) \asymp 1$ and $L^{(j)}(s, \chi) = \sum_{n \geq 2} \frac{\chi(n)(-\log n)^j}{n^s} \ll 1$, we get

$$\left| \Gamma^{(k)}(s) \cos\left(\frac{\pi}{2}(s-\kappa)\right) L(s, \bar{\chi}) \right| \asymp \left| \Gamma(s) \log^k(s) e^{\pi \frac{|t|}{2}} \right| \tag{2.18}$$

and

$$\left| \sum_{j=0}^{k-1} \Gamma^{(j)}(s) R_{j,k}(s) \right| \ll \left| \Gamma(s) \log^{k-1}(s) e^{\pi \frac{|t|}{2}} \right|. \tag{2.19}$$

As a consequence, one has

$$\begin{aligned} &L^{(k)}(1-s, \chi) \\ &= (-1)^k \frac{2\tau(\chi)}{i^\kappa q} \left(\frac{2\pi}{q}\right)^{-s} \Gamma(s) \log^k(s) \cos\left(\frac{\pi}{2}(s-\kappa)\right) L(s, \bar{\chi}) \left(1 + O\left(\frac{1}{|\log s|}\right)\right) \end{aligned} \tag{2.20}$$

in the region $\{s \in \mathbb{C}, \sigma \geq 1 + \delta, |t| \geq 1\}$. It follows from (2.20) that $L^k(1-s, \chi) \rightarrow \infty$ as $\sigma \rightarrow \infty$. Thus, there exists $E_1 = E_1(k, a, q) \leq 0$ such that $|L^{(k)}(s, \chi)| > |a|$ for $\sigma \leq E_1$ and $|t| \geq 1$. Next, since

$L^{(k)}(s, \chi) = \sum_{n \geq 2} \frac{\chi(n)(-\log n)^k}{n^s} \rightarrow 0$ as $\sigma \rightarrow \infty$ there exists $E_2 = E_2(k, a, q) \geq 1$ such that $|L^{(k)}(s, \chi)| < |a|$.

□

Remark. It can also be seen by Rouché’s theorem that there is $N_k = N_k(a, q) < 0$ such that $L^{(k)}(s, \chi) = a$ has only one zero in the region $\{s \in \mathbb{C}, -1 - 2n - \kappa < \sigma < 1 - 2n - \kappa, -1 < t < 1\}$ for $-n < N_k$. Moreover, apart from these a -points, there are only finitely many other a -points in the half-plane $\sigma \leq C$ for any $C < 0$.

From Lemma 2.1, equation (2.11) and by Jensen’s formula, we deduce easily the following lemma.

Lemma 2.2 *For any complex number a and any sufficiently large T , we have*

$$N_{k,\chi}(a; 1, T + 1) - N_{k,\chi}(a; 1, T) \ll \log(qT), \tag{2.21}$$

where

$$N_{k,\chi}(a; 1, T) = \sum_{\rho_{a,\chi}^{(k)}: 1 < \gamma_{a,\chi}^{(k)} < T} 1. \tag{2.22}$$

3. Proof of Theorem 1.1

To prove Theorem 1.1, we use the same argument as in [10]. For this purpose, we need to extend some lemmas for k -th derivative of Dirichlet L -function $L^k(s, \chi)$. The case $k = 0$ was already proved by Kaptan et al. [6], so here we assume $k \neq 0$.

Lemma 3.1 *Let $(b_n)_n$ be a sequence of complex numbers such that $b_n \ll n^\epsilon$ for any $\epsilon > 0$. Let $a > 1$ and m be an integer. Then, for $1 \leq T_1 \leq T$ and $|m| = O(T)$ as $T \rightarrow \infty$, one has*

$$\begin{aligned} & \frac{1}{2\pi} \int_{T_1}^T \Lambda(1 - a - it, \chi) \left(\log \left(\frac{qt}{2\pi} \right) \right)^m \sum_{n=1}^{\infty} \frac{b_n}{n^{a+it}} dt \\ &= \frac{\tau(\chi)}{q} \sum_{1 \leq n \leq \frac{qT}{2\pi}} b_n e^{-\frac{2\pi in}{q}} (\log n)^m + O\left((qT)^{a-\frac{1}{2}} (\log qT)^m\right) + O\left(q^{2a-1} (\log q)^m\right). \end{aligned}$$

Proof The case m nonnegative is treated by Kaptan in [8, Lemma 2.14] which is based on [5, Lemma 2] (see also [10, Lemma 2.2]). For the case when m is negative, we use the same argument of Kaptan and [11, Lemma 3.5] to obtain the result. □

An elementary computation yields the following lemma.

Lemma 3.2 *For $k, i_1, i_2, \dots, i_k, m \in \mathbb{N}, v \in \{0, 1, \dots, k\}, \sigma > 1$ and χ be a Dirichlet character modulo q , let us define*

$$\sum_{n=1}^{\infty} \frac{c_n(i_1, i_2, \dots, i_k; v; m; \chi)}{n^s} := \frac{L^{(v+1)}(s, \chi)L^{(m)}(s, \chi)}{L} \prod_{w=1}^k \left(\frac{L^{(w)}(s, \chi)}{L} \right)^{i_w}.$$

We have

$$\chi'(n)c_n(i_1, i_2, \dots, i_k; v; m; \chi) = c_n(i_1, i_2, \dots, i_k; v; m; \chi' \chi),$$

for every Dirichlet character χ' modulo q , with

$$|c_n(i_1, i_2, \dots, i_k; v; m; \chi)| \leq (\log n)^{K+m+1},$$

where

$$K := i_1 + 2i_2 + \dots + ki_k + v.$$

Lemma 3.3 Let χ be a Dirichlet character modulo q . Let $k, i_1, i_2, \dots, i_k, m \in \mathbb{N}, v \in \{0, 1, \dots, k\}$. For fixed k , if $i_1 + i_2 + \dots + i_k \leq \frac{\log x}{\log \log x}$, then as $T \rightarrow \infty$, we have

$$\sum_{n \leq x} c_n(i_1, i_2, \dots, i_k; v; m; \chi) = O_k(x(\log x)^{K+m}),$$

if χ is nonprincipal and

$$\sum_{n \leq x} c_n(i_1, i_2, \dots, i_k; v; m; \chi) = \frac{\varphi(q)}{q} S(i_1, i_2, \dots, i_k; v; m) x(\log x)^{K+m+1} + O_k(x(\log x)^{K+m})$$

if χ is the principal character, where

$$S(i_1, i_2, \dots, i_k; v; m) = \frac{(-1)^{K+m+1} (v+1)! m! \prod_{w=1}^k (w!)^{i_w}}{(K+m+1)!}.$$

Proof Let χ be a nonprincipal character modulo q . Lemma 3.2 with Perron's formula [15, chapter 3.12], yields

$$\begin{aligned} \sum_{n \leq x} c_n(i_1, i_2, \dots, i_k; v; m; \chi) &= \int_{1+\frac{1}{\log x}-iU}^{1+\frac{1}{\log x}+iU} \frac{L^{(v+1)}(s, \chi) L^{(m)}(s, \chi)}{L} \prod_{w=1}^k \left(\frac{L^{(w)}(s, \chi)}{L} \right)^{i_w} \frac{x^s}{s} ds \\ &+ O_k\left(\frac{x}{U} (\log x)^{K+m+2}\right), \end{aligned}$$

where $q \leq U \leq x$. Now, let C be the rectangle with vertices $1 + \frac{1}{\log x} - iU$, $1 + \frac{1}{\log x} + iU$, $\sigma_0 + iU$, and $\sigma_0 - iU$.

Case 1. Assume that $L(s, \chi)$ has no exceptional zero. We take $\sigma_0 = 1 - \frac{c}{5 \log qU}$, where c is the constant such that $L(s, \chi) \neq 0$ for $\sigma > 1 - \frac{c}{\log qU}$ (see [2, page 93]). Thus, the integrand is analytic on and inside C and we have the bound $\frac{L^{(w)}(s, \chi)}{L} \ll (\log qU)^w$. Then, we have by Cauchy's formula

$$M = \frac{1}{2\pi i} \int_C \frac{L^{(v+1)}(s, \chi) L^{(m)}(s, \chi)}{L} \prod_{w=1}^k \left(\frac{L^{(w)}(s, \chi)}{L} \right)^{i_w} \frac{x^s}{s} ds = 0.$$

Now, using that $L^{(m)}(s, \chi) \ll (qU)^{\frac{1}{2}(1-\sigma)+\epsilon}$, we get

$$\begin{aligned} \int_{1+\frac{1}{\log x}+iU}^{1-\frac{1}{5 \log qU}+iU} \frac{L^{(v+1)}(s, \chi) L^{(m)}(s, \chi)}{L} \prod_{w=1}^k \left(\frac{L^{(w)}(s, \chi)}{L} \right)^{i_w} \frac{x^s}{s} ds &\ll (\log qU)^{K+1} \int_{1+\frac{1}{\log x}}^{1-\frac{1}{5 \log qU}} (qU)^{\frac{1}{2}(1-\sigma)+\epsilon} \frac{x^\sigma}{|\sigma + iU|} d\sigma \\ &\ll x(\log qU)^{K+1} (qU)^{\frac{c}{10 \log qU}} \left(\frac{1}{\log x} + \frac{c}{5 \log qU} \right) \\ &\ll \frac{x}{U^{1-\epsilon}} (\log qU)^K. \end{aligned}$$

Analogously, we have

$$\int_{1-\frac{1}{5\log qU}-iU}^{1+\frac{1}{\log x}-iU} \frac{L^{(v+1)}}{L}(s, \chi)L^{(m)}(s, \chi) \prod_{w=1}^k \left(\frac{L^{(w)}}{L}(s, \chi)\right)^{i_w} \frac{x^s}{s} ds \ll \frac{x}{U^{1-\epsilon}} (\log qU)^K$$

and

$$\begin{aligned} \int_{1-\frac{1}{5\log qU}-iU}^{1-\frac{1}{5\log qU}-iU} \frac{L^{(v+1)}}{L}(s, \chi)L^{(m)}(s, \chi) \prod_{w=1}^k \left(\frac{L^{(w)}}{L}(s, \chi)\right)^{i_w} \frac{x^s}{s} ds &\ll (\log qU)^{K+1} \int_{-U}^U (|qt|)^{\frac{1}{2}(1-\sigma_0)+\epsilon} \frac{x^{\sigma_0}}{|\sigma_0+it|} dt \\ &\ll x(\log qU)^{K+1} (qU)^{\frac{c}{10\log qU}} x^{\sigma_0} \int_{-U}^U \frac{1}{|\sigma_0+it|} dt \\ &\ll xU^\epsilon (\log qU)^{K+1} \exp\left(\frac{-c \log x}{10 \log qU}\right). \end{aligned}$$

Let $U = (\log x)^2$. Then, from all above estimates, we obtain

$$\sum_{n \leq x} c_n(i_1, i_2, \dots, i_k; v; m; \chi) = O_k(x(\log x)^{K+m})$$

Case 2. Suppose that there is an exceptional zero β , with $\beta \geq 1 - \frac{c}{4\log qU}$. Therefore, we take $\sigma_0 = 1 - \frac{c}{3\log qU}$. Thus, the integrand has a pole at β of order $L + 1$, where $L = i_1 + i_2 + \dots + i_k$. Hence,

$$\begin{aligned} M &= \frac{1}{L!} \frac{d^L}{ds^L} \left\{ (s - \beta)^{L+1} \frac{L^{(v+1)}}{L}(s, \chi)L^{(m)}(s, \chi) \prod_{w=1}^k \left(\frac{L^{(w)}}{L}(s, \chi)\right)^{i_w} \frac{x^s}{s} \right\}_{s=\beta} \\ &= \frac{1}{L!} \sum_{j_1+j_2+j_3=L} \binom{L}{j_1, j_2, j_3} \frac{d^{j_1}}{ds^{j_1}} \left\{ (s - \beta)^{L+1} \frac{L^{(v+1)}}{L}(s, \chi)L^{(m)}(s, \chi) \prod_{w=1}^k \left(\frac{L^{(w)}}{L}(s, \chi)\right)^{i_w} \right\}_{s=\beta} \\ &\quad \times \frac{d^{j_2}}{ds^{j_2}} \{x^s\}_{s=\beta} \frac{d^{j_3}}{ds^{j_3}} \left\{ \frac{1}{s} \right\}_{s=\beta} \\ &= (-1)^L \frac{x^\beta}{\beta^{L+1}} \sum_{j_1=0}^L \frac{(-1)^{j_1}}{j_1!} \beta^{j_1} \frac{d^{j_1}}{ds^{j_1}} \left\{ (s - \beta)^{L+1} \frac{L^{(v+1)}}{L}(s, \chi)L^{(m)}(s, \chi) \prod_{w=1}^k \left(\frac{L^{(w)}}{L}(s, \chi)\right)^{i_w} \right\}_{s=\beta} \\ &\quad \times \sum_{j_2=0}^{L-j_1} \frac{(-1)^{j_2}}{j_2!} \beta^{j_2} (\log x)^{j_2}. \end{aligned}$$

By Cauchy’s formula on a disk of radius 1 centered at $s = \beta$, we deduce

$$\begin{aligned} \left| \frac{d^{j_1}}{ds^{j_1}} \left\{ (s - \beta)^{L+1} \frac{L^{(v+1)}}{L}(s, \chi)L^{(m)}(s, \chi) \prod_{w=1}^k \left(\frac{L^{(w)}}{L}(s, \chi)\right)^{i_w} \right\} \right| &\leq j_1! \max_{|s-\beta|=1} \left| \frac{L^{(v+1)}}{L}(s, \chi)L^{(m)}(s, \chi) \prod_{w=1}^k \left(\frac{L^{(w)}}{L}(s, \chi)\right)^{i_w} \right| \\ &\ll_k j_1!. \end{aligned}$$

The last equation yields to

$$\begin{aligned} M &\ll_k \frac{x^\beta}{\beta^{L+1}} \sum_{j_1=0}^L \beta^{j_1} \sum_{j_2=0}^{L-j_1} \frac{\beta^{j_2}}{j_2!} (\log x)^{j_2} \\ &\ll_k \frac{x^\beta}{\beta} (\log x)^L \\ &\ll_k x (\log x)^L. \end{aligned}$$

As above, we obtain

$$\sum_{n \leq x} c_n(i_1, i_2, \dots, i_k; v; m; \chi) = O_k(x(\log x)^{K+m})$$

Case 3. Suppose the existence of an exceptional zero β , with $\beta < 1 - \frac{c}{4 \log qU}$. Therefore, proceeding similarly as in case 1, we get

$$\sum_{n \leq x} c_n(i_1, i_2, \dots, i_k; v; m; \chi) = O_k(x(\log x)^{K+m}).$$

The proof of Lemma 3.3 when χ is principal is closely similar to that in [10, Lemma 2.4]. □

Lemma 3.4 *Let χ be a Dirichlet character modulo q . Let $k, i_1, i_2, \dots, i_k, m \in \mathbb{N}$ and $v \in \{0, 1, \dots, k\}$. For fixed k , if $i_1 + i_2 + \dots + i_k \leq \frac{\log x}{\log \log x}$, then as $T \rightarrow \infty$, we have*

$$\sum_{n \leq x} \frac{c_n(i_1, i_2, \dots, i_k; v; m; \chi)}{(\log n)^{K-r}} = O_{k,r,m}(x(\log x)^{r+m})$$

if χ is nonprincipal and

$$\sum_{n \leq x} \frac{c_n(i_1, i_2, \dots, i_k; v; m; \chi)}{(\log n)^{K-r}} = \frac{\varphi(q)}{q} S(i_1, i_2, \dots, i_k; v; m) x(\log x)^{r+m+1} + O_{k,r,m}(x(\log x)^{r+m})$$

if χ is a principal character.

Proof of Theorem 1.1. The basic idea of the proof is to interpret the sum of $L^{(j)}(\rho_\chi^{(k)}, \chi)$ as a sum of residues. By Cauchy's theorem, we have

$$\sum_{\substack{0 < \gamma_\chi^{(k)} < T \\ -b < \beta_\chi^{(k)} < a}} L^{(j)}(\rho_\chi^{(k)}, \chi) = \frac{1}{2\pi i} \int_R L^{(j)}(s, \chi) \frac{L^{(k+1)}}{L^{(k)}}(s, \chi) ds$$

where the integration is taken over a rectangular contour in counterclockwise direction denoted by R with vertices $-b + ic, a + ic, a + iT, -b + iT$ with some constants $a, b, c > 0$ such that $\frac{1}{L^{(k)}(a + it, \chi)} \ll_k 1$,

$0 < b < \frac{1}{8}$ and $L^{(k)}(s, \chi)$ has no zero on the lines $t = T$ and $t = c$. From [16, Theorem 3], we deduce that there are finitely many zeros of $L^{(k)}(s, \chi)$ in the region $\sigma < -b$ and $t > c$, then we have

$$\begin{aligned} \sum_{0 < \gamma_\chi^{(k)} < T} L^{(j)}(\rho^{(k)}, \chi) &= \frac{1}{2\pi i} \int_R L^{(j)}(s, \chi) \frac{L^{(k+1)}}{L^{(k)}}(s, \chi) ds + O(1) \\ &= \frac{1}{2\pi i} \left\{ \int_{-b+ic}^{a+ic} + \int_{a+ic}^{a+iT} + \int_{a+iT}^{-b+iT} + \int_{-b+iT}^{-b+ic} \right\} L^{(j)}(s, \chi) \frac{L^{(k+1)}}{L^{(k)}}(s, \chi) ds + O(1) \\ &= I_1 + I_2 + I_3 + I_4 + O(1) \end{aligned}$$

The first integral I_1 is independent of T , so $I_1 = O(1)$. Next, we consider I_2 , using that $\frac{1}{L^{(k)}(a + it, \chi)} \ll_k 1$ and $L^{(j)}(s, \chi) \ll 1$, we get $I_2 = O(T)$. Now, using equation (2.12) and take the horizontal sides of the rectangular contour to be a distance $\gg \frac{1}{\log qT}$ from any zero of $L^{(k)}(s, \chi)$, one has

$$\begin{aligned} I_3 &= \frac{1}{2\pi i} \int_{a+iT}^{b+iT} \sum_{\substack{\gamma_\chi^{(k)} \\ |\gamma_\chi^{(k)} - t| < 1}} \frac{L^{(j)}(s, \chi)}{s - \rho_\chi^{(k)}} ds + O\left(\int_{a+iT}^{b+iT} \log(qt) L^{(j)}(s, \chi) ds\right) \\ &= O\left((qT)^{\frac{1}{2}+b+\epsilon} \log qT \sum_{\substack{\gamma_\chi^{(k)} \\ |\gamma_\chi^{(k)} - T| < 1}} 1\right) + O\left((qT)^{\frac{1}{2}+b+\epsilon} \log qT\right). \end{aligned}$$

By Lemma 2.2, we obtain

$$I_3 = O\left((q)T^{\frac{1}{2}+b+\epsilon}(\log qT)^2\right).$$

This leads $I_3 \ll T$, since $0 < b < \frac{1}{8}$. For the fourth integral I_4 , by using equations (2.6), (2.7), and (2.9), we obtain

$$\begin{aligned} \bar{I}_4 &= -\frac{1}{2\pi i} \int_{1+b+ic}^{1+b+iT} L^{(j)}(1-s, \bar{\chi}) \frac{L^{(k+1)}}{L^{(k)}}(1-s, \bar{\chi}) ds \\ &= \frac{(-1)^j}{2\pi i} \sum_{m=0}^j \binom{j}{m} \int_{1+b+ic}^{1+b+iT} \Lambda(1-s, \bar{\chi}) \ell^{j-m+1} L^{(m)}(s, \chi) ds \\ &\quad + \frac{(-1)^j}{2\pi i} \sum_{m=0}^j \binom{j}{m} \int_{1+b+ic}^{1+b+iT} \Lambda(1-s, \bar{\chi}) \ell^{j-m} \frac{G'_k}{G_k}(s, \ell, \chi) L^{(m)}(s, \chi) ds + O(T) \\ &= S_1 + S_2 + O(T). \end{aligned}$$

Lemma 3.1 gives

$$\begin{aligned}
 S_1 &= (-1)^j \sum_{m=0}^j \binom{j}{m} \frac{\tau(\bar{\chi})}{q} \sum_{1 \leq n \leq \frac{qT}{2\pi}} (-1)^m \chi(n) e^{-\frac{2\pi i n}{q}} (\log n)^{j+1} + O\left(T^{b+\frac{1}{2}} (\log qT)^{j+1}\right) \\
 &= (-1)^j \frac{\tau(\bar{\chi})}{q} \sum_{1 \leq n \leq \frac{qT}{2\pi}} \chi(n) e^{-\frac{2\pi i n}{q}} (\log n)^{j+1} \sum_{m=0}^j \binom{j}{m} (-1)^m + O\left(T^{b+\frac{1}{2}} (\log qT)^{j+1}\right) \\
 &= \begin{cases} O\left(T^{b+\frac{1}{2}} (\log qT)^{j+1}\right) & \text{if } j \geq 1, \\ \frac{\tau(\bar{\chi})}{q} \sum_{1 \leq n \leq \frac{qT}{2\pi}} \chi(n) e^{-\frac{2\pi i n}{q}} \log n + O\left(T^{b+\frac{1}{2}} \log qT\right) & \text{if } j = 0. \end{cases}
 \end{aligned}$$

Recall that (see [2, page 146])

$$e^{-\frac{2\pi i n}{q}} = \frac{1}{\varphi(q)} \sum_{\chi' \equiv q} \tau(\bar{\chi}') \chi'(-n),$$

when $(n, q) = 1$. The last formula yields

$$\begin{aligned}
 \frac{\tau(\bar{\chi})}{q} \sum_{1 \leq n \leq \frac{qT}{2\pi}} \chi(n) e^{-\frac{2\pi i n}{q}} \log n &= \frac{\tau(\bar{\chi})}{q\varphi(q)} \sum_{\chi' \equiv q} \tau(\bar{\chi}') \chi'(-1) \sum_{1 \leq n \leq \frac{qT}{2\pi}} \chi(n) \chi'(n) \log n \\
 &= \sum_{\chi' \neq \bar{\chi}} \frac{\tau(\bar{\chi}) \tau(\bar{\chi}') \chi'(-1)}{q\varphi(q)} \sum_{1 \leq n \leq \frac{qT}{2\pi}} \chi(n) \chi'(n) \log n \\
 &\quad + \frac{\tau(\bar{\chi}) \tau(\chi) \overline{\chi(-1)}}{q\varphi(q)} \sum_{1 \leq n \leq \frac{qT}{2\pi}} \chi_0(n) \log n.
 \end{aligned}$$

Using the following estimate

$$\sum_{1 \leq n \leq x} \chi_0(n) \log n = \frac{\varphi(q)}{q} x \log(x) + O\left(\frac{\varphi(q)}{q} x\right) + O(q^\epsilon \log(x))$$

and Pólya-Vinogradov inequality

$$\sum_{n \leq x} \chi(n) \ll 2\sqrt{q} \log q$$

for every nonprincipal character modulo q , we obtain

$$S_1 = \begin{cases} O\left(T^{b+\frac{1}{2}} (\log qT)^{j+1}\right) & \text{if } j \geq 1, \\ \frac{T}{2\pi} \log\left(\frac{qT}{2\pi}\right) + O\left(T^{b+\frac{1}{2}} \log qT\right) & \text{if } j = 0. \end{cases}$$

Now, we estimate S_2 . We have

$$\begin{aligned} S_2 &= \frac{(-1)^j}{2\pi i} \sum_{m=0}^j \binom{j}{m} \int_{1+b+ic}^{1+b+iT} \Lambda(1-s, \bar{\chi}) \ell^{j-m} \frac{G'_k}{G_k}(s, \ell, \chi) L^{(m)}(s, \chi) ds + O(T) \\ &= (-1)^j \sum_{m=0}^j \binom{j}{m} \sum_{u \leq \frac{\log T}{\log \log T}} \sum_{v=0}^k (-1)^u \binom{k}{v} \sum_{i_1+i_2+\dots+i_k=u} \binom{u}{i_1, i_2, \dots, i_k} \prod_{w=1}^k \binom{k}{w}^{i_w} \\ &\quad \times \frac{1}{2\pi} \int_c^T \Lambda(-b-it, \bar{\chi}) \ell^{j-K-m} L^{(m)}(1+b+it, \chi) \frac{L^{(v+1)}}{L}(1+b+it, \chi) \prod_{w=1}^k \left(\frac{L^{(w)}}{L}(1+b+it, \chi) \right)^{i_w} dt \\ &\quad + O_{j,k} \left(T^{\frac{1}{2}+b+\epsilon} \right). \end{aligned}$$

From Lemma 3.1, we get

$$\begin{aligned} S_2 &= (-1)^j \sum_{m=0}^j \binom{j}{m} \sum_{u \leq \frac{\log T}{\log \log T}} \sum_{v=0}^k (-1)^u \binom{k}{v} \sum_{i_1+i_2+\dots+i_k=u} \binom{u}{i_1, i_2, \dots, i_k} \prod_{w=1}^k \binom{k}{w}^{i_w} \\ &\quad \times \frac{\tau(\bar{\chi})}{q} \sum_{1 \leq n \leq \frac{qT}{2\pi}} C_n(i_1, i_2, \dots, i_k; v; m; \chi) e^{-\frac{2\pi i n}{q}} (\log n)^{j-K-m} + O_{j,k} \left(T^{\frac{1}{2}+b+\epsilon} \right). \end{aligned}$$

Since

$$e^{-\frac{2\pi i n}{q}} = \frac{1}{\varphi(q)} \sum_{\chi' \equiv q} \tau(\bar{\chi}') \chi'(-n)$$

when $(n, q) = 1$, we obtain

$$\begin{aligned} S_2 &= (-1)^j \sum_{m=0}^j \binom{j}{m} \sum_{u \leq \frac{\log T}{\log \log T}} \sum_{v=0}^k (-1)^u \binom{k}{v} \sum_{i_1+i_2+\dots+i_k=u} \binom{u}{i_1, i_2, \dots, i_k} \prod_{w=1}^k \binom{k}{w}^{i_w} \\ &\quad \times \left\{ \sum_{\chi' \neq \bar{\chi}} \frac{\tau(\bar{\chi})\tau(\chi')\chi'(-1)}{q\varphi(q)} \sum_{1 \leq n \leq \frac{qT}{2\pi}} \frac{\chi'(n)c_n(i_1, i_2, \dots, i_k; v; m; \chi)}{(\log n)^{K+m-j}} + \frac{\tau(\bar{\chi})\tau(\chi)\bar{\chi}(-1)}{q\varphi(q)} \sum_{1 \leq n \leq \frac{qT}{2\pi}} \frac{\bar{\chi}(n)c_n(i_1, i_2, \dots, i_k; v; m; \chi)}{(\log n)^{K+m-j}} \right\} \\ &\quad + O_{j,k} \left(T^{\frac{1}{2}+b+\epsilon} \right). \end{aligned}$$

By Lemma 3.4, we deduce

$$\begin{aligned} S_2 &= (-1)^j \frac{T}{2\pi} \left(\log \frac{qT}{2\pi} \right)^{j+1} \sum_{m=0}^j \binom{j}{m} \sum_{u \leq \frac{\log T}{\log \log T}} \sum_{v=0}^k (-1)^u \binom{k}{v} \\ &\quad \times \sum_{i_1+i_2+\dots+i_k=u} \binom{u}{i_1, i_2, \dots, i_k} \prod_{w=1}^k \binom{k}{w}^{i_w} \frac{(-1)^{K+m+1}(v+1)!m! \prod_{w=1}^k (w!)^{i_w}}{(K+m+1)!} + O_{j,k} (T(\log qT)^j) \end{aligned}$$

This last sum S_2 was evaluated by Karabulut and Yildirim in [10]

$$S_2 = (-1)^j \frac{T}{2\pi} \left(\log \frac{qT}{2\pi} \right)^{j+1} B(j, k) + O_{j,k} (T(\log qT)^j)$$

Combining S_1 and S_2 , we obtain

$$I_4 = (-1)^j (\delta_{j,0} + B(j, k)) \frac{T}{2\pi} \left(\log \frac{qT}{2\pi} \right)^{j+1} + O_{j,k} \left(T (\log qT)^j \right).$$

Finally, theorem 1.1 follows from estimates of I_1, I_2, I_3 , and I_4 .

4. Proof of Theorem 1.2

Let a be a complex number. We write $s = \sigma + it$, $\rho_{a,\chi}^{(k)} = \beta_{a,\chi}^{(k)} + i\gamma_{a,\chi}^{(k)}$ with real numbers $\sigma, t, \beta_{a,\chi}^{(k)}$ and $\gamma_{a,\chi}^{(k)}$. The case $a = 0$ was already proven in Theorem 1.1, so here we assume $a \neq 0$. By the residue theorem, for a sufficiently large constant B and constant $b \in (1, 9/8)$, we have

$$\sum_{\substack{1 < \gamma_{a,\chi}^{(k)} < T \\ 1-b < \beta_{a,\chi}^{(k)} < B}} L^{(j)} \left(\rho_{a,\chi}^{(k)}, \chi \right) = \frac{1}{2\pi i} \int_{\mathbf{R}} L^{(j)}(s, \chi) \frac{L^{(k+1)}(s, \chi)}{L^{(k)}(s, \chi) - a} ds, \tag{4.1}$$

where the integration is taken over a rectangular contour in counterclockwise direction denoted by \mathbf{R} with vertices $1 - b + i, B + i, B + iT, 1 - b + iT$. Since there are finitely many a -points in $\{s \in \mathbb{C}; \operatorname{Re}(s) \leq 1 - b, \operatorname{Im}(s) \geq 1\}$, we have

$$\sum_{1 < \gamma_{a,\chi}^{(k)} < T} L^{(j)} \left(\rho_{a,\chi}^{(k)}, \chi \right) = \frac{1}{2\pi i} \int_{\mathbf{R}} L^{(j)}(s, \chi) \frac{L^{(k+1)}(s, \chi)}{L^{(k)}(s, \chi) - a} ds + O(1).$$

Hence,

$$\begin{aligned} & \sum_{1 < \gamma_{a,\chi}^{(k)} < T} L^{(j)} \left(\rho_{a,\chi}^{(k)}, \chi \right) \tag{4.2} \\ &= \frac{1}{2\pi i} \left\{ \int_{1-b+i}^{B+i} + \int_{B+i}^{B+iT} + \int_{B+iT}^{1-b+iT} + \int_{1-b+iT}^{1-b+i} \right\} L^{(j)}(s, \chi) \frac{L^{(k+1)}(s, \chi)}{L^{(k)}(s, \chi) - a} ds + O(1) \\ &:= \frac{1}{2\pi i} (I_1 + I_2 + I_3 + I_4) + O(1). \end{aligned}$$

The integral I_1 is independent of T , so we have $I_1 = O(1)$. Next, we consider I_2 . Since $L^{(k)}(s, \chi) \rightarrow 0$ as $\sigma \rightarrow \infty$ if $k \geq 1$, we choose in this case B such that $|L^{(k)}(B + it, \chi)| < \frac{|a|}{2}$, then we have $\frac{1}{L^{(k)}(B+it,\chi)-a} \ll_k 1$. Using this and $L^{(j)}(s, \chi) \ll 1$, we get

$$I_2 = O(T).$$

For the case $k = 0$, recall that, for $\sigma \rightarrow \infty$, we have $L(s, \chi) = 1 + o(1)$ and $L'(s, \chi) \ll 2^{-\sigma}$ uniformly in t . Hence, there are no a -points for sufficiently large σ provided that $a \neq 1$. For the case $a = 1$, we define $m = \min\{n \geq 2, \chi(n) \neq 0\}$. We observe, for $\sigma \rightarrow \infty$, $L(s, \chi) - 1 = \frac{\chi(m)}{m^{\sigma+it}}(1 + o(1))$. Hence, we choose B a fixed constant sufficiently large such that there are no a -points of $L(s, \chi)$ in the half-plane $\sigma > B - 1$. Therefore, we deduce that

$$I_2 = O(T).$$

From equation (2.12), we get

$$I_3 = \sum_{|\gamma_{a,\chi}^{(k)} - T| < 1} \int_{B+iT}^{1-b+iT} \frac{L^{(j)}(s, \chi)}{s - \rho_{a,\chi}^{(k)}} ds + O \left(\int_{B+iT}^{1-b+iT} (\log qt) L^{(j)}(s, \chi) ds \right).$$

Now, we change the path of integration. If $\gamma_{a,\chi}^{(k)} < T$, we change the path to the upper semicircle with center $\rho_{a,\chi}^{(k)}$ and radius 1. If $\gamma_{a,\chi}^{(k)} > T$, we change the path to the lower semicircle with center $\rho_{a,\chi}^{(k)}$ and radius 1. Then, we have

$$\frac{1}{s - \rho_{a,\chi}^{(k)}} \ll 1$$

on the new path. This estimate and the bound (21) yields

$$I_3 = O \left((qT)^{b-\frac{1}{2}+\epsilon} \sum_{|\gamma_{a,\chi}^{(k)} - T| < 1} 1 \right) + O \left((qT)^{b-\frac{1}{2}+\epsilon} \log qT \right).$$

By Lemma 2.2, we obtain

$$I_3 = O \left((qT)^{b-\frac{1}{2}+\epsilon} \log qT \right).$$

This yields $I_3 \ll T$, since $1 < b < 9/8$.

Finally, we estimate I_4 . By equation (2.20) and Stirling's formula, for fixed $1 < b < 9/8$ and large $|t| > 2$, we have

$$\left| L^{(k)}(1 - b + it, \chi) \right| \asymp |qt|^{b-1/2} |\log |t||^k. \tag{4.3}$$

Therefore, there exists a constant A such that

$$\left| \frac{a}{L^{(k)}(1 - b + it, \chi)} \right| < 1$$

holds for any $|t| \geq A$. We divide the path of the integral into two parts

$$I_4 = \left(\int_{1-b+iT}^{1-b+iA} + \int_{1-b+iA}^{1-b+i} \right) L^{(j)}(s, \chi) \frac{L^{(k+1)}(s, \chi)}{L^{(k)}(s, \chi) - a} ds.$$

The second term is $O(1)$ since it is independent of T . Since the integrand of the first term has a geometric series, we have

$$I_4 = - \sum_{n=0}^{\infty} a^n \int_{1-b+iA}^{1-b+iT} \frac{L^{(j)}(s, \chi) L^{(k+1)}(s, \chi)}{(L^{(k)}(s, \chi))^{n+1}} ds + O(1).$$

By (4.3), the integrand can be estimated as

$$\frac{L^{(j)}(s, \chi) L^{(k+1)}(s, \chi)}{(L^{(k)}(s, \chi))^{n+1}} \asymp |qt|^{(b-1/2)(1-n)} (\log t)^{-kn+j+1}. \tag{4.4}$$

Hence, each integral can be calculated as

$$\int_{1-b+iA}^{1-b+iT} \frac{L^{(j)}(s, \chi)L^{(k+1)}(s, \chi)}{(L^{(k)}(s, \chi))^{n+1}} ds \ll (qT)^{(b-1/2)(1-n)+1+\varepsilon}$$

for any small $\varepsilon > 0$. It follows from the last estimate that the sum for $n \geq 2$ is bounded as

$$\sum_{n=2}^{\infty} a^n \int_{1-b+iA}^{1-b+iT} \frac{L^{(j)}(s, \chi)L^{(k+1)}(s, \chi)}{(L^{(k)}(s, \chi))^{n+1}} ds \ll T^{-(b-1/2)+1+\varepsilon} \ll T^{1/2}.$$

Therefore, we get

$$\begin{aligned} I_4 &= - \int_{1-b+iA}^{1-b+iT} \frac{L^{(j)}(s, \chi)L^{(k+1)}(s, \chi)}{L^{(k)}(s, \chi)} ds - a \int_{1-b+iA}^{1-b+iT} \frac{L^{(j)}(s, \chi)L^{(k+1)}(s, \chi)}{(L^{(k)}(s, \chi))^2} ds + O\left(T^{1/2}\right) \\ &:= -K_1 - aK_2 + O\left(T^{1/2}\right). \end{aligned}$$

We already studied K_1 in Theorem 1.1 and we get the estimate

$$K_1 = -2\pi i \left\{ \delta_{j,0} \frac{T}{2\pi} \log \frac{qT}{2\pi} + (-1)^j B(j, k) \frac{T}{2\pi} \left(\log \frac{T}{2\pi} \right)^{j+1} + O\left(T(\log qT)^j\right) \right\}.$$

It remains to evaluate K_2 . By equation (4.4), for $k \geq 1$, one has

$$K_2 \ll \int_{1-b+iA}^{1-b+iT} |\log t|^j |ds| \ll T(\log T)^j.$$

In the case $k = 0$, we use equations (2.1) and (2.6) to obtain

$$\frac{L^{(j)}(s, \chi)L'(s, \chi)}{L^2(s, \chi)} = (-1)^{j+1} \ell^{j+1} \left(1 + O\left(\frac{1}{|t|}\right) \right) \tag{4.5}$$

for fixed σ and $|t| \gg 1$, where $\ell := \log(q|t|/2\pi)$. Then, we have

$$\begin{aligned} K_2 &= \int_{1-b+iA}^{1-b+iT} \left((-\ell)^{j+1} + O\left((\log q|t|)^j\right) \right) ds \\ &= (-1)^{j+1} iT \left(\log \frac{qT}{2\pi} \right)^{j+1} + O\left(T(\log qT)^j\right). \end{aligned}$$

Combining estimates of K_1 and K_2 , we get

$$I_4 = (-1)^j 2\pi i (\delta_{j,0} + a\delta_{k,0} + B(j, k)) \frac{T}{2\pi} \left(\log \frac{qT}{2\pi} \right)^{j+1} + O\left(T(\log qT)^j\right).$$

Finally, Theorem 1.2 follows from estimates of I_1, I_2, I_3 and I_4 .

5. Concluding remarks

The a -points of an L -function $L(s)$ are the roots of the equation $L(s) = a$. We refer to Steuding’s book [14, chapter 7] for some results about a -points of L -functions from the Selberg class. Therefore, it is an interesting question to extend Theorem 1.1 and mainly Theorem 1.2 to the other class of Dirichlet L -functions (the Selberg class with some further condition) and its higher derivative. This problem will be considered in a sequel to this paper since it is done for the Riemann zeta function and its k -th derivative in [6] and [12].

Acknowledgment

The authors would like to express their sincere gratitude to the referee for her/his many valuable suggestions which increased the clarity of the presentation.

References

- [1] Conrey JB, Ghosh A Zeros of derivatives of the Riemann zeta-function near the critical line. *Analytic Number Theory* (Allerton Park, IL 1989), Progr. Math. Birkhauser Boston, Boston, MA 1990; 85: 95–110.
- [2] Davenport H. *Multiplicative Number Theory*. 2nd ed. revised by H. L. Montgomery. Springer, 1980.
- [3] Fujii A. On the distribution of values of derivative of the Riemann zeta function at its zeros.I. *Proceedings of the Steklov Institute of Mathematics* 2012; 276: 51-76.
- [4] Garunkštis R, Steuding J. On the roots of the equation $\zeta(s) = a$. *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg* 2014; 84: 1-15.
- [5] Gonek SM. Mean values of the Riemann zeta-function and its derivatives. *Inventiones Mathematicae* 1984; 75: 123-141.
- [6] Jakhlouti MT, Mazhouda K. Distribution of the values of the derivative of the Dirichlet L -functions at its a -points. *Bulletin of the Korean Mathematical Society* 2017; 54 (4): 1141-1158.
- [7] Kaptan DA, Karabulut Y, Yildirim C. Some mean value theorems for the Riemann zeta-function and Dirichlet L -functions. *Commentarii Mathematici Universitatis Sancti Pauli* 2011; 60 (1-2): 83-87.
- [8] Kaptan DA. Some mean values related to Dirichlet L -functions. MSc, Boğazici University, İstanbul, Turkey, 2009; <http://seyhan.library.boun.edu.tr/record=b1558882> S5.
- [9] Karabulut Y. Some mean value problems concerning the Riemann zeta function. MSc, Boğazici University, İstanbul, Turkey, 2009; <http://seyhan.library.boun.edu.tr/record=b1558899> S5.
- [10] Karabulut Y, Yildirim C. On some averages at the zeros of the derivatives of the Riemann zeta-function. *Journal of Number Theory* 2011; 131: 1939-1961.
- [11] Levinson N. More than one third of the zeros of Riemann's zeta function are on $\sigma = \frac{1}{2}$. *Advances in Mathematics* 1974; 13: 386-436.
- [12] Mazhouda K, Onozuka T. On some sums at the a -points of derivatives of the Riemann zeta-function. To appear in *Analysis, Geometry and Number Theory*, 2020.
- [13] Onozuka T. On the a -points of the derivatives of the Riemann zeta function. *European Journal of Mathematics* 2017; 3 (1): 53-76.
- [14] Steuding J. *Value-distribution of L -functions*. Lecture Notes in Mathematics 1877, Springer 2007.
- [15] Titchmarsh EC. *The Theory of the Riemann Zeta-function*. 2nd ed, revised by D. R. Heath-Brown. Oxford, UK: Oxford University Press, 1986.
- [16] Yildirim C. Zeros of derivatives of Dirichlet L -functions. *Turkish Journal of Math* 1996; 20: 521-534.