

## On $q$ - and $h$ -deformations of 3d-superspaces

SALİH ÇELİK\* 

Department of Mathematics, Yıldız Technical University, İstanbul, Turkey

Received: 18.07.2019

Accepted/Published Online: 08.12.2020

Final Version: 21.01.2021

**Abstract:** In this paper, we introduce nonstandard deformations of  $(1+2)$ - and  $(2+1)$ -superspaces via a contraction using standard deformations of them. This deformed superspaces are denoted by  $\mathbb{A}_h^{1|2}$  and  $\mathbb{A}_{h'}^{2|1}$ , respectively. We find a two-parameter  $R$ -matrix satisfying quantum Yang–Baxter equation and thus obtain a new two-parameter nonstandard deformation of the supergroup  $GL(1|2)$ . Finally, we get a new superalgebra derived from the Hopf superalgebra of functions on the quantum superspace  $\mathbb{A}_{p,q}^{1|2}$ .

**Key words:** Quantum superspace, Hopf superalgebra, quantum supergroup, quantum Lie superalgebra, super  $\star$ -algebra

### 1. Introduction

There are two distinct deformations for general Lie (super)groups as standard and nonstandard (or Jordanian). One of them is the well-known quantum ( $q$ -deformed) group and the other is the so-called Jordanian ( $h$ -deformed) one. Specially, quantum groups  $GL_q(2)$  [10] and  $GL_h(2)$  [9] have been obtained by deforming the coordinates of a plane to be noncommutative objects. In [1], the authors showed that the  $h$ -deformed group can be obtained from the  $q$ -deformed Lie group through a singular limit  $q \rightarrow 1$  of a linear transformation. This method is known as the contraction procedure. Using this method, one- and two-parameter  $h$ -deformations of supergroup  $GL(1|1)$  were obtained in [7] and [2], respectively.

In this paper, we give some standard (as  $q$ -deformation) deformations of  $(1+2)$ -superspace using the Hopf superalgebra structure of  $\mathcal{O}(\mathbb{A}^{1|2})$  and nonstandard (as  $h$ -deformation) deformations using standard deformations via a contraction. We also introduce an  $(h, h')$ -deformed supergroup acting on these two-parameter  $h$ -deformed superspaces. Finally, we define involutions on  $h$ -deformed superspaces and use the generators of  $(p, q)$ -deformed superalgebra  $\mathcal{O}(\mathbb{A}_{p,q}^{1|2})$  to get a new Lie superalgebra.

Throughout the paper, we will fix a base field  $\mathbb{K}$ . The reader may consider it as the set of real numbers,  $\mathbb{R}$ , or the set of complex numbers,  $\mathbb{C}$ . We will denote by  $\mathbb{G}$  the Grassmann numbers and by  $\mathbb{K}'$  the set  $\mathbb{K} \cup \mathbb{G}$ .

### 2. On $(p, q)$ -deformation of superspaces $\mathbb{A}^{1|2}$ and $\mathbb{A}^{2|1}$

In order to define superalgebras and Hopf superalgebras, some minor changes are made in familiar definitions. These are briefly mentioned in the following.

A supervector space  $\mathcal{X}$  over a field  $\mathbb{K}$  is a  $\mathbb{Z}_2$ -graded vector space  $\mathcal{X}$  together with two subspaces  $\mathcal{X}_0$  and  $\mathcal{X}_1$  of  $\mathcal{X}$  such that  $\mathcal{X} = \mathcal{X}_0 \oplus \mathcal{X}_1$ . If a space  $\mathcal{X}$  is a superspace, then we denote by  $\tau(a)$  the  $\mathbb{Z}_2$ -grade of

\*Correspondence: sacelik@yildiz.edu.tr

the element  $a \in \mathcal{X}$ . If  $\tau(a) = 0$ , then we will call the element  $a$  even and if  $\tau(a) = 1$ , it is called odd.

If  $f : \mathcal{X} \rightarrow \mathcal{Y}$  is a linear map of supervector spaces and it satisfies

$$\tau(f(v)) = \tau(f) + \tau(v) \pmod{2}$$

for all  $v \in \mathcal{X}$ , then  $f$  is called a supervector space homomorphism.

A superalgebra (or  $\mathbb{Z}_2$ -graded algebra)  $\mathcal{A}$  over  $\mathbb{K}$  is a supervector space over  $\mathbb{K}$  with a map  $\mathcal{A} \times \mathcal{A} \rightarrow \mathcal{A}$  such that  $\mathcal{A}_i \cdot \mathcal{A}_j \subset \mathcal{A}_{i+j}$  for  $i, j = 0, 1$ . The superalgebra  $\mathcal{A}$  is called supercommutative if

$$ab = (-1)^{\tau(a)\tau(b)}ba$$

for homogeneous elements  $a, b \in \mathcal{A}$ .

Let  $f : \mathcal{A} \rightarrow \mathcal{B}$  be a map of definite degree of superalgebras. If it is a supervector space homomorphism and it obeys

$$f(ab) = (-1)^{\tau(a)\tau(f)}f(a)f(b), \quad \forall a, b \in \mathcal{A},$$

then  $f$  is called a superalgebra homomorphism.

### 2.1. The algebra of polynomials on the quantum superspace $\mathbb{A}_q^{1|2}$

Let  $\mathbb{K}\langle X, \Theta_1, \Theta_2 \rangle$  be a free algebra with unit generated by  $X$ ,  $\Theta_1$ , and  $\Theta_2$ , where the coordinate  $X$  is even and the coordinates  $\Theta_1$  and  $\Theta_2$  are odd.

**Definition 2.1** [11] Let  $I_q$  be the two-sided ideal of  $\mathbb{K}\langle X, \Theta_1, \Theta_2 \rangle$  generated by the elements  $X\Theta_1 - q\Theta_1X$ ,  $X\Theta_2 - q\Theta_2X$ ,  $\Theta_1\Theta_2 + q^{-1}\Theta_2\Theta_1$ ,  $\Theta_1^2$ , and  $\Theta_2^2$ . The quantum superspace  $\mathbb{A}_q^{1|2}$  with the function algebra

$$\mathcal{O}(\mathbb{A}_q^{1|2}) = \mathbb{K}\langle X, \Theta_1, \Theta_2 \rangle / I_q$$

is called  $\mathbb{Z}_2$ -graded quantum space (or quantum superspace).

This associative algebra over the complex number is known as the algebra of polynomials over quantum (1+2)-superspace. In accordance with the above definition, we have

$$X\Theta_i = q\Theta_iX, \quad \Theta_i\Theta_j = -q^{i-j}\Theta_j\Theta_i, \quad (i, j = 1, 2) \tag{2.1}$$

where  $q \in \mathbb{K} - \{0\}$ .

**Example 2.2** If we consider the generators of the algebra  $\mathcal{O}(\mathbb{A}_q^{1|2})$  as linear maps, then we can find the matrix representations of them. In fact, it can be seen that there exists a representation  $\rho : \mathcal{O}(\mathbb{A}_q^{1|2}) \rightarrow M(3, \mathbb{K}')$  such that matrices

$$\rho(X) = \begin{pmatrix} q & 0 & 0 \\ 0 & q & 0 \\ 0 & 0 & q^2 \end{pmatrix}, \quad \rho(\Theta_1) = \begin{pmatrix} 0 & 0 & \varepsilon_1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \rho(\Theta_2) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \varepsilon_2 \\ 0 & 0 & 0 \end{pmatrix} \tag{2.2}$$

representing the coordinate functions satisfy relations (2.1) for all  $\varepsilon_1, \varepsilon_2$ .

**Remark 2.3** In the next section, we will assume that  $\varepsilon_1$  and  $\varepsilon_2$  are two Grassmann numbers.

The following definition gives the product rule for tensor product of  $\mathbb{Z}_2$ -graded algebras.

**Definition 2.4** *The product rule is defined by*

$$(a_1 \otimes a_2)(a_3 \otimes a_4) = (-1)^{\tau(a_2)\tau(a_3)}(a_1a_3 \otimes a_2a_4)$$

in the  $\mathbb{Z}_2$ -graded algebra  $\mathcal{A} \otimes \mathcal{A}$ , where  $\mathcal{A}$  is the  $\mathbb{Z}_2$ -graded algebra and  $a_i$ 's are homogeneous elements in  $\mathcal{A}$ .

A Hopf superalgebra is a supervector space  $\mathcal{A}$  over  $\mathbb{K}$  with two algebra homomorphisms  $\Delta : \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$ , called the coproduct,  $\epsilon : \mathcal{A} \rightarrow \mathbb{K}$ , called the counit, and an algebra antihomomorphism  $S : \mathcal{A} \rightarrow \mathcal{A}$ , called the antipode, such that

$$\begin{aligned} (\Delta \otimes \text{id}) \circ \Delta &= (\text{id} \otimes \Delta) \circ \Delta, \\ m \circ (\epsilon \otimes \text{id}) \circ \Delta &= \text{id} = m \circ (\text{id} \otimes \epsilon) \circ \Delta, \\ m \circ (S \otimes \text{id}) \circ \Delta &= \eta \circ \epsilon = m \circ (\text{id} \otimes S) \circ \Delta, \end{aligned}$$

and  $\Delta(\mathbf{1}) = \mathbf{1} \otimes \mathbf{1}$ ,  $\epsilon(\mathbf{1}) = 1$ ,  $S(\mathbf{1}) = \mathbf{1}$ , where  $m$  is the multiplication map,  $\text{id}$  is the identity map and  $\eta : \mathbb{K} \rightarrow \mathcal{A}$ .

*Note.* An element of a Hopf superalgebra  $\mathcal{A}$  is expressed as a product on the generators and its antipode  $S$  is calculated with the property

$$S(ab) = (-1)^{\tau(a)\tau(b)}S(b)S(a), \quad \forall a, b \in \mathcal{A}.$$

We denote the unital extension of  $\mathcal{O}(\mathbb{A}_q^{1|2})$  by  $\mathcal{F}(\mathbb{A}_q^{1|2})$  adding the unit and  $x^{-1}$ , the inverse of  $x$ , which obeys  $xx^{-1} = \mathbf{1} = x^{-1}x$ . The following theorem says that the superalgebra  $\mathcal{F}(\mathbb{A}_q^{1|2})$  has a Hopf algebra structure [4]:

**Theorem 2.5** [4] *The superalgebra  $\mathcal{F}(\mathbb{A}_q^{1|2})$  is a Hopf superalgebra with the defining coproduct, counit, and antipode on the algebra  $\mathcal{F}(\mathbb{A}_q^{1|2})$  as follows:*

(1) *The coproduct  $\Delta : \mathcal{F}(\mathbb{A}_q^{1|2}) \rightarrow \mathcal{F}(\mathbb{A}_q^{1|2}) \otimes \mathcal{F}(\mathbb{A}_q^{1|2})$  is defined by*

$$\Delta(X) = X \otimes X, \quad \Delta(\Theta_1) = \Theta_1 \otimes X + X \otimes \Theta_1, \quad \Delta(\Theta_2) = \Theta_2 \otimes X^2 + X^2 \otimes \Theta_2. \quad (2.3)$$

(2) *The counit  $\epsilon : \mathcal{F}(\mathbb{A}_q^{1|2}) \rightarrow \mathbb{K}$  is given by*

$$\epsilon(X) = 1, \quad \epsilon(\Theta_i) = 0, \quad (i = 1, 2).$$

(3) *The algebra  $\mathcal{F}(\mathbb{A}_q^{1|2})$  admits a  $\mathbb{K}$ -algebra antihomomorphism (antipode)  $S : \mathcal{F}(\mathbb{A}_q^{1|2}) \rightarrow \mathcal{F}(\mathbb{A}_{q^{-1}}^{1|2})$  defined by*

$$S(X) = X^{-1}, \quad S(\Theta_1) = -X^{-1}\Theta_1X^{-1}, \quad S(\Theta_2) = -X^{-2}\Theta_2X^{-2}.$$

## 2.2. The algebra of polynomials on the quantum superspace $\mathbb{A}_{p,q}^{2|1}$

Let  $\mathbb{K}\langle \Phi, Y_1, Y_2 \rangle$  be a free algebra with unit generated by  $\Phi$ ,  $Y_1$  and  $Y_2$ , where  $\tau(\Phi) = 1$  and  $\tau(Y_1) = 0 = \tau(Y_2)$ .

**Definition 2.6** [5] Let  $\Lambda(\mathbb{A}_q^{1|2})$  be the algebra with the generators  $\Phi$ ,  $Y_1$ , and  $Y_2$  satisfying the relations

$$\Phi^2 = 0, \quad \Phi Y_1 = qp^{-1}Y_1\Phi, \quad \Phi Y_2 = pqY_2\Phi, \quad Y_1Y_2 = pq^{-1}Y_2Y_1. \quad (2.4)$$

We call  $\Lambda(\mathbb{A}_q^{1|2})$  exterior algebra of the  $\mathbb{Z}_2$ -graded space  $\mathbb{A}_q^{1|2}$ .

**Remark 2.7** The exterior algebra  $\Lambda(\mathbb{A}_q^{1|2})$  of the superspace  $\mathbb{A}_q^{1|2}$  can be thought of as a two-parameter deformation of the  $(2+1)$ -superspace  $\mathbb{A}^{2|1}$ . Thus, we denote this algebra by  $\mathcal{O}(\mathbb{A}_{p,q}^{2|1})$ .

**Example 2.8** If we consider the generators of the algebra  $\mathcal{O}(\mathbb{A}_{p,q}^{2|1})$  as linear maps, then we can find the matrix representations of them. In fact, it can be seen that there exists a representation  $\rho : \mathcal{O}(\mathbb{A}_{p,q}^{2|1}) \rightarrow M(3, \mathbb{K}')$  such that matrices

$$\rho(\Phi) = \begin{pmatrix} 0 & 0 & \epsilon \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \rho(Y_1) = \begin{pmatrix} q & 0 & 0 \\ 0 & p & 0 \\ 0 & 0 & p \end{pmatrix}, \quad \rho(Y_2) = \begin{pmatrix} 0 & c & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

representing the coordinate functions satisfy relations (2.4) for all  $c, \epsilon$ .

### 3. Two-parameter $h$ -deformation of the superspaces

In this section, we introduce a two-parameter  $h$ -deformation of the superspace  $\mathbb{A}^{1|2}$  (and its dual) from the  $(p, q)$ -deformation via a contraction similar to the method of [1].

We consider the  $q$ -deformed algebra of functions on the quantum superspace  $\mathbb{A}_q^{1|2}$  generated by  $X$ ,  $\Theta_1$ , and  $\Theta_2$  with the relations (2.1) and we introduce new even coordinate  $x$  and odd coordinates  $\theta_1$ ,  $\theta_2$  with the change of basis in the coordinates of the  $q$ -superspace using the following  $g$  matrix:

$$\mathbf{X} = \begin{pmatrix} X \\ \Theta_1 \\ \Theta_2 \end{pmatrix} = \begin{pmatrix} 1 & 0 & \tilde{h}' \\ 0 & 1 & 0 \\ \tilde{h} & 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ \theta_1 \\ \theta_2 \end{pmatrix} = g\mathbf{x}, \quad \tilde{h} = \frac{h}{q-1}, \quad \tilde{h}' = \frac{h'}{pq-1} \quad (3.1)$$

where  $h$  and  $h'$  ( $h \neq 0 \neq h'$ ) are two new deformation parameters that will be replaced with  $q$  and  $p$  ( $q \neq 1 \neq pq$ ) in the limits  $q \rightarrow 1$  and  $p \rightarrow 1$ .

We now assume that the parameters  $h$  and  $h'$  are both Grassmann numbers ( $h^2 = 0 = h'^2$ ,  $hh' = -h'h$ ) and anticommute with  $\theta_i$  for  $i = 1, 2$ . When the relations (2.1) are used, one gets

$$x\theta_1 = q\theta_1x, \quad x\theta_2 = q\theta_2x + hx^2, \quad \theta_2\theta_1 = -q\theta_1\theta_2, \quad \theta_1^2 = 0, \quad \theta_2^2 = -h\theta_2x. \quad (3.2)$$

Note that the parameter  $h'$  does not enter the above relations. By taking the limit  $q \rightarrow 1$ , we obtain the following exchange relations, which define the  $h$ -superspace  $\mathbb{A}_h^{1|2}$ :

**Definition 3.1** [4] Let  $\mathcal{O}(\mathbb{A}_h^{1|2})$  be the algebra with the generators  $x$ ,  $\theta_1$ , and  $\theta_2$  satisfying the relations

$$x\theta_1 = \theta_1x, \quad x\theta_2 = \theta_2x + hx^2, \quad \theta_1\theta_2 = -\theta_2\theta_1, \quad \theta_1^2 = 0, \quad \theta_2^2 = -h\theta_2x. \quad (3.3)$$

We call  $\mathcal{O}(\mathbb{A}_h^{1|2})$  the algebra of functions on the  $\mathbb{Z}_2$ -graded quantum space  $\mathbb{A}_h^{1|2}$ .

**Example 3.2** Let us assume that  $\varepsilon_1$  and  $\varepsilon_2$  are two Grassmann numbers. If the  $g$  matrix in (3.1) is used, the matrix representation in (2.2) takes the following form:

$$\rho(x) = q \begin{pmatrix} 1 - \tilde{h}\tilde{h}' & 0 & 0 \\ 0 & 1 - \tilde{h}\tilde{h}' & -q^{-1}\tilde{h}'\varepsilon_2 \\ 0 & 0 & q(1 - \tilde{h}\tilde{h}') \end{pmatrix}, \quad \rho(\theta_1) = \begin{pmatrix} 0 & 0 & \varepsilon_1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \rho(\theta_2) = - \begin{pmatrix} q\tilde{h} & 0 & 0 \\ 0 & q\tilde{h} & -(1 + \tilde{h}\tilde{h}')\varepsilon_2 \\ 0 & 0 & q^2\tilde{h} \end{pmatrix}. \tag{3.4}$$

These matrices satisfy the relations (3.2), for all  $\varepsilon_1$  and  $\varepsilon_2$ .

**Proof** Existing claims come from the fact that  $\rho$  is an algebra homomorphism. □

In the case of dual (exterior)  $h'$ -superspace, we use the transformation

$$\hat{\mathbf{X}} = g\hat{\mathbf{x}} \tag{3.5}$$

with the components  $\varphi$ ,  $y_1$ , and  $y_2$  of  $\hat{\mathbf{x}}$ . The definition is given below.

**Definition 3.3** Let  $\mathcal{O}(\mathbb{A}_{h'}^{2|1}) := \Lambda(\mathbb{A}_h^{1|2})$  be the algebra with the generators  $\varphi$ ,  $y_1$ , and  $y_2$  satisfying the relations

$$\varphi y_1 = y_1 \varphi, \quad \varphi y_2 = y_2 \varphi + h' y_2^2, \quad y_1 y_2 = y_2 y_1, \quad \varphi^2 = h' y_2 \varphi \tag{3.6}$$

where  $\tau(\varphi) = 1$  and  $\tau(y_1) = 0 = \tau(y_2)$ . We call  $\Lambda(\mathbb{A}_h^{1|2})$  the quantum exterior algebra of the  $\mathbb{Z}_2$ -graded quantum space  $\mathbb{A}_h^{1|2}$ .

**Remark 3.4** The parameter  $h$  does not enter the relations (3.6). The exterior algebra  $\Lambda(\mathbb{A}_h^{1|2})$  of the superspace  $\mathbb{A}_h^{1|2}$  can be thought of as an  $h'$ -deformation of the (2+1)-superspace  $\mathbb{A}^{2|1}$ .

#### 4. An $R$ -matrix and its properties

The relations in (2.1) can be written in a compact form as follows:

$$p \mathbf{X} \otimes \mathbf{X} = \hat{R}_{p,q} \mathbf{X} \otimes \mathbf{X} \tag{4.1}$$

with an  $R$ -matrix given by [6]

$$\hat{R}_{p,q} = \begin{pmatrix} p & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & p-1 & 0 & q & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & pq & 0 & 0 \\ 0 & pq^{-1} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -pq^{-1} & 0 \\ 0 & 0 & q^{-1} & 0 & 0 & 0 & p-1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -q & 0 & p-1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \end{pmatrix}$$

where  $p, q \in \mathbb{K} - \{0\}$ . This matrix satisfies the graded braid equation and the matrix  $R_{p,q} = P\hat{R}_{p,q}$  satisfies the graded Yang–Baxter equation where  $P$  is the super permutation matrix.

It can be considered that a change of basis in the quantum superspaces leads to a two-parameter  $R$ -matrix. The corresponding  $R$ -matrix can be obtained as

$$\hat{R}_{h,h'} = \lim_{(p,q) \rightarrow (1,1)} \left[ (g \otimes g)^{-1} \hat{R}_{p,q} (g \otimes g) \right]$$

where it is assumed that  $\otimes$  is graded. As a result, we obtain the following  $R$ -matrix

$$\hat{R}_{h,h'} = \begin{pmatrix} 1 + hh' & 0 & h' & 0 & 0 & 0 & -h' & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ h & 0 & hh' & 0 & 0 & 0 & 1 & 0 & -h' \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ -h & 0 & 1 & 0 & 0 & 0 & hh' & 0 & -h' \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & -h & 0 & 0 & 0 & -h & 0 & hh' - 1 \end{pmatrix}.$$

The equation in (4.1) with the new  $R$ -matrix  $\hat{R}_{h,h'}$  takes the form

$$\mathbf{x} \otimes \mathbf{x} = \hat{R}_{h,h'} \mathbf{x} \otimes \mathbf{x},$$

that is, the relations (3.3) are equivalent to this equation.

The  $R$ -matrix  $\hat{R}_{h,h'}$  has some interesting properties. Some of them are listed below, where sometimes we write  $\hat{R} = \hat{R}_{h,h'}$  for simplicity.

1. The matrix  $\hat{R}_{h,h'}$  satisfies the graded braid equation  $\hat{R}_{12} \hat{R}_{23} \hat{R}_{12} = \hat{R}_{23} \hat{R}_{12} \hat{R}_{23}$ , where  $\hat{R}_{12} = \hat{R} \otimes I_3$  and  $\hat{R}_{12} = I_3 \otimes \hat{R}$ .
2. The matrix  $R_{h,h'} = P \hat{R}_{h,h'}$  satisfies the graded Yang–Baxter equation  $R_{12} R_{13} R_{23} = R_{23} R_{13} R_{12}$ , where  $R_{13}$  acts both on the first and third spaces.
3. The matrix  $\hat{R}_{h,h'}$  holds  $\hat{R}_{h,h'}^2 = I_9$ ; thus, it has two eigenvalues  $\pm 1$ .
4. If we set  $hh' = 0$ , then the matrix  $R_{h,h'}$  can be decomposed in the form

$$R_{h,h'} = R(h)R(h')$$

where

$$R(h) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -h & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ h & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & h & 0 & 0 & 0 & h & 0 & 1 \end{pmatrix}, \quad R(h') = R^{\text{st}}(h)|_{h=h'}.$$

It can be checked that these matrices satisfy the graded Yang–Baxter equation.

5. If  $P_{\pm}$  are the projections onto the eigenspaces  $\pm 1$  of  $\hat{R}_{h,h'}$ , then we have

$$\hat{R}_{h,h'} = P_+ - P_-.$$

Let  $\mathcal{O}(\mathbb{A}^{1|2})$  and  $\mathcal{O}(\mathbb{A}^{2|1})$  be the quotients of algebras generated by  $x, \theta_1, \theta_2$  and  $\varphi, y_1, y_2$  modulo the two-sided ideals generated by  $\text{Ker}P_-$  and  $\text{Ker}P_+$ , respectively. Then  $\mathcal{O}(\mathbb{A}^{1|2})$  and  $\mathcal{O}(\mathbb{A}^{2|1})$  are isomorphic to  $\mathcal{O}(\mathbb{A}_h^{1|2})$  with defining relations (3.3) and  $\mathcal{O}(\mathbb{A}_{h'}^{2|1})$  with defining relations (3.6), respectively. That is, we can write

$$P_- \mathbf{x} \otimes \mathbf{x} = 0 \quad \text{and} \quad (-1)^{\tau(\hat{\mathbf{x}})} P_+ \hat{\mathbf{x}} \otimes \hat{\mathbf{x}} = 0.$$

### 5. The quantum superbialgebra $\mathcal{O}(M_{h,h'}(1|2))$

Let  $T$  be a 3x3 matrix in  $\mathbb{Z}_2$ -graded space given by

$$T = \begin{pmatrix} a & \alpha & \beta \\ \gamma & b & c \\ \delta & d & e \end{pmatrix} = (t_{ij})$$

where  $a, b, c, d, e$  are even and  $\alpha, \beta, \gamma,$  and  $\delta$  are odd. The coordinate ring of such matrices over a field  $\mathbb{K}$  is simply the polynomial ring in nine variables, that is  $\mathcal{O}(M(1|2)) = \mathbb{K}[a, b, c, d, e, \alpha, \beta, \gamma, \delta]$ .

In this section, we will assume that the matrix entries of  $T$  belong to a free superalgebra and define a two-parameter  $h$ -analogue of  $\mathcal{O}(M(1|2))$ . To do so, let  $x, \theta_1, \theta_2$  be elements of the superalgebra  $\mathcal{O}(\mathbb{A}_h^{1|2})$  subject to the relations (3.3) and  $\varphi, y_1, y_2$  be elements of  $\mathcal{O}(\mathbb{A}_{h'}^{2|1})$  subject to the relations (3.6), and  $t_{ij}$  be nine generators which supercommute with the elements of  $\mathcal{O}(\mathbb{A}_h^{1|2})$  and  $\mathcal{O}(\mathbb{A}_{h'}^{2|1})$ . It is well known that the supermatrix  $T$  defines the linear transformations  $T : \mathbb{A}_h^{1|2} \rightarrow \mathbb{A}_h^{1|2}$  and  $T : \mathbb{A}_{h'}^{2|1} \rightarrow \mathbb{A}_{h'}^{2|1}$ . Let  $\mathbf{x} = (x, \theta_1, \theta_2)^t$  and  $\hat{\mathbf{x}} = (\varphi, y_1, y_2)^t$ . Thus, we can give the following theorem.

**Theorem 5.1** *Under the above hypotheses, the following conditions are equivalent:*

- (i)  $T\mathbf{x} = \mathbf{x}' \in \mathbb{A}_h^{1|2}$  and  $T\hat{\mathbf{x}} = \hat{\mathbf{x}}' \in \mathbb{A}_{h'}^{2|1}$ ,

(ii) the relations are satisfied

$$\begin{aligned}
 a\alpha &= (1 + hh')\alpha a - h'(\alpha\delta + da), & a\beta &= \beta a + h'(a^2 - ea - \beta\delta) - h\beta^2, \\
 a\gamma &= (1 + hh')\gamma a + h(\gamma\beta - ca), & ac &= ca - hc\beta - h'\gamma a + hh'\gamma\beta, \\
 a\delta &= \delta a + h(a^2 - ea + \delta\beta) + h'\delta^2, & ad &= da + h\alpha a + h'd\delta - hh'\alpha\delta, \\
 ae &= ea + h\beta(a - e) + h'(e - a)\delta, & \alpha\beta &= -(1 + hh')\beta\alpha + h'(\beta d + e\alpha), \\
 \alpha\gamma &= -\gamma\alpha, & \alpha c &= c\alpha, & \alpha\delta &= -\delta\alpha - ha\alpha + h'\delta d - hh'ad, \\
 \alpha d &= d\alpha + h'd^2, & \alpha e &= e\alpha + h\beta\alpha + h'ed - hh'd\beta, \\
 \beta\gamma &= -\gamma\beta + hc\beta - h'\gamma a - hh'ca, & \beta c &= (1 - hh')c\beta - h'(\gamma\beta + ca), \\
 \beta\delta &= -\delta\beta + (h\beta + h'\delta)(e - a), & \beta d &= d\beta + h\alpha\beta + h'de - hh'e\alpha, \\
 \beta e &= e\beta + h'(e^2 - ea - \delta\beta) - h\beta^2, & \gamma c &= c\gamma + hc^2, \\
 \gamma\delta &= -(1 + hh')\delta\gamma + h(e\gamma + \delta c), & \gamma d &= d\gamma, \\
 \gamma e &= e\gamma + hec - h'\delta\gamma - hh'c\delta, & c\delta &= \delta c - hec - h'\delta\gamma - hh'\gamma e, \\
 cd &= dc, & ce &= (1 - hh')ec + h'(e\gamma - \delta c), & \delta d &= (1 - hh')d\delta + h(\alpha\delta - da), \\
 \delta e &= e\delta + h(e^2 - ea + \beta\delta) + h'\delta^2, & de &= (1 - hh')ed + h(\beta d - e\alpha), \\
 \alpha^2 &= h'\alpha d, & \beta^2 &= h'\beta(e - a), & \gamma^2 &= h\gamma c, & \delta^2 &= h\delta(e - a), \\
 bt_{ij} &= t_{ij}b, & a(h\beta + h'\delta) &= (h\beta + h'\delta)a, & e(h\beta + h'\delta) &= (h\beta + h'\delta)e.
 \end{aligned} \tag{5.1}$$

**Proof** A direct verification shows that the relations (5.1) respect the ideals defining  $\mathbb{A}_h^{1|2}$  and  $\mathbb{A}_{h'}^{2|1}$ . □

Standard FRT construction [8], namely, the relations (5.1), is obtained via the matrix  $\hat{R}_{h,h'}$  given in Section 4:

**Theorem 5.2** A 3x3-matrix  $T$  is a  $\mathbb{Z}_2$ -graded quantum supermatrix if and only if

$$\hat{R}_{h,h'}T_1T_2 = T_1T_2\hat{R}_{h,h'}$$

where  $T_1 = T \otimes I_3$  and  $T_2 = PT_1P$ .

**Definition 5.3** The superalgebra  $\mathcal{O}(M_{h,h'}(1|2))$  is the quotient of the free algebra  $\mathbb{K}\{a, b, c, d, e, \alpha, \beta, \gamma, \delta\}$  by the two-sided ideal  $J_{h,h'}$  generated by the relations (5.1) of Theorem 5.1.

**Remark 5.4** The quantum matrix space  $M_{p,q}(1|2)$  is obtained in [6]. It is clear that a change of basis in the quantum superspace leads to the similarity transformation  $T = g^{-1}T'g$ , where  $T' \in M_{p,q}(1|2)$ . Therefore, the entries of the transformed quantum matrix  $T$  fulfill the commutation relations (5.1) of the matrix elements of the matrix  $T$  in  $M(1|2)$ .

**Theorem 5.5** The superalgebra  $\mathcal{O}(M_{h,h'}(1|2))$  with the following two algebra homomorphisms of superalgebras (1) the coproduct  $\Delta : \mathcal{O}(M_{h,h'}(1|2)) \rightarrow \mathcal{O}(M_{h,h'}(1|2)) \otimes \mathcal{O}(M_{h,h'}(1|2))$  determined by  $\Delta(t_{ij}) = \sum_{k=1}^3 t_{ik} \otimes t_{kj}$ ,



(2) the counit  $\epsilon : \mathcal{O}(M_{h,h'}(1|2)) \rightarrow \mathbb{K}$  determined by  $\epsilon(t_{ij}) = \delta_{ij}$  becomes a super bialgebra.

**Proof** It can be easily checked the properties of the costructures hold:

(i) The coproduct  $\Delta$  is coassociative in the sense of

$$(\Delta \otimes \text{id}) \circ \Delta = (\text{id} \otimes \Delta) \circ \Delta$$

where  $\text{id}$  denotes the identity map on  $M_{h,h'}(1|2)$  and  $\Delta(ab) = \Delta(a)\Delta(b)$ ,  $\Delta(\mathbf{1}) = \mathbf{1} \otimes \mathbf{1}$ .

(ii) The counit  $\epsilon$  has the property

$$m \circ (\epsilon \otimes \text{id}) \circ \Delta = \text{id} = m \circ (\text{id} \otimes \epsilon) \circ \Delta$$

where  $m$  stands for the algebra product and  $\epsilon(ab) = \epsilon(a)\epsilon(b)$ ,  $\epsilon(\mathbf{1}) = 1$ . □

It is well known that  $\mathcal{O}(\mathbb{A}^{1|2})$  is comodule algebra over the bialgebra  $\mathcal{O}(M(1|2))$ . The following theorem gives a quantum version of this fact.

**Theorem 5.6** *There exist algebra homomorphisms*

$$\begin{aligned} \delta_L : \mathcal{O}(\mathbb{A}_h^{1|2}) &\longrightarrow \mathcal{O}(M_{h,h'}(1|2)) \otimes \mathcal{O}(\mathbb{A}_h^{1|2}), & \delta_L(x_i) &= \sum_{k=1}^3 t_{ik} \otimes x_k, \\ \tilde{\delta}_L : \mathcal{O}(\mathbb{A}_{h'}^{2|1}) &\longrightarrow \mathcal{O}(M_{h,h'}(1|2)) \otimes \mathcal{O}(\mathbb{A}_{h'}^{2|1}), & \tilde{\delta}_L(\hat{x}_i) &= \sum_{k=1}^3 t_{ik} \otimes \hat{x}_k \end{aligned}$$

where  $x_i \in \{x, \theta_1, \theta_2\}$  and  $\hat{x}_i \in \{\varphi, y_1, y_2\}$ .

**Proof** Using the relations (3.3) and (3.6) together with (5.1), it is enough to check that

$$\delta_L(x\theta_1 - \theta_1x) = \delta_L(x)\delta_L(\theta_1) - \delta_L(\theta_1)\delta_L(x) = 0,$$

etc., in  $\mathcal{O}(M_{h,h'}(1|2)) \otimes \mathcal{O}(\mathbb{A}_h^{1|2})$ . To see that  $\delta_L$  defines a comodule structure we check that

$$(\Delta \otimes \text{id}) \circ \delta_L = (\text{id} \otimes \delta_L) \circ \delta_L, \quad m \circ (\epsilon \otimes \text{id}) \circ \delta_L = \text{id}.$$

□

A quantum supergroup (Hopf superalgebra) can be regarded as a generalization of the notion of a supergroup. It is defined by

$$\mathcal{O}(\text{GL}_{h,h'}(1|2)) = \mathcal{O}(M_{h,h'}(1|2))[t]/(t \text{sdet}_{h,h'} - 1).$$

This case is also inviting to generalize the corresponding notions of differential geometry [12]. A differential calculus on  $\mathcal{O}(\text{GL}_{h,h'}(1|2))$  will be discussed in the next work.

### 6. A Lie superalgebra derived from $\mathcal{F}(\mathbb{A}_{p,q}^{1|2})$

It is known that an element of a Lie group can be represented by exponential of an element of its Lie algebra.

In [3], by virtue of this fact, using the generators of the superalgebra  $\mathcal{F}(\mathbb{A}_q^{1|1})$ , a new superalgebra is obtained

from this algebra. In this section, we will obtain a new superalgebra from  $\mathcal{F}(\mathbb{A}_{p,q}^{1|2})$ . Thus, let us begin with the definition of  $\mathcal{F}(\mathbb{A}_{p,q}^{1|2})$  which is an extension to two parameters of  $\mathcal{F}(\mathbb{A}_q^{1|2})$ .

**Definition 6.1** Let  $I_{p,q}$  be the two-sided ideal of  $\mathbb{K}\langle X, \Theta_1, \Theta_2 \rangle$  generated by the elements  $X\Theta_1 - q\Theta_1X$ ,  $X\Theta_2 - p\Theta_2X$ ,  $\Theta_1\Theta_2 + pq^{-2}\Theta_2\Theta_1$ ,  $\Theta_1^2$ , and  $\Theta_2^2$ . The quantum superspace  $\mathbb{A}_{p,q}^{1|2}$  with the function algebra

$$\mathcal{O}(\mathbb{A}_{p,q}^{1|2}) = \mathbb{K}\langle X, \Theta_1, \Theta_2 \rangle / I_{p,q}$$

is called quantum superspace.

In accordance with this definition, we have

$$X\Theta_1 = q\Theta_1X, \quad X\Theta_2 = p\Theta_2X, \quad \Theta_1\Theta_2 = -pq^{-2}\Theta_2\Theta_1, \quad \Theta_i^2 = 0 \tag{6.1}$$

where  $p, q \in \mathbb{K} - \{0\}$ .

**Example 6.2** If we consider the generators of the algebra  $\mathcal{O}(\mathbb{A}_{p,q}^{1|2})$  as linear maps, then we can find the matrix representations of them. In fact, it can be seen that there exists a representation  $\rho : \mathcal{O}(\mathbb{A}_{p,q}^{1|2}) \rightarrow M(3, \mathbb{K}')$  such that matrices

$$\rho(X) = \begin{pmatrix} q & 0 & 0 \\ 0 & p & 0 \\ 0 & 0 & pq \end{pmatrix}, \quad \rho(\Theta_1) = \begin{pmatrix} 0 & 0 & \varepsilon_1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \rho(\Theta_2) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \varepsilon_2 \\ 0 & 0 & 0 \end{pmatrix}$$

representing the coordinate functions satisfy relations (6.1) for all  $\varepsilon_1, \varepsilon_2$ .

Let  $\mathbb{K}\langle u, \xi_1, \xi_2 \rangle$  be a free algebra generated by  $u, \xi_1, \xi_2$ , where  $\tau(u) = 0, \tau(\xi_1) = 1 = \tau(\xi_2)$ . Let  $\mathcal{L}$  be the quotient of the free algebra  $\mathbb{K}\langle u, \xi_1, \xi_2 \rangle$  by the two-sided ideal  $J_0$  generated by the elements  $u\xi_k - \xi_k u, \xi_1\xi_2 + \xi_2\xi_1, \xi_k^2$  for  $k = 1, 2$ .

Now, we will show that the Hopf superalgebra of Theorem 2.5 can be embedded into the enveloping superalgebra of a Lie superalgebra, with Lie structure and a deformed coproduct. Thus, let us define the generators of the algebra  $\mathcal{F}(\mathbb{A}_{p,q}^{1|2})$  as

$$X := e^u, \quad \Theta_k := e^{ku}\xi_k,$$

for  $k = 1, 2$ . The first equality implies that the generator  $X$  is invertible. Then, by direct calculations we can prove the following lemma.

**Lemma 6.3** The generators  $u, \xi_1, \xi_2$  have the following commutation relations (Lie (anti-)brackets), for  $j, k = 1, 2$

$$[u, \xi_k] = \mathbf{i} \hbar_k \xi_k, \quad [\xi_j, \xi_k]_+ = 0, \tag{6.2}$$

where  $q = e^{\mathbf{i}\hbar_1}, p = e^{\mathbf{i}\hbar_2}$  with  $\mathbf{i} = \sqrt{-1}$  and  $\hbar_1, \hbar_2 \in \mathbb{R}$ .

We denote the algebra for which the generators obey the relations (6.2) by  $\mathcal{L}_{\hbar_1, \hbar_2} := \mathcal{L}(\mathbb{A}_{p,q}^{1|2})$ . Let  $U(\mathcal{L}_{\hbar_1, \hbar_2})$  be the algebra defined by (6.2). The Hopf superalgebra structure of  $U(\mathcal{L}_{\hbar_1, \hbar_2})$  can be read off from Theorem 2.5:

**Theorem 6.4** *The superalgebra  $U(\mathcal{L}_{\hbar_1, \hbar_2})$  is a Hopf superalgebra with coproduct, counit, and antipode on the algebra  $\mathcal{L}_{\hbar_1, \hbar_2}$  defined by*

$$\Delta(u_i) = u_i \otimes \mathbf{1} + \mathbf{1} \otimes u_i, \quad \epsilon(u_i) = 0, \quad S(u_i) = -u_i.$$

for  $u_i \in \{u, \xi_1, \xi_2\}$ .

**Example 6.5** *There exists a Lie algebra homomorphism  $\mu$  from  $\mathcal{L}_{\hbar_1, \hbar_2}$  into  $M(3, \mathbb{K}')$ .*

**Proof** We see that there exists an algebra homomorphism  $\rho$  from  $\mathcal{F}(\mathbb{A}_{p,q}^{1|2})$  into  $M(3, \mathbb{K}')$  such that the relations (6.1) hold. As a consequence of this fact, there exists a Lie algebra homomorphism  $\mu$  from  $\mathcal{L}_{\hbar_1, \hbar_2}$  into  $M(3, \mathbb{K}')$ . The action of  $\mu$  on the generators of  $\mathcal{L}_{\hbar_1, \hbar_2}$  is of the form

$$\mu(u) = \begin{pmatrix} i\hbar_2 & 0 & 0 \\ 0 & i\hbar_1 & 0 \\ 0 & 0 & i(\hbar_1 + \hbar_2) \end{pmatrix}, \quad \mu(\xi_1) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ e^{-i(\hbar_1 + \hbar_2)\varepsilon_1} & 0 & 0 \end{pmatrix}, \quad \mu(\xi_2) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & e^{-2i(\hbar_1 + \hbar_2)\varepsilon_2} & 0 \end{pmatrix} \quad (6.3)$$

where  $\varepsilon_1$  and  $\varepsilon_2$  are two Grassmann numbers. To see that the relations (6.2) are preserved under the action of  $\mu$ , we use the fact that

$$\mu[a, b] = [\mu(a), \mu(b)],$$

for all  $a, b \in \mathcal{L}_{\hbar_1, \hbar_2}$ . □

### 7. $\star$ -Structures on the algebras $\mathcal{O}(\mathbb{A}_h^{1|2})$ and $\mathcal{O}(\mathbb{A}_{h'}^{2|1})$

It is possible to define the star operation (or involution) on the Grassmann generators. However, there are two possibilities to do so\*. If  $\alpha$  and  $\beta$  are two Grassmann generators and  $\lambda$  is a complex number and  $\bar{\lambda}$  its complex conjugate, the star operation, denoted by  $\star$ , is defined by

$$(\lambda\alpha)^\star = \bar{\lambda}\alpha^\star, \quad (\alpha\beta)^\star = \beta^\star\alpha^\star, \quad (\alpha^\star)^\star = \alpha$$

and the superstar operation, denoted by  $\#$ , is defined by

$$(\lambda\alpha)^\# = \bar{\lambda}\alpha^\#, \quad (\alpha\beta)^\# = \alpha^\#\beta^\#, \quad (\alpha^\#)^\# = -\alpha.$$

It is easily shown that there exists a star operation on the algebra  $\mathcal{O}(\mathbb{A}_q^{1|2})$  if  $q$  is a complex number of modulus one:

**Proposition 7.1** (i) *If  $\bar{q} = q^{-1}$  then the algebra  $\mathcal{O}(\mathbb{A}_q^{1|2})$  equipped with the involution determined by*

$$X^\star = X, \quad \Theta_i^\star = \Theta_i \quad (i = 1, 2) \quad (7.1)$$

becomes a  $\star$ -algebra.

(ii) *If  $\bar{p} = p^{-1}$  and  $\bar{q} = q^{-1}$  then the algebra  $\mathcal{O}(\mathbb{A}_{p,q}^{2|1})$  equipped with the involution determined by*

$$\Phi^\star = \Phi, \quad Y_i^\star = -Y_i \quad (i = 1, 2) \quad (7.2)$$

becomes a  $\star$ -algebra.

---

\*arXiv.org e-Print archive (1996). Dictionary on Lie Superalgebras [online]. Website <https://arxiv.org/abs/hep-th/9607161> [18 July 1996].

**7.1.  $\star$ -Structures on the algebra  $\mathcal{O}(\mathbb{A}_h^{1|2})$**

As noted in Section 3, the relations in (3.3) do not include the parameter  $h'$ . Thus, we can rearrange the change of basis in the coordinates (see, equation (3.1)) as

$$\begin{pmatrix} X \\ \Theta_1 \\ \Theta_2 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ \frac{h}{q-1} & 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ \theta_1 \\ \theta_2 \end{pmatrix}. \tag{7.3}$$

This case can help us to define a star operation on the algebra  $\mathcal{O}(\mathbb{A}_h^{1|2})$  by a coordinate transformation using the generators of the algebra  $\mathcal{O}(\mathbb{A}_q^{1|2})$  and to prove the following lemma.

**Lemma 7.2** *For a certain special choice of  $h$ , there exists an involution on the algebra  $\mathcal{O}(\mathbb{A}_h^{1|2})$ .*

**Proof** Using the equation (7.3), we introduce the coordinates  $x$ ,  $\theta_1$ , and  $\theta_2$  with the change of basis in the coordinates of the superspace  $\mathbb{A}_q^{1|2}$  as follows:

$$x = X, \quad \theta_1 = \Theta_1, \quad \theta_2 = \Theta_2 - \frac{h}{q-1} X.$$

Then, with  $|q| = 1$  and (7.1)

$$\theta_2^* = \Theta_2^* - \frac{q\bar{h}}{1-q} X^* = \theta_2 + \frac{h+q\bar{h}}{q-1} x$$

so that, if we demand that  $\bar{h} = -h$ , we obtain  $\theta_2^* = \theta_2 - hx$ . Note that

$$(x^*)^* = x, \quad (\theta_1^*)^* = \theta_1, \quad (\theta_2^*)^* = \theta_2,$$

for all  $h$ . □

**Proposition 7.3** *If  $\bar{h} = -h$ , then the algebra  $\mathcal{O}(\mathbb{A}_h^{1|2})$  supplied with the involution determined by*

$$x^* = x, \quad \theta_1^* = \theta_1, \quad \theta_2^* = \theta_2 - hx \tag{7.4}$$

*becomes a  $\star$ -algebra.*

**Proof** Since  $\bar{h} = -h$ , we have

$$\begin{aligned} (x\theta_1 - \theta_1x)^* &= \theta_1x - x\theta_1, \\ (x\theta_2 - \theta_2x - hx^2)^* &= (\theta_2 - hx)x - x(\theta_2 - hx) + hx^2 = (\theta_2x - x\theta_2 + hx^2), \\ (\theta_1\theta_2 + \theta_2\theta_1)^* &= (\theta_2 - hx)\theta_1 + \theta_1(\theta_2 - hx) = \theta_2\theta_1 + \theta_1\theta_2, \\ (\theta_2^2 + h\theta_2x)^* &= (\theta_2 - hx)(\theta_2 - hx) + x(\theta_2 - hx)(-h) = \theta_2^2 + h\theta_2x. \end{aligned}$$

Hence, the ideal  $(x\theta_1 - \theta_1x, x\theta_2 - \theta_2x - hx^2, \theta_1\theta_2 + \theta_2\theta_1, \theta_1^2, \theta_2^2 + h\theta_2x)$  is  $\star$ -invariant and the quotient algebra

$$\mathbb{K}\langle x, \theta_1, \theta_2 \rangle / (x\theta_1 - \theta_1x, x\theta_2 - \theta_2x - hx^2, \theta_1\theta_2 + \theta_2\theta_1, \theta_1^2, \theta_2^2 + h\theta_2x)$$

becomes a  $\star$ -algebra. □

**Remark 7.4** *Of course, we can consider the change of basis in the coordinates of the superspace  $\mathbb{A}_q^{1|2}$  in (3.1). In this case, since*

$$\begin{aligned} x^* &= (1 + \tilde{h}\tilde{h}' - \overline{\tilde{h}\tilde{h}'} )x + (\tilde{h}' - \overline{\tilde{h}'})\theta_2, \\ \theta_1^* &= \theta_1, \\ \theta_2^* &= (1 - \overline{\tilde{h}}(\tilde{h}' - \overline{\tilde{h}'}))\theta_2 + (\tilde{h} - \overline{\tilde{h}})x, \end{aligned}$$

*we have again (7.4) with the choices  $\bar{h} = -h$  and  $\bar{h}' = h'$ .*

**7.2.  $\star$ -Structure on the algebra  $\mathcal{O}(\mathbb{A}_{h'}^{2|1})$**

Since the relations in (3.6) do not include the parameter  $h$ , we can rearrange the change of basis in the coordinates (see, equation (3.1)) as

$$\begin{pmatrix} \Phi \\ Y_1 \\ Y_2 \end{pmatrix} = \begin{pmatrix} 1 & 0 & \frac{h'}{pq-1} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \varphi \\ y_1 \\ y_2 \end{pmatrix}. \tag{7.5}$$

There exists a special case, where the algebra  $\mathcal{O}(\mathbb{A}_{h'}^{2|1})$  admits an involution. The proofs of the following lemma and proposition can be done in a similar way to Lemma 7.2 and Proposition 7.3.

**Lemma 7.5** *If  $\bar{h}' = h'$ , there exists an involution on the algebra  $\mathcal{O}(\mathbb{A}_{h'}^{2|1})$ .*

**Proposition 7.6** *If  $\bar{h}' = h'$ , then the algebra  $\mathcal{O}(\mathbb{A}_h^{2|1})$  supplied with the involution determined by*

$$\varphi^* = \varphi - h'y_2, \quad y_i^* = -y_i, \quad (i = 1, 2) \tag{7.6}$$

*becomes a  $\star$ -algebra.*

**References**

[1] Aghamohammadi A, Khorrami M, Shariati A.  $h$ -deformation as a contraction of  $q$ -deformation. Jordanian deformation of  $SL(2)$  as a contraction of its Drinfeld-Jimbo deformation. Journal of Physics A: Mathematical and General 1995; 28 (8): L225-L231.

[2] Celik S. Two-parametric extension of  $h$ -deformation of  $GL(1|1)$ . Letters in Mathematical Physics 1997; 42: 299-308.

[3] Celik S. Differential geometry of the Lie algebra of the quantum superplane. Balkan Physics Letters 2003; 11: 119-127. (arXiv:math/0201170)

[4] Celik S. Bicovariant differential calculus on the quantum superspace  $\mathbb{R}_q(1|2)$ . Journal of Algebra and its Applications 2016; 15: 1650172 (17 pages).

[5] Celik Sultan A. Differential calculi on super-Hopf algebra  $\mathcal{F}(\mathbb{R}_q(1|2))$ . Advanced in Applied Clifford Algebras 2018; 28: 85:1-16.

[6] Celik Sultan A. A two-parameter deformation of supergroup  $GL(1|2)$ . Journal of the Institute of Science and Technology 2018; 8(4): 271-280.

[7] Dabrowski L, Parashar P.  $h$ -deformation of  $GL(1|1)$ . Letters in Mathematical Physics 1996; 38: 331-336.

- [8] Faddeev LD, Reshetikhin NY, Takhtajan LA. Quantization of Lie groups and Lie algebras. Leningrad Mathematical Journal 1990; 1 (1): 193-225.
- [9] Kupershmidt BA. The quantum group  $GL_h(2)$ . Journal of Physics A: Mathematical and General 1992; 25: L1239-L1244.
- [10] Manin YI. Quantum groups and noncommutative geometry. Montreal, Canada: Centre de Recherches Mathematiques, Universite de Montreal, 1988.
- [11] Manin YI. Multiparametric quantum deformation of the general linear supergroup. Communications in Mathematical Physics 1989; 123 (1): 163-175.
- [12] Woronowicz SL. Differential calculus on compact matrix pseudogroups (quantum groups). Communications in Mathematical Physics 1989; 122 (1): 125-170.