

## Hilbert series of tangent cones for Gorenstein monomial curves in $\mathbb{A}^4(K)$

Anargyros KATSABEKIS\* 

Department of Mathematics, Faculty of Science, Bilkent University, Ankara, Turkey

Received: 14.11.2020

Accepted/Published Online: 27.12.2020

Final Version: 21.01.2021

**Abstract:** In this paper, we study the Hilbert series of the tangent cone of Gorenstein monomial curves in the 4-dimensional affine space. We give an explicit formula for the reduced Hilbert series of the tangent cone of a noncomplete intersection Gorenstein monomial curve whose tangent cone is Cohen–Macaulay.

**Key words:** Hilbert series, tangent cone, Gorenstein monomial curve

### 1. Introduction

Let  $n_1 < n_2 < \dots < n_d$  be positive integers with  $\gcd(n_1, \dots, n_d) = 1$ . Consider the polynomial ring  $R = K[x_1, \dots, x_d]$  in  $d$  variables over a field  $K$ . We shall denote by  $\mathbf{x}^{\mathbf{u}}$  the monomial  $x_1^{u_1} \cdots x_d^{u_d}$  of  $R$ , with  $\mathbf{u} = (u_1, \dots, u_d) \in \mathbb{N}^d$ , where  $\mathbb{N}$  stands for the set of nonnegative integers. Consider the affine monomial curve in the  $d$ -dimensional affine space  $\mathbb{A}^d(K)$  defined parametrically by  $x_1 = t^{n_1}, \dots, x_d = t^{n_d}$ . The toric ideal of  $C$ , denoted by  $I_C$ , is the kernel of the  $K$ -algebra homomorphism  $\varphi : R \rightarrow K[t]$  given by

$$\varphi(x_i) = t^{n_i} \text{ for all } 1 \leq i \leq d.$$

The ideal  $I_C$  is generated by all the binomials  $\mathbf{x}^{\mathbf{u}} - \mathbf{x}^{\mathbf{v}}$  such that  $\varphi(\mathbf{x}^{\mathbf{u}}) = \varphi(\mathbf{x}^{\mathbf{v}})$  see for example, [7, Lemma 4.1]. Given a polynomial  $f \in I_C$ , we let  $f^*$  be the homogeneous summand of  $f$  of least degree. We shall denote by  $I_C^*$  the ideal in  $R$  generated by the polynomials  $f^*$  for  $f \in I_C$ .

Let  $m = \langle t^{n_1}, \dots, t^{n_d} \rangle$  be the maximal ideal of the one-dimensional local ring  $A = K[[t^{n_1}, \dots, t^{n_d}]]$ . The Hilbert function  $H_A$  of  $A$  is defined by  $H_A(i) = \dim_{A/m}(m^i/m^{i+1})$  for every  $i \in \mathbb{N}$ , which coincides, by its definition, with the Hilbert function of the associated graded ring  $\text{gr}_m(A) = \bigoplus_{i \geq 0} m^i/m^{i+1}$ . It is worth noting that  $\text{gr}_m(A)$  is isomorphic to the quotient  $R/I_C^*$ . We recall that  $I_C^*$  is the defining ideal of the tangent cone of  $C$  at the origin.

Given an ideal  $J \subset R$ , we shall denote by  $\text{HS}(R/J, z)$  the Hilbert series of the ring  $R/J$ , namely  $\text{HS}(R/J, z) = \sum_{i \in \mathbb{N}} H_{R/J}(i)z^i$  where  $H_{R/J}$  is the Hilbert function of  $R/J$ . By the Hilbert–Serre theorem,  $\text{HS}(R/J, z)$  is a rational function of the form  $\text{HS}(R/J, z) = \frac{p(z)}{(1-z)^d}$  for some  $p(z) \in \mathbb{Z}[z]$ . In particular, by reducing this rational function we get  $\text{HS}(R/J, z) = \frac{h(z)}{(1-z)^e}$  for some  $h(z) \in \mathbb{Z}[z]$ , where  $e$  is the Krull dimension of  $R/J$ .

\*Correspondence: katsampekis@bilkent.edu.tr

2010 AMS Mathematics Subject Classification: 13H10, 14H20, 20M14

In this paper, we study the reduced Hilbert series of the ring  $R/I_C^*$ . Since the Krull dimension of the above ring is equal to 1, we can write  $HS(R/I_C^*, z) = \frac{h(z)}{1-z}$ . We restrict ourselves to Gorenstein noncomplete intersection monomial curves in  $\mathbb{A}^4(K)$ . Recall that a monomial curve  $C$  is called Gorenstein if the associated local ring  $A$  is Gorenstein. Furthermore we assume that  $R/I_C^*$  is a Cohen–Macaulay ring. The significance of this class is underscored by the following result: If  $R/I_C^*$  is Cohen–Macaulay, then the Hilbert function of  $A$  is nondecreasing. Our aim is to give an explicit formula for the numerator of the reduced Hilbert series of the ring  $R/I_C^*$  depending only on a minimal generating set of  $I_C$ .

In [4] Bresinsky provided a minimal generating set of  $I_C$  consisting of five generators. Actually, there are 6 permutations of the above generator set. In [1] the authors provided necessary and sufficient conditions for the Cohen–Macaulayness of  $R/I_C^*$  in all six permutations. We compute a standard basis for  $I_C$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$ . Using [3, Proposition 2.2] we determine a formula for the numerator of the reduced Hilbert series of  $R/LM(I_C^*)$ , where  $LM(I_C^*)$  is the ideal generated by the leading monomials of the polynomials of  $I_C^*$  with respect to the aforementioned order. By [5, Theorem 5.2.6] the reduced Hilbert series of  $R/I_C^*$  coincides with the reduced Hilbert series of  $R/LM(I_C^*)$ .

**2. Formulas for the reduced Hilbert series**

In this section we first recall Bresinsky’s theorem, which gives the explicit description of  $I_C$  when  $C$  is a Gorenstein noncomplete intersection monomial curve in  $\mathbb{A}^4(K)$ . By Kunz [6] it is well known that the Gorenstein property of a monomial curve is equivalent to the symmetric property of  $\mathbb{N}\{n_1, \dots, n_4\}$ .

**Theorem 2.1** ([4]) *Let  $C$  be a monomial curve having the parametrization*

$$x_1 = t^{n_1}, x_2 = t^{n_2}, x_3 = t^{n_3}, x_4 = t^{n_4}.$$

*The semigroup  $\mathbb{N}\{n_1, \dots, n_4\}$  is symmetric and  $C$  is a noncomplete intersection curve if and only if  $I_C$  is minimally generated by the set*

$$\{f_1 = x_1^{a_1} - x_3^{a_{13}}x_4^{a_{14}}, f_2 = x_2^{a_2} - x_1^{a_{21}}x_4^{a_{24}}, f_3 = x_3^{a_3} - x_1^{a_{31}}x_2^{a_{32}}, f_4 = x_4^{a_4} - x_2^{a_{42}}x_3^{a_{43}}, f_5 = x_1^{a_{21}}x_3^{a_{43}} - x_2^{a_{32}}x_4^{a_{14}}\}$$

*where the polynomials  $f_i$  are unique up to isomorphism,  $a_{ij} > 0$  and also*

$$a_1 = a_{21} + a_{31}, a_2 = a_{32} + a_{42}, a_3 = a_{13} + a_{43}, a_4 = a_{14} + a_{24}.$$

**Remark 2.2** ([2]) *Theorem 2.1 implies that for any noncomplete intersection Gorenstein monomial curve in  $\mathbb{A}^4(K)$ , the variables can be renamed to obtain generators exactly of the given form, and this means that there are six isomorphic possible permutations which can be considered within three cases:*

- (1)  $f_1 = (1, (3, 4))$ 
  - (a)  $f_2 = (2, (1, 4)), f_3 = (3, (1, 2)), f_4 = (4, (2, 3)), f_5 = ((1, 3), (2, 4))$
  - (b)  $f_2 = (2, (1, 3)), f_3 = (3, (2, 4)), f_4 = (4, (1, 2)), f_5 = ((1, 4), (2, 3))$
- (2)  $f_1 = (1, (2, 3))$ 
  - (a)  $f_2 = (2, (3, 4)), f_3 = (3, (1, 4)), f_4 = (4, (1, 2)), f_5 = ((2, 4), (1, 3))$

(b)  $f_2 = (2, (1, 4)), f_3 = (3, (2, 4)), f_4 = (4, (1, 3)), f_5 = ((1, 2), (4, 3))$

(3)  $f_1 = (1, (2, 4))$

(a)  $f_2 = (2, (1, 3)), f_3 = (3, (1, 4)), f_4 = (4, (2, 3)), f_5 = ((1, 2), (3, 4))$

(b)  $f_2 = (2, (3, 4)), f_3 = (3, (1, 2)), f_4 = (4, (1, 3)), f_5 = ((2, 3), (1, 4))$

Here, the notation  $f_i = (i, (j, k))$  and  $f_5 = ((i, j), (k, l))$  denote the generators  $f_i = x_i^{a_i} - x_j^{a_{ij}} x_k^{a_{ik}}$  and  $f_5 = x_i^{a_{ki}} x_j^{a_{lj}} - x_k^{a_{jk}} x_l^{a_{il}}$ . Thus, given a Gorenstein monomial curve  $C$ , if we have the extra condition  $n_1 < n_2 < n_3 < n_4$ , then the generator set of  $I_C$  is exactly given by one of these six permutations.

In [1] they provided necessary and sufficient conditions for the Cohen–Macaulayness of  $R/I_C^*$ . More precisely they proved the following.

**Theorem 2.3** ([1]) (1) Suppose that  $I_C$  is given as in case 1(a). Then  $R/I_C^*$  is Cohen–Macaulay if and only if  $a_2 \leq a_{21} + a_{24}$ .

(2) Suppose that  $I_C$  is given as in case 1(b). (i) Assume that  $a_{42} \leq a_{32}$ . Then  $R/I_C^*$  is Cohen–Macaulay if and only if

1.  $a_2 \leq a_{21} + a_{23}$ ,

2.  $a_{42} + a_{13} \leq a_{21} + a_{34}$ , and

3. either  $a_{34} < a_{14}$  and  $a_3 + a_{13} \leq a_{21} + a_{32} - a_{42} + 2a_{34}$  or  $a_{14} \leq a_{34}$  and  $a_3 + a_{13} \leq a_1 + a_{32} + a_{34} - a_{14}$ .

(ii) Assume that  $a_{32} < a_{42}$  and  $a_{14} \leq a_{34}$ . Then  $R/I_C^*$  is Cohen–Macaulay if and only if

1.  $a_2 \leq a_{21} + a_{23}$ ,

2.  $a_{42} + a_{13} \leq a_{21} + a_{34}$ , and

3.  $a_3 + a_{13} \leq a_1 + a_{32} + a_{34} - a_{14}$ .

(3) Suppose that  $I_C$  is given as in case 2(a). (i) Assume that  $a_{34} \leq a_{24}$ . Then  $R/I_C^*$  is Cohen–Macaulay if and only if

1.  $a_3 \leq a_{31} + a_{34}$ ,

2.  $a_{12} + a_{34} \leq a_{41} + a_{23}$ , and

3. either  $a_{23} < a_{13}$  and  $a_2 + a_{12} \leq a_{41} + 2a_{23} + a_{24} - a_{34}$  or  $a_{13} \leq a_{23}$  and  $a_2 + a_{12} \leq a_1 + a_{23} - a_{13} + a_{24}$ .

(ii) Assume that  $a_{24} < a_{34}$  and  $a_{13} \leq a_{23}$ . Then  $R/I_C^*$  is Cohen–Macaulay if and only if

1.  $a_3 \leq a_{31} + a_{34}$ ,

2.  $a_{12} + a_{34} \leq a_{41} + a_{23}$ , and

3.  $a_2 + a_{12} \leq a_1 + a_{23} - a_{13} + a_{24}$ .

(4) Suppose that  $I_C$  is given as in case 2(b). (i) Assume that  $a_{24} \leq a_{34}$ . Then  $R/I_C^*$  is Cohen–Macaulay if and only if

1.  $a_2 \leq a_{21} + a_{24}$  and
2. either  $a_{32} < a_{12}$  and  $a_3 + a_{13} \leq a_{41} + 2a_{32} + a_{34} - a_{24}$  or  $a_{12} \leq a_{32}$  and  $a_3 + a_{13} \leq a_1 + a_{32} - a_{12} + a_{34}$ .

(ii) Assume that  $a_{34} < a_{24}$  and  $a_{12} \leq a_{32}$ . Then  $R/I_C^*$  is Cohen–Macaulay if and only if

1.  $a_2 \leq a_{21} + a_{24}$  and
2.  $a_3 + a_{13} \leq a_1 + a_{32} - a_{12} + a_{34}$ .

(5) Suppose that  $I_C$  is given as in case 3(a). Then  $R/I_C^*$  is Cohen–Macaulay if and only if  $a_2 \leq a_{21} + a_{23}$  and  $a_3 \leq a_{31} + a_{34}$ .

(6) Suppose that  $I_C$  is given as in case 3(b). (i) Assume that  $a_{43} \leq a_{23}$ . Then  $R/I_C^*$  is Cohen–Macaulay if and only if

1.  $a_{12} + a_{43} \leq a_{31} + a_{24}$  and
2. either  $a_{24} < a_{14}$  and  $a_2 + a_{12} \leq a_{31} + 2a_{24} + a_{23} - a_{43}$  or  $a_{14} \leq a_{24}$  and  $a_2 + a_{12} \leq a_1 + a_{23} + a_{24} - a_{14}$ .

(ii) Assume that  $a_{23} < a_{43}$  and  $a_{14} \leq a_{24}$ . Then  $R/I_C^*$  is Cohen–Macaulay if and only if

1.  $a_{12} + a_{43} \leq a_{31} + a_{24}$  and
2.  $a_2 + a_{12} \leq a_1 + a_{23} + a_{24} - a_{14}$ .

For the rest of this section, we assume that  $R/I_C^*$  is a Cohen–Macaulay ring.

In the sequel, we will make repeatedly use of the next proposition.

**Proposition 2.4** ([3, Proposition 2.2]) Let  $I \subset R$  be a monomial ideal and let  $I = \langle J, \mathbf{x}^{\mathbf{u}} \rangle$  for a monomial ideal  $J$  and a monomial  $\mathbf{x}^{\mathbf{u}}$ . For an ideal  $M \subset R$  denote by  $p(M)$  the numerator of the Hilbert series of  $R/M$ . Then  $p(I) = p(J) - z^{\deg(\mathbf{x}^{\mathbf{u}})}p(J : \langle \mathbf{x}^{\mathbf{u}} \rangle)$ .

**Theorem 2.5** Suppose that  $I_C$  is given as in case 1(a). Then the reduced Hilbert series of  $R/I_C^*$  is  $HS(R/I_C^*, z) = \frac{h(z)}{1-z}$  for

$$h(z) = \prod_{i=2}^4 (1 + z + z^2 + \dots + z^{a_i-1}) - z^{a_{14}+a_{32}} \sum_{i=0}^{a_{42}-1} z^i \sum_{i=0}^{a_3-1} z^i \sum_{i=0}^{a_{24}-1} z^i - z^{a_{13}+a_{14}} \sum_{i=0}^{a_{32}-1} z^i \sum_{i=0}^{a_{43}-1} z^i \sum_{i=0}^{a_{24}-1} z^i.$$

**Proof.** By [2, Lemma 2.7],

$$G = \{x_1^{a_1} - x_3^{a_{13}}x_4^{a_{14}}, x_2^{a_2} - x_1^{a_{21}}x_4^{a_{24}}, x_3^{a_3} - x_1^{a_{31}}x_2^{a_{32}}, x_4^{a_4} - x_2^{a_{42}}x_3^{a_{43}}, x_1^{a_{21}}x_3^{a_{43}} - x_2^{a_{32}}x_4^{a_{14}}\}$$

is a standard basis for  $I_C$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$ . From [5, Lemma 5.5.11] the ideal  $I_C^*$  is generated by the least homogeneous summands of the elements in  $G$ . In addition,  $LM(I_C^*) = \langle x_3^{a_{13}}x_4^{a_{14}}, x_2^{a_2}, x_3^{a_3}, x_4^{a_4}, x_2^{a_{32}}x_4^{a_{14}} \rangle$ . Let

$$J_0 = \langle LM(I_C^*) \rangle, J_1 = \langle x_2^{a_2}, x_3^{a_3}, x_4^{a_4}, x_2^{a_{32}}x_4^{a_{14}} \rangle, J_2 = \langle x_2^{a_2}, x_3^{a_3}, x_4^{a_4} \rangle.$$

Then  $J_i = \langle J_{i+1}, q_i \rangle$ , where  $q_0 = x_3^{a_{13}}x_4^{a_{14}}$  and  $q_1 = x_2^{a_{32}}x_4^{a_{14}}$ . By Proposition 2.4,

$$p(J_i) = p(J_{i+1}) - z^{\deg(q_i)}p(J_{i+1} : \langle q_i \rangle), \text{ for } 0 \leq i \leq 1.$$

Note that  $\deg(q_0) = a_{13} + a_{14}$  and  $\deg(q_1) = a_{14} + a_{32}$ . In this case,  $J_1 : \langle q_0 \rangle = \langle x_2^{a_{32}}, x_3^{a_{43}}, x_4^{a_{24}} \rangle$  and  $J_2 : \langle q_1 \rangle = \langle x_2^{a_{42}}, x_3^{a_{43}}, x_4^{a_{24}} \rangle$ . We have that  $p(J_2) = \prod_{i=2}^4 (1 - z^{a_i})$  and  $p(J_2 : \langle q_1 \rangle) = (1 - z^{a_{42}})(1 - z^{a_{43}})(1 - z^{a_{24}})$ , so  $p(J_1) = \prod_{i=2}^4 (1 - z^{a_i}) - z^{a_{32}+a_{14}}(1 - z^{a_{42}})(1 - z^{a_{43}})(1 - z^{a_{24}})$ . Furthermore  $p(J_1 : \langle q_0 \rangle) = (1 - z^{a_{32}})(1 - z^{a_{43}})(1 - z^{a_{24}})$ , so  $p(J_0) = \prod_{i=2}^4 (1 - z^{a_i}) - z^{a_{14}+a_{32}}(1 - z^{a_{42}})(1 - z^{a_{43}})(1 - z^{a_{24}}) - z^{a_{13}+a_{14}}(1 - z^{a_{32}})(1 - z^{a_{43}})(1 - z^{a_{24}})$ . Now using the fact that  $1 - z^b = (1 - z) \sum_{i=0}^{b-1} z^i$  we get

$$p(J_0) = (1 - z)^3 \sum_{i=0}^{a_2-1} z^i \sum_{i=0}^{a_3-1} z^i \sum_{i=0}^{a_4-1} z^i - (1 - z)^3 z^{a_{14}+a_{32}} \sum_{i=0}^{a_{42}-1} z^i \sum_{i=0}^{a_3-1} z^i \sum_{i=0}^{a_{24}-1} z^i - (1 - z)^3 z^{a_{13}+a_{14}} \sum_{i=0}^{a_{32}-1} z^i \sum_{i=0}^{a_{43}-1} z^i \sum_{i=0}^{a_{24}-1} z^i.$$

Thus  $p(J_0) = (1 - z)^3 h(z)$ , and therefore  $\text{HS}(R/\text{LM}(I_C^*), z) = \frac{h(z)}{1-z}$ . □

We continue with case 1(b). If  $a_3 \leq a_{32} + a_{34}$ , then we can use [2, Remark 2.9] to find the reduced Hilbert series of  $R/I_C^*$ .

**Theorem 2.6** *Suppose that  $I_C$  is given as in case 1(b) and also that  $a_3 \leq a_{32} + a_{34}$ . Then the reduced Hilbert series of  $R/I_C^*$  is  $\text{HS}(R/I_C^*, z) = \frac{h(z)}{1-z}$  for*

$$h(z) = \prod_{i=2}^4 (1 + z + z^2 + \dots + z^{a_i-1}) - z^{a_{13}+a_{42}} \sum_{i=0}^{a_{32}-1} z^i \sum_{i=0}^{a_{23}-1} z^i \sum_{i=0}^{a_4-1} z^i - z^{a_{13}+a_{14}} \sum_{i=0}^{a_{42}-1} z^i \sum_{i=0}^{a_{23}-1} z^i \sum_{i=0}^{a_{34}-1} z^i.$$

**Proof.** By [2, Remark 2.9],

$$G = \{x_1^{a_1} - x_3^{a_{13}}x_4^{a_{14}}, x_2^{a_2} - x_1^{a_{21}}x_3^{a_{23}}, x_3^{a_3} - x_2^{a_{32}}x_4^{a_{34}}, x_4^{a_4} - x_1^{a_{41}}x_2^{a_{42}}, x_1^{a_{21}}x_4^{a_{34}} - x_2^{a_{42}}x_3^{a_{13}}\}$$

is a standard basis for  $I_C$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$ . From [5, Lemma 5.5.11] the ideal  $I_C^*$  is generated by the least homogeneous summands of the elements in  $G$ . Also  $\text{LM}(I_C^*) = \langle x_3^{a_{13}}x_4^{a_{14}}, x_2^{a_2}, x_3^{a_3}, x_4^{a_4}, x_2^{a_{42}}x_3^{a_{13}} \rangle$ . Let

$$J_0 = \text{LM}(I_C^*), J_1 = \langle x_2^{a_2}, x_3^{a_3}, x_4^{a_4}, x_2^{a_{42}}x_3^{a_{13}} \rangle, J_2 = \langle x_2^{a_2}, x_3^{a_3}, x_4^{a_4} \rangle.$$

Then  $J_i = \langle J_{i+1}, q_i \rangle$ , where  $q_0 = x_3^{a_{13}}x_4^{a_{14}}$  and  $q_1 = x_2^{a_{42}}x_3^{a_{13}}$ . By Proposition 2.4,

$$p(J_i) = p(J_{i+1}) - z^{\deg(q_i)}p(J_{i+1} : \langle q_i \rangle), \text{ for } 0 \leq i \leq 1.$$

Note that  $\deg(q_0) = a_{13} + a_{14}$  and  $\deg(q_1) = a_{13} + a_{42}$ . In this case,  $J_1 : \langle q_0 \rangle = \langle x_2^{a_{42}}, x_3^{a_{23}}, x_4^{a_{34}} \rangle$  and  $J_2 : \langle q_1 \rangle = \langle x_2^{a_{32}}, x_3^{a_{23}}, x_4^{a_4} \rangle$ . We have that  $p(J_2) = \prod_{i=2}^4 (1 - z^{a_i})$  and  $p(J_2 : \langle q_1 \rangle) = (1 -$

$z^{a_{32}}(1 - z^{a_{23}})(1 - z^{a_4})$ , so  $p(J_1) = \prod_{i=2}^4(1 - z^{a_i}) - z^{a_{13}+a_{42}}(1 - z^{a_{32}})(1 - z^{a_{23}})(1 - z^{a_4})$ . Furthermore  $p(J_1 :< q_0 >) = (1 - z^{a_{42}})(1 - z^{a_{23}})(1 - z^{a_{34}})$ , so

$$p(J_0) = \prod_{i=2}^4(1 - z^{a_i}) - z^{a_{13}+a_{42}}(1 - z^{a_{32}})(1 - z^{a_{23}})(1 - z^{a_4}) - z^{a_{13}+a_{14}}(1 - z^{a_{42}})(1 - z^{a_{23}})(1 - z^{a_{34}}).$$

Using the fact that  $1 - z^b = (1 - z) \sum_{i=0}^{b-1} z^i$  we get  $p(J_0) = (1 - z)^3 h(z)$ . Thus  $\text{HS}(R/\text{LM}(I_C^*), z) = \frac{h(z)}{1-z}$ .  $\square$

**Proposition 2.7** *Suppose that  $I_C$  is given as in case 1(b). Let  $a_3 > a_{32} + a_{34}$ ,  $a_{32} < a_{42}$  and  $a_{14} \leq a_{34}$ . Then the set*

$$G = \{f_1 = x_1^{a_1} - x_3^{a_{13}}x_4^{a_{14}}, f_2 = x_2^{a_2} - x_1^{a_{21}}x_3^{a_{23}}, f_3 = x_3^{a_3} - x_2^{a_{32}}x_4^{a_{34}}, \\ f_4 = x_4^{a_4} - x_1^{a_{41}}x_2^{a_{42}}, f_5 = x_1^{a_{21}}x_4^{a_{34}} - x_2^{a_{42}}x_3^{a_{13}}, f_6 = x_3^{a_3+a_{13}} - x_1^{a_1}x_2^{a_{32}}x_4^{a_{34}-a_{14}}\}$$

is a standard basis for  $I_C$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$ .

**Proof.** We will apply the standard basis algorithm [5] to the set  $G = \{f_1, \dots, f_6\}$ . Here  $\text{ecart}(g)$  denotes  $\text{deg}(g) - \text{deg}(\text{LM}(g))$ . We will show that  $\text{NF}(\text{spoly}(f_i, f_j)|G) = 0$ , for all  $i, j$  with  $1 \leq i < j \leq 6$ . Here  $\text{LM}(f_1) = x_3^{a_{13}}x_4^{a_{14}}$ ,  $\text{LM}(f_2) = x_2^{a_2}$ ,  $\text{LM}(f_3) = x_2^{a_{32}}x_4^{a_{34}}$ ,  $\text{LM}(f_4) = x_4^{a_4}$ ,  $\text{LM}(f_5) = x_2^{a_{42}}x_3^{a_{13}}$  and  $\text{LM}(f_6) = x_3^{a_3+a_{13}}$ . Therefore  $\text{NF}(\text{spoly}(f_i, f_j)|G) = 0$  as  $\text{LM}(f_i)$  and  $\text{LM}(f_j)$  are relatively prime, for  $(i, j) \in \{(1, 2), (2, 4), (2, 6), (3, 6), (4, 5), (4, 6)\}$ .

- $\text{spoly}(f_1, f_3) = x_3^{a_3+a_{13}} - x_1^{a_1}x_2^{a_{32}}x_4^{a_{34}-a_{14}} = f_6$ , so  $\text{NF}(\text{spoly}(f_1, f_3)|G) = 0$ .
- $\text{spoly}(f_1, f_4) = x_1^{a_{41}}x_2^{a_{42}}x_3^{a_{13}} - x_1^{a_1}x_4^{a_{34}}$ . It holds that  $a_{34}+a_{21} \geq a_{42}+a_{13}$ , so  $a_{34}+a_{21}+a_{41} \geq a_{41}+a_{42}+a_{13}$  and therefore  $a_1 + a_{34} \geq a_{41} + a_{42} + a_{13}$ . Thus  $\text{LM}(\text{spoly}(f_1, f_4)) = x_1^{a_{41}}x_2^{a_{42}}x_3^{a_{13}}$ . Only  $\text{LM}(f_5)$  divides  $\text{LM}(\text{spoly}(f_1, f_4))$ . Using the fact that  $a_{21} = a_1 - a_{41}$  we get  $\text{ecart}(\text{spoly}(f_1, f_4)) = \text{ecart}(f_5)$ . Since  $\text{spoly}(f_5, \text{spoly}(f_1, f_4)) = 0$ , we get  $\text{NF}(\text{spoly}(f_1, f_4)|G) = 0$ .
- $\text{spoly}(f_1, f_5) = x_1^{a_{21}}x_4^{a_4} - x_1^{a_1}x_2^{a_{42}}$ . Since  $a_4 < a_{41} + a_{42}$ , we get that  $a_4 + a_{21} < a_{21} + a_{41} + a_{42} = a_1 + a_{42}$ . Thus  $\text{LM}(\text{spoly}(f_1, f_5)) = x_1^{a_{21}}x_4^{a_4}$ . Only  $\text{LM}(f_4)$  divides  $\text{LM}(\text{spoly}(f_1, f_5))$  and  $\text{ecart}(\text{spoly}(f_1, f_5)) = \text{ecart}(f_4)$ . Then  $\text{spoly}(f_4, \text{spoly}(f_1, f_5)) = 0$  and also  $\text{NF}(\text{spoly}(f_1, f_5)|G) = 0$ .
- $\text{spoly}(f_1, f_6) = x_1^{a_1}x_2^{a_{32}}x_4^{a_{34}} - x_1^{a_1}x_3^{a_3}$ . Then  $\text{LM}(\text{spoly}(f_1, f_6)) = x_1^{a_1}x_2^{a_{32}}x_4^{a_{34}}$  and only  $\text{LM}(f_3)$  divides  $\text{LM}(\text{spoly}(f_1, f_6))$ . Note that  $\text{ecart}(\text{spoly}(f_1, f_6)) = \text{ecart}(f_3)$ . Then  $\text{spoly}(f_3, \text{spoly}(f_1, f_6)) = 0$  and  $\text{NF}(\text{spoly}(f_1, f_6)|G) = 0$ .
- $\text{spoly}(f_2, f_3) = x_2^{a_{42}}x_3^{a_3} - x_1^{a_{21}}x_3^{a_{23}}x_4^{a_{34}}$ . Since  $a_{42} + a_{13} \leq a_{21} + a_{34}$ , we have  $a_{42} + a_3 \leq a_{21} + a_{34} + a_{23}$ . Thus  $\text{LM}(\text{spoly}(f_2, f_3)) = x_2^{a_{42}}x_3^{a_3}$  and only  $\text{LM}(f_5)$  divides  $\text{LM}(\text{spoly}(f_2, f_3))$ . Using the fact that  $a_3 = a_{13} + a_{23}$  we get  $\text{ecart}(\text{spoly}(f_2, f_3)) = \text{ecart}(f_5)$ . Then  $\text{spoly}(f_5, \text{spoly}(f_2, f_3)) = 0$  and also  $\text{NF}(\text{spoly}(f_2, f_3)|G) = 0$ .
- $\text{spoly}(f_2, f_5) = x_1^{a_{21}}x_2^{a_{32}}x_4^{a_{34}} - x_1^{a_{21}}x_3^{a_3}$ . Since  $a_{32} + a_{34} < a_3$ , we get  $a_{32} + a_{34} + a_{21} < a_3 + a_{21}$ . Thus  $\text{LM}(\text{spoly}(f_2, f_5)) = x_1^{a_{21}}x_2^{a_{32}}x_4^{a_{34}}$  and only  $\text{LM}(f_3)$  divides  $\text{LM}(\text{spoly}(f_2, f_5))$ . Furthermore  $\text{ecart}(\text{spoly}(f_2, f_5)) = \text{ecart}(f_3)$ . Then  $\text{spoly}(f_3, \text{spoly}(f_2, f_5)) = 0$  and also  $\text{NF}(\text{spoly}(f_2, f_5)|G) = 0$ .

- $\text{spoly}(f_3, f_4) = x_1^{a_{41}} x_2^{a_2} - x_3^{a_3} x_4^{a_{14}}$ . Suppose that  $a_{41} + a_2 \leq a_3 + a_{14}$ . Then  $(a_{41} + a_2)n_2 > a_{41}n_1 + a_2n_2$  and also  $(a_{41} + a_2)n_2 > a_3n_3 + a_{14}n_4$ , since  $a_{41}n_1 + a_2n_2 = a_3n_3 + a_{14}n_4$ . But  $a_3n_3 + a_{14}n_4 > (a_3 + a_{14})n_3$ , so  $(a_{41} + a_2)n_2 > (a_3 + a_{14})n_3$ . Moreover  $(a_{41} + a_2)n_2 < (a_{41} + a_2)n_3$  and  $(a_{41} + a_2)n_3 \leq (a_3 + a_{14})n_3$ , hence  $(a_{41} + a_2)n_2 < (a_3 + a_{14})n_3$  a contradiction. Thus  $a_3 + a_{14} < a_{41} + a_2$  and therefore  $\text{LM}(\text{spoly}(f_3, f_4)) = x_3^{a_3} x_4^{a_{14}}$ . Only  $\text{LM}(f_1)$  divides  $\text{LM}(\text{spoly}(f_3, f_4))$  and  $\text{ecart}(\text{spoly}(f_3, f_4)) \leq \text{ecart}(f_1)$ , since  $a_3 = a_{13} + a_{23}$ ,  $a_2 - a_{23} \leq a_{21}$  and  $a_1 = a_{21} + a_{41}$ . Let  $h = \text{spoly}(f_1, \text{spoly}(f_3, f_4)) = x_1^{a_{41}} x_2^{a_2} - x_1^{a_1} x_3^{a_{23}}$ . Since  $a_2 \leq a_{21} + a_{23}$ , we deduce that  $a_2 + a_{41} \leq a_{21} + a_{41} + a_{23} = a_1 + a_{23}$ . Thus  $\text{LM}(h) = x_1^{a_{41}} x_2^{a_2}$  and only  $\text{LM}(f_2)$  divides  $\text{LM}(h)$ . Using the fact that  $a_1 = a_{21} + a_{41}$  we get  $\text{ecart}(h) = \text{ecart}(f_2)$ . Then  $\text{spoly}(f_2, h) = 0$  and  $\text{NF}(\text{spoly}(f_3, f_4)|G) = 0$ .
- $\text{spoly}(f_5, f_6) = x_1^{a_1} x_2^{a_2} x_4^{a_{34} - a_{14}} - x_1^{a_{21}} x_3^{a_3} x_4^{a_{34}}$ . Recall that  $a_3 + a_{14} < a_{41} + a_2$ . Then  $a_1 + a_2 + a_{34} - a_{14} = a_{21} + a_{41} + a_2 + a_{34} - a_{14} > a_{21} + a_3 + a_{14} + a_{34} - a_{14} = a_{21} + a_3 + a_{34}$  and therefore  $\text{LM}(\text{spoly}(f_5, f_6)) = x_1^{a_{21}} x_3^{a_3} x_4^{a_{34}}$ . Only  $\text{LM}(f_1)$  divides  $\text{LM}(\text{spoly}(f_5, f_6))$  and  $\text{ecart}(\text{spoly}(f_5, f_6)) \leq \text{ecart}(f_1)$ , since  $a_3 = a_{13} + a_{23}$  and  $a_2 - a_{21} - a_{23} \leq 0$ . Let  $g = \text{spoly}(f_1, \text{spoly}(f_5, f_6)) = x_1^{a_1 + a_{21}} x_3^{a_{23}} x_4^{a_{34} - a_{14}} - x_1^{a_1} x_2^{a_2} x_4^{a_{34} - a_{14}}$ . Then  $\text{LM}(g) = x_1^{a_1} x_2^{a_2} x_4^{a_{34} - a_{14}}$ . Only  $\text{LM}(f_2)$  divides  $\text{LM}(g)$  and  $\text{ecart}(g) = \text{ecart}(f_2)$ . Then  $\text{spoly}(f_2, g) = 0$  and  $\text{NF}(\text{spoly}(f_5, f_6)|G) = 0$ .
- $\text{spoly}(f_3, f_5) = x_2^{a_{42} - a_{32}} x_3^{a_3 + a_{13}} - x_1^{a_{21}} x_4^{2a_{34}}$ . We distinguish the following cases:
  1.  $\text{LM}(\text{spoly}(f_3, f_5)) = x_1^{a_{21}} x_4^{2a_{34}}$ . Only  $\text{LM}(f_4)$  divides  $\text{LM}(\text{spoly}(f_3, f_5))$  and  $\text{ecart}(\text{spoly}(f_3, f_5)) \leq \text{ecart}(f_4)$ , since  $a_3 + a_{13} \leq a_1 + a_{32} + a_{34} - a_{14}$ ,  $a_1 - a_{21} = a_{41}$  and  $a_{14} + a_{34} = a_4$ . Let  $g = \text{spoly}(f_4, \text{spoly}(f_3, f_5)) = x_1^{a_1} x_2^{a_{42}} x_4^{a_{34} - a_{14}} - x_2^{a_{42} - a_{32}} x_3^{a_3 + a_{13}}$ . Then  $a_3 + a_{13} + a_{42} - a_{32} \leq a_1 + a_{42} + a_{34} - a_{14}$  and therefore  $\text{LM}(g) = x_2^{a_{42} - a_{32}} x_3^{a_3 + a_{13}}$ . Only  $\text{LM}(f_6)$  divides  $\text{LM}(g)$  and also  $\text{ecart}(g) = \text{ecart}(f_6)$ . Finally  $\text{spoly}(f_6, g) = 0$  and  $\text{NF}(\text{spoly}(f_3, f_5)|G) = 0$ .
  2.  $\text{LM}(\text{spoly}(f_3, f_5)) = x_2^{a_{42} - a_{32}} x_3^{a_3 + a_{13}}$ . Only  $\text{LM}(f_6)$  divides the monomial  $\text{LM}(\text{spoly}(f_3, f_5))$  and  $\text{ecart}(\text{spoly}(f_3, f_5)) \leq \text{ecart}(f_6)$ . Let  $h = \text{spoly}(f_6, \text{spoly}(f_3, f_5)) = x_1^{a_1} x_2^{a_{42}} x_4^{a_{34} - a_{14}} - x_1^{a_{21}} x_4^{2a_{34}}$ . Then  $a_1 + a_{42} + a_{34} - a_{14} \geq a_{21} + 2a_{34}$ , since  $a_1 = a_{21} + a_{41}$ ,  $a_{41} + a_{42} \geq a_4$  and  $a_4 - a_{14} = a_{34}$ . We have that  $a_1 + a_{42} + a_{34} - a_{14} = a_{21} + a_{41} + a_{42} + a_{34} - a_{14} \geq a_{21} + a_4 + a_{34} - a_{14} = a_{21} + a_{14} + a_{34} + a_{34} - a_{14} = a_{21} + 2a_{34}$ . So  $\text{LM}(h) = x_1^{a_{21}} x_4^{2a_{34}}$ . Only  $\text{LM}(f_4)$  divides  $\text{LM}(h)$  and also  $\text{ecart}(h) = \text{ecart}(f_4)$ . Then  $\text{spoly}(f_4, h) = 0$  and  $\text{NF}(\text{spoly}(f_3, f_5)|G) = 0$ .  $\square$

**Theorem 2.8** Suppose that  $I_C$  is given as in case 1(b). Let  $a_3 > a_{32} + a_{34}$ ,  $a_{32} < a_{42}$  and  $a_{14} \leq a_{34}$ . Then the reduced Hilbert series of  $R/I_C^*$  is  $\text{HS}(R/I_C^*, z) = \frac{h(z)}{1-z}$  for

$$\begin{aligned}
 h(z) = & \sum_{i=0}^{a_2-1} z^i \sum_{i=0}^{a_3+a_{13}-1} z^i \sum_{i=0}^{a_4-1} z^i - z^{a_{13}+a_{14}} \sum_{i=0}^{a_2-1} z^i \sum_{i=0}^{a_3-1} z^i \sum_{i=0}^{a_{34}-1} z^i - \\
 & z^{a_{13}+a_{42}} \sum_{i=0}^{a_{32}-1} z^i \sum_{i=0}^{a_3-1} z^i \sum_{i=0}^{a_{14}-1} z^i - z^{a_{32}+a_{34}} \sum_{i=0}^{a_{42}-1} z^i \sum_{i=0}^{a_{13}-1} z^i \sum_{i=0}^{a_{14}-1} z^i.
 \end{aligned}$$

**Proof.** By Proposition 2.7,  $G = \{x_1^{a_1} - x_3^{a_{13}} x_4^{a_{14}}, x_2^{a_2} - x_1^{a_{21}} x_3^{a_{23}}, x_3^{a_3} - x_2^{a_{32}} x_4^{a_{34}}, x_4^{a_4} - x_1^{a_{41}} x_2^{a_{42}}, x_1^{a_{21}} x_4^{a_{34}} - x_2^{a_{42}} x_3^{a_{13}}, x_3^{a_3 + a_{13}} - x_1^{a_1} x_2^{a_{32}} x_4^{a_{34} - a_{14}}\}$  is a standard basis for  $I_C$  with respect to the negative degree reverse

lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$ . Then  $I_C^*$  is generated by the least homogeneous summands of the elements in  $G$ . Moreover,  $\text{LM}(I_C^*) = \langle x_3^{a_{13}} x_4^{a_{14}}, x_2^{a_2}, x_3^{a_3+a_{13}}, x_4^{a_4}, x_2^{a_{32}} x_4^{a_{34}}, x_2^{a_{42}} x_3^{a_{13}} \rangle$ . Let

$$J_0 = \text{LM}(I_C^*), J_1 = \langle x_3^{a_{13}} x_4^{a_{14}}, x_2^{a_2}, x_3^{a_3+a_{13}}, x_4^{a_4}, x_2^{a_{42}} x_3^{a_{13}} \rangle,$$

$$J_2 = \langle x_3^{a_{13}} x_4^{a_{14}}, x_2^{a_2}, x_3^{a_3+a_{13}}, x_4^{a_4} \rangle, J_3 = \langle x_2^{a_2}, x_3^{a_3+a_{13}}, x_4^{a_4} \rangle.$$

Then  $J_i = \langle J_{i+1}, q_i \rangle$ , where  $q_0 = x_2^{a_{32}} x_4^{a_{34}}$ ,  $q_1 = x_2^{a_{42}} x_3^{a_{13}}$  and  $q_2 = x_3^{a_{13}} x_4^{a_{14}}$ . Therefore

$$p(J_i) = p(J_{i+1}) - z^{\deg(q_i)} p(J_{i+1} : \langle q_i \rangle), \text{ for } 0 \leq i \leq 2.$$

In this case,  $J_1 : \langle q_0 \rangle = \langle x_2^{a_{42}}, x_3^{a_{13}}, x_4^{a_{14}} \rangle$ ,  $J_2 : \langle q_1 \rangle = \langle x_2^{a_{32}}, x_3^{a_3}, x_4^{a_{14}} \rangle$  and  $J_3 : \langle q_2 \rangle = \langle x_2^{a_2}, x_3^{a_3}, x_4^{a_{34}} \rangle$ . We have that  $p(J_1 : \langle q_0 \rangle) = (1 - z^{a_{42}})(1 - z^{a_{13}})(1 - z^{a_{14}})$ ,  $p(J_2 : \langle q_1 \rangle) = (1 - z^{a_{32}})(1 - z^{a_3})(1 - z^{a_{14}})$ ,  $p(J_3 : \langle q_2 \rangle) = (1 - z^{a_2})(1 - z^{a_3})(1 - z^{a_{34}})$  and  $p(J_3) = (1 - z^{a_2})(1 - z^{a_3+a_{13}})(1 - z^{a_4})$ . Therefore

$$p(J_0) = (1 - z^{a_2})(1 - z^{a_3+a_{13}})(1 - z^{a_4}) - z^{a_{13}+a_{14}}(1 - z^{a_2})(1 - z^{a_3})(1 - z^{a_{34}}) - z^{a_{13}+a_{42}}(1 - z^{a_{32}})(1 - z^{a_3})(1 - z^{a_{14}}) - z^{a_{32}+a_{34}}(1 - z^{a_{42}})(1 - z^{a_{13}})(1 - z^{a_{14}}).$$

One can easily show that  $p(J_0) = (1 - z)^3 h(z)$ . Thus  $\text{HS}(R/\text{LM}(I_C^*), z) = \frac{h(z)}{1-z}$ . □

**Proposition 2.9** *Suppose that  $I_C$  is given as in case 1(b) and also that  $a_3 > a_{32} + a_{34}$ . Assume that  $a_{42} \leq a_{32}$ . (1) If  $a_{34} < a_{14}$ , then*

$$G = \{f_1 = x_1^{a_1} - x_3^{a_{13}} x_4^{a_{14}}, f_2 = x_2^{a_2} - x_1^{a_{21}} x_3^{a_{23}}, f_3 = x_3^{a_3} - x_2^{a_{32}} x_4^{a_{34}}, f_4 = x_4^{a_4} - x_1^{a_{41}} x_2^{a_{42}}, f_5 = x_1^{a_{21}} x_4^{a_{34}} - x_2^{a_{42}} x_3^{a_{13}}, f_6 = x_3^{a_3+a_{13}} - x_1^{a_{21}} x_2^{a_{32}-a_{42}} x_4^{2a_{34}}\}$$

is a standard basis for  $I_C$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$ .

(2) If  $a_{14} \leq a_{34}$ , then

$$G = \{f_1 = x_1^{a_1} - x_3^{a_{13}} x_4^{a_{14}}, f_2 = x_2^{a_2} - x_1^{a_{21}} x_3^{a_{23}}, f_3 = x_3^{a_3} - x_2^{a_{32}} x_4^{a_{34}}, f_4 = x_4^{a_4} - x_1^{a_{41}} x_2^{a_{42}}, f_5 = x_1^{a_{21}} x_4^{a_{34}} - x_2^{a_{42}} x_3^{a_{13}}, f_6 = x_3^{a_3+a_{13}} - x_1^{a_1} x_2^{a_{32}} x_4^{a_{34}-a_{14}}\}$$

is a standard basis for  $I_C$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$ .

**Proof.** (1) We will apply the standard basis algorithm to the set  $G = \{f_1, \dots, f_6\}$ . We will show that  $\text{NF}(\text{spoly}(f_i, f_j)|G) = 0$ , for all  $i, j$  with  $1 \leq i < j \leq 6$ . Here  $\text{LM}(f_1) = x_3^{a_{13}} x_4^{a_{14}}$ ,  $\text{LM}(f_2) = x_2^{a_2}$ ,  $\text{LM}(f_3) = x_2^{a_{32}} x_4^{a_{34}}$ ,  $\text{LM}(f_4) = x_4^{a_4}$ ,  $\text{LM}(f_5) = x_2^{a_{42}} x_3^{a_{13}}$  and  $\text{LM}(f_6) = x_3^{a_3+a_{13}}$ . Therefore  $\text{NF}(\text{spoly}(f_i, f_j)|G) = 0$  as  $\text{LM}(f_i)$  and  $\text{LM}(f_j)$  are relatively prime, for  $(i, j) \in \{(1, 2), (2, 4), (2, 6), (3, 6), (4, 5), (4, 6)\}$ .

- $\text{spoly}(f_1, f_3) = x_3^{a_3+a_{13}} x_4^{a_{14}-a_{34}} - x_1^{a_1} x_2^{a_{32}}$ . Then  $\text{LM}(\text{spoly}(f_1, f_3)) = x_3^{a_3+a_{13}} x_4^{a_{14}-a_{34}}$ . Only  $\text{LM}(f_6)$  divides  $\text{LM}(\text{spoly}(f_1, f_3))$  and also  $\text{ecart}(f_6) \leq \text{ecart}(\text{spoly}(f_1, f_3))$ , since  $a_1 = a_{21} + a_{41}$ ,  $a_4 - a_{42} \leq a_{41}$  and  $a_4 = a_{14} + a_{34}$ . Let  $g = \text{spoly}(f_6, \text{spoly}(f_1, f_3)) = x_1^{a_{21}} x_2^{a_{32}-a_{42}} x_4^{a_4} - x_1^{a_1} x_2^{a_{32}}$ . Only  $\text{LM}(f_4)$  divides  $\text{LM}(g) = x_1^{a_{21}} x_2^{a_{32}-a_{42}} x_4^{a_4}$  and also  $\text{ecart}(g) = \text{ecart}(f_4)$ . Furthermore  $\text{spoly}(f_4, g) = 0$  and  $\text{NF}(\text{spoly}(f_1, f_3)|G) = 0$ .



- $\text{spoly}(f_1, f_4) = x_1^{a_{41}} x_2^{a_{42}} x_3^{a_{13}} - x_1^{a_1} x_4^{a_{34}}$ . In this case  $\text{LM}(\text{spoly}(f_1, f_4)) = x_1^{a_{41}} x_2^{a_{42}} x_3^{a_{13}}$ . Only  $\text{LM}(f_5)$  divides  $\text{LM}(\text{spoly}(f_1, f_4))$ . Using the fact that  $a_1 = a_{21} + a_{41}$  we get  $\text{ecart}(\text{spoly}(f_1, f_4)) = \text{ecart}(f_5)$ . The computation  $\text{spoly}(f_5, \text{spoly}(f_1, f_4)) = 0$  implies that  $\text{NF}(\text{spoly}(f_1, f_4)|G) = 0$ .
- $\text{spoly}(f_1, f_5) = x_1^{a_{21}} x_4^{a_4} - x_1^{a_1} x_2^{a_{42}}$ . Only  $\text{LM}(f_4)$  divides  $\text{LM}(\text{spoly}(f_1, f_5)) = x_1^{a_{21}} x_4^{a_4}$  and  $\text{ecart}(\text{spoly}(f_1, f_5)) = \text{ecart}(f_4)$ . Then  $\text{spoly}(f_4, \text{spoly}(f_1, f_5)) = 0$  and also  $\text{NF}(\text{spoly}(f_1, f_5)|G) = 0$ .
- $\text{spoly}(f_1, f_6) = x_1^{a_{21}} x_2^{a_{32}-a_{42}} x_4^{a_{14}+2a_{34}} - x_1^{a_1} x_3^{a_3}$ . We have that  $a_{21} + a_{32} - a_{42} + a_{14} + 2a_{34} < a_1 + a_3$ , since  $a_{14} + 2a_{34} = a_{34} + a_4$ ,  $a_{32} + a_{34} < a_3$  and  $a_4 - a_{42} \leq a_{41}$ . So  $\text{LM}(\text{spoly}(f_1, f_6)) = x_1^{a_{21}} x_2^{a_{32}-a_{42}} x_4^{a_{14}+2a_{34}}$ . Only  $\text{LM}(f_4)$  divides  $\text{LM}(\text{spoly}(f_1, f_6))$  and  $\text{ecart}(\text{spoly}(f_1, f_6)) > \text{ecart}(f_4)$ , since  $a_3 - a_{32} - a_{34} > 0$ . Let  $h = \text{spoly}(f_4, \text{spoly}(f_1, f_6)) = x_1^{a_1} x_3^{a_3} - x_1^{a_1} x_2^{a_{32}} x_4^{a_{34}}$ , then  $\text{LM}(h) = x_1^{a_1} x_2^{a_{32}} x_4^{a_{34}}$ . Only  $\text{LM}(f_3)$  divides  $\text{LM}(h)$  and  $\text{ecart}(h) = \text{ecart}(f_3)$ . Then  $\text{spoly}(f_3, h) = 0$  and  $\text{NF}(\text{spoly}(f_1, f_6)|G) = 0$ .
- $\text{spoly}(f_2, f_3) = x_2^{a_{42}} x_3^{a_3} - x_1^{a_{21}} x_3^{a_{23}} x_4^{a_{34}}$ . Then  $\text{LM}(\text{spoly}(f_2, f_3)) = x_2^{a_{42}} x_3^{a_3}$  and only  $\text{LM}(f_5)$  divides  $\text{LM}(\text{spoly}(f_2, f_3))$ . Furthermore  $\text{ecart}(\text{spoly}(f_2, f_3)) = \text{ecart}(f_5)$ . Then  $\text{spoly}(f_5, \text{spoly}(f_2, f_3)) = 0$  and also  $\text{NF}(\text{spoly}(f_2, f_3)|G) = 0$ .
- $\text{spoly}(f_2, f_5) = x_1^{a_{21}} x_2^{a_{32}} x_4^{a_{34}} - x_1^{a_{21}} x_3^{a_3}$ . Thus  $\text{LM}(\text{spoly}(f_2, f_5)) = x_1^{a_{21}} x_2^{a_{32}} x_4^{a_{34}}$  and only  $\text{LM}(f_3)$  divides  $\text{LM}(\text{spoly}(f_2, f_5))$ . Furthermore  $\text{ecart}(\text{spoly}(f_2, f_5)) = \text{ecart}(f_3)$ . Then  $\text{spoly}(f_3, \text{spoly}(f_2, f_5)) = 0$  and  $\text{NF}(\text{spoly}(f_2, f_5)|G) = 0$ .
- $\text{spoly}(f_3, f_4) = x_1^{a_{41}} x_2^{a_2} - x_3^{a_3} x_4^{a_{14}}$ . Since  $a_3 + a_{14} < a_{41} + a_2$ , we have that  $\text{LM}(\text{spoly}(f_3, f_4)) = x_3^{a_3} x_4^{a_{14}}$ . Only  $\text{LM}(f_1)$  divides  $\text{LM}(\text{spoly}(f_3, f_4))$  and  $\text{ecart}(\text{spoly}(f_3, f_4)) \leq \text{ecart}(f_1)$ . Let  $g = \text{spoly}(f_1, \text{spoly}(f_3, f_4)) = x_1^{a_{41}} x_2^{a_2} - x_1^{a_1} x_3^{a_{23}}$ . Then  $\text{LM}(g) = x_1^{a_{41}} x_2^{a_2}$  and only  $\text{LM}(f_2)$  divides  $\text{LM}(g)$ . Also  $\text{ecart}(g) = \text{ecart}(f_2)$ . Then  $\text{spoly}(f_2, g) = 0$  and  $\text{NF}(\text{spoly}(f_3, f_4)|G) = 0$ .
- $\text{spoly}(f_3, f_5) = x_1^{a_{21}} x_2^{a_{32}-a_{42}} x_4^{2a_{34}} - x_3^{a_3+a_{13}} = -f_6$ . Then  $\text{NF}(\text{spoly}(f_3, f_5)|G) = 0$ .
- $\text{spoly}(f_5, f_6) = x_1^{a_{21}} x_2^{a_{32}} x_4^{2a_{34}} - x_1^{a_{21}} x_3^{a_3} x_4^{a_{34}}$ . Then  $\text{LM}(\text{spoly}(f_5, f_6)) = x_1^{a_{21}} x_2^{a_{32}} x_4^{2a_{34}}$ . Only  $\text{LM}(f_3)$  divides  $\text{LM}(\text{spoly}(f_5, f_6))$  and  $\text{ecart}(\text{spoly}(f_5, f_6)) = \text{ecart}(f_3)$ . We have that  $\text{spoly}(f_3, \text{spoly}(f_5, f_6)) = 0$  and  $\text{NF}(\text{spoly}(f_5, f_6)|G) = 0$ .

(2) It is enough to prove that  $\text{NF}(\text{spoly}(f_i, f_j)|G) = 0$ , for all  $i, j$  with  $1 \leq i < j \leq 6$ . Here  $\text{LM}(f_1) = x_3^{a_{13}} x_4^{a_{14}}$ ,  $\text{LM}(f_2) = x_2^{a_2}$ ,  $\text{LM}(f_3) = x_2^{a_{32}} x_4^{a_{34}}$ ,  $\text{LM}(f_4) = x_4^{a_4}$ ,  $\text{LM}(f_5) = x_2^{a_{42}} x_3^{a_{13}}$  and  $\text{LM}(f_6) = x_3^{a_3+a_{13}}$ . So  $\text{NF}(\text{spoly}(f_i, f_j)|G) = 0$  as  $\text{LM}(f_i)$  and  $\text{LM}(f_j)$  are relatively prime, for

$$(i, j) \in \{(1, 2), (2, 4), (2, 6), (3, 6), (4, 5), (4, 6)\}.$$

- $\text{spoly}(f_1, f_3) = x_3^{a_3+a_{13}} - x_1^{a_1} x_2^{a_{32}} x_4^{a_{34}-a_{14}} = f_6$ . Then  $\text{NF}(\text{spoly}(f_1, f_3)|G) = 0$ .
- $\text{spoly}(f_1, f_4) = x_1^{a_{41}} x_2^{a_{42}} x_3^{a_{13}} - x_1^{a_1} x_4^{a_{34}}$ . Then  $\text{LM}(\text{spoly}(f_1, f_4)) = x_1^{a_{41}} x_2^{a_{42}} x_3^{a_{13}}$  and only  $\text{LM}(f_5)$  divides  $\text{LM}(\text{spoly}(f_1, f_4))$ . Also  $\text{ecart}(\text{spoly}(f_1, f_4)) = \text{ecart}(f_5)$ . Moreover  $\text{spoly}(f_5, \text{spoly}(f_1, f_4)) = 0$  and  $\text{NF}(\text{spoly}(f_1, f_4)|G) = 0$ .
- $\text{spoly}(f_1, f_5) = x_1^{a_{21}} x_4^{a_4} - x_1^{a_1} x_2^{a_{42}}$ . Then  $\text{LM}(\text{spoly}(f_1, f_5)) = x_1^{a_{21}} x_4^{a_4}$  and only  $\text{LM}(f_4)$  divides  $\text{LM}(\text{spoly}(f_1, f_5))$ . Also  $\text{ecart}(\text{spoly}(f_1, f_5)) = \text{ecart}(f_4)$ . Then  $\text{spoly}(f_4, \text{spoly}(f_1, f_5)) = 0$  and  $\text{NF}(\text{spoly}(f_1, f_5)|G) = 0$ .

- $\text{spoly}(f_1, f_6) = x_1^{a_1} x_2^{a_{32}} x_4^{a_{34}} - x_1^{a_1} x_3^{a_3}$ . Then  $\text{LM}(\text{spoly}(f_1, f_6)) = x_1^{a_1} x_2^{a_{32}} x_4^{a_{34}}$  and only  $\text{LM}(f_3)$  divides  $\text{LM}(\text{spoly}(f_1, f_6))$ . Moreover  $\text{ecart}(\text{spoly}(f_1, f_6)) = \text{ecart}(f_3)$ . Then  $\text{spoly}(f_3, \text{spoly}(f_1, f_6)) = 0$  and  $\text{NF}(\text{spoly}(f_1, f_6)|G) = 0$ .
- $\text{spoly}(f_2, f_3) = x_2^{a_{42}} x_3^{a_3} - x_1^{a_{21}} x_3^{a_{23}} x_4^{a_{34}}$ . Then  $\text{LM}(\text{spoly}(f_2, f_3)) = x_2^{a_{42}} x_3^{a_3}$  and only  $\text{LM}(f_5)$  divides  $\text{LM}(\text{spoly}(f_2, f_3))$ . Also  $\text{ecart}(\text{spoly}(f_2, f_3)) = \text{ecart}(f_5)$ . Then  $\text{spoly}(f_5, \text{spoly}(f_2, f_3)) = 0$  and also  $\text{NF}(\text{spoly}(f_2, f_3)|G) = 0$ .
- $\text{spoly}(f_2, f_5) = x_1^{a_{21}} x_2^{a_{32}} x_4^{a_{34}} - x_1^{a_{21}} x_3^{a_3}$ . Then  $\text{LM}(\text{spoly}(f_2, f_5)) = x_1^{a_{21}} x_2^{a_{32}} x_4^{a_{34}}$  and only  $\text{LM}(f_3)$  divides  $\text{LM}(\text{spoly}(f_2, f_5))$ . Furthermore  $\text{ecart}(\text{spoly}(f_2, f_5)) = \text{ecart}(f_3)$ . Then  $\text{spoly}(f_3, \text{spoly}(f_2, f_5)) = 0$  and  $\text{NF}(\text{spoly}(f_2, f_5)|G) = 0$ .
- $\text{spoly}(f_3, f_4) = x_1^{a_{41}} x_2^{a_2} - x_3^{a_3} x_4^{a_{14}}$ . Then  $\text{LM}(\text{spoly}(f_3, f_4)) = x_3^{a_3} x_4^{a_{14}}$  and only  $\text{LM}(f_1)$  divides  $\text{LM}(\text{spoly}(f_3, f_4))$ . Also  $\text{ecart}(\text{spoly}(f_3, f_4)) \leq \text{ecart}(f_1)$ , since  $a_2 \leq a_{21} + a_{23}$ ,  $a_1 = a_{21} + a_{41}$  and  $a_3 = a_{13} + a_{23}$ . Let  $h = \text{spoly}(f_1, \text{spoly}(f_3, f_4)) = x_1^{a_{41}} x_2^{a_2} - x_1^{a_1} x_3^{a_{23}}$ , then  $\text{LM}(h) = x_1^{a_{41}} x_2^{a_2}$  and only  $\text{LM}(f_2)$  divides  $\text{LM}(h)$ . Moreover  $\text{ecart}(h) = \text{ecart}(f_2)$ . Then  $\text{spoly}(f_2, h) = 0$  and also  $\text{NF}(\text{spoly}(f_3, f_4)|G) = 0$ .
- $\text{spoly}(f_5, f_6) = x_1^{a_1} x_2^{a_2} x_4^{a_{34}-a_{14}} - x_1^{a_{21}} x_3^{a_3} x_4^{a_{34}}$ . Then  $\text{LM}(\text{spoly}(f_5, f_6)) = x_1^{a_{21}} x_3^{a_3} x_4^{a_{34}}$  and only  $\text{LM}(f_1)$  divides  $\text{LM}(\text{spoly}(f_5, f_6))$ . Furthermore  $\text{ecart}(\text{spoly}(f_5, f_6)) \leq \text{ecart}(f_1)$ , since  $a_2 \leq a_{21} + a_{23}$  and  $a_3 = a_{13} + a_{23}$ . Let  $g = \text{spoly}(f_1, \text{spoly}(f_5, f_6)) = x_1^{a_1} x_2^{a_2} x_4^{a_{34}-a_{14}} - x_1^{a_1+a_{21}} x_3^{a_{23}} x_4^{a_{34}-a_{14}}$ . Then  $\text{LM}(g) = x_1^{a_1} x_2^{a_2} x_4^{a_{34}-a_{14}}$  and only  $\text{LM}(f_2)$  divides  $\text{LM}(g)$ . Also  $\text{ecart}(g) = \text{ecart}(f_2)$ . Then  $\text{spoly}(f_2, g) = 0$  and  $\text{NF}(\text{spoly}(f_5, f_6)|G) = 0$ .
- $\text{spoly}(f_3, f_5) = x_1^{a_{21}} x_2^{a_{32}-a_{42}} x_4^{2a_{34}} - x_3^{a_3+a_{13}}$ . We distinguish the following cases:  
 (1)  $\text{LM}(\text{spoly}(f_3, f_5)) = x_3^{a_3+a_{13}}$ , then only  $\text{LM}(f_6)$  divides  $\text{LM}(\text{spoly}(f_3, f_5))$  and also  $\text{ecart}(\text{spoly}(f_3, f_5)) \leq \text{ecart}(f_6)$ , since  $a_{34} = a_4 - a_{14}$ ,  $a_4 - a_{42} \leq a_{41}$  and  $a_1 = a_{21} + a_{41}$ . Let  $g = \text{spoly}(f_6, \text{spoly}(f_3, f_5)) = x_1^{a_{21}} x_2^{a_{32}-a_{42}} x_4^{2a_{34}} - x_1^{a_1} x_2^{a_{32}} x_4^{a_{34}-a_{14}}$ . Then  $\text{LM}(g) = x_1^{a_{21}} x_2^{a_{32}-a_{42}} x_4^{2a_{34}}$  and only  $\text{LM}(f_4)$  divides  $\text{LM}(g)$ . Also  $\text{ecart}(g) = \text{ecart}(f_4)$ . We have that  $\text{spoly}(f_4, g) = 0$  and  $\text{NF}(\text{spoly}(f_3, f_5)|G) = 0$ .  
 (2)  $\text{LM}(\text{spoly}(f_3, f_5)) = x_1^{a_{21}} x_2^{a_{32}-a_{42}} x_4^{2a_{34}}$ . Only  $\text{LM}(f_4)$  divides  $\text{LM}(\text{spoly}(f_3, f_5))$  and also

$$\text{ecart}(\text{spoly}(f_3, f_5)) < \text{ecart}(f_4),$$

since  $a_3 + a_{13} \leq a_1 + a_{32} + a_{34} - a_{14}$ ,  $a_1 = a_{21} + a_{41}$  and  $a_4 = a_{14} + a_{34}$ . Then  $\text{spoly}(f_4, \text{spoly}(f_3, f_5)) = x_3^{a_3+a_{13}} - x_1^{a_1} x_2^{a_{32}} x_4^{a_{34}-a_{14}} = f_6$ . So  $\text{NF}(\text{spoly}(f_3, f_5)|G) = 0$ . □

**Theorem 2.10** Suppose that  $I_C$  is given as in case 1(b) and also that  $a_3 > a_{32} + a_{34}$ . Assume that  $a_{42} \leq a_{32}$ .

Then the reduced Hilbert series of  $R/I_C^*$  is  $\text{HS}(R/I_C^*, z) = \frac{h(z)}{1-z}$  for

$$h(z) = \sum_{i=0}^{a_2-1} z^i \sum_{i=0}^{a_3+a_{13}-1} z^i \sum_{i=0}^{a_4-1} z^i - z^{a_{13}+a_{42}} \sum_{i=0}^{a_{32}-1} z^i \sum_{i=0}^{a_3-1} z^i \sum_{i=0}^{a_4-1} z^i - z^{a_{32}+a_{34}} \sum_{i=0}^{a_{42}-1} z^i \sum_{i=0}^{a_{13}-1} z^i \sum_{i=0}^{a_{14}-1} z^i - z^{a_{13}+a_{14}} \sum_{i=0}^{a_{42}-1} z^i \sum_{i=0}^{a_3-1} z^i \sum_{i=0}^{a_{34}-1} z^i.$$

**Proof.** By Proposition 2.9,  $\text{LM}(I_C^*)$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$  is written as  $\text{LM}(I_C^*) = \langle x_3^{a_{13}} x_4^{a_{14}}, x_2^{a_2}, x_3^{a_3+a_{13}}, x_4^{a_4}, x_2^{a_{42}} x_3^{a_{13}}, x_2^{a_{32}} x_4^{a_{34}} \rangle$ . Let

$$J_0 = \text{LM}(I_C^*), J_1 = \langle x_2^{a_2}, x_3^{a_3+a_{13}}, x_4^{a_4}, x_2^{a_{32}} x_4^{a_{34}}, x_2^{a_{42}} x_3^{a_{13}} \rangle,$$

$$J_2 = \langle x_2^{a_2}, x_3^{a_3+a_{13}}, x_4^{a_4}, x_2^{a_{42}} x_3^{a_{13}} \rangle, J_3 = \langle x_2^{a_2}, x_3^{a_3+a_{13}}, x_4^{a_4} \rangle.$$

Then  $J_i = \langle J_{i+1}, q_i \rangle$ , where  $q_0 = x_3^{a_{13}} x_4^{a_{14}}$ ,  $q_1 = x_2^{a_{32}} x_4^{a_{34}}$  and  $q_2 = x_2^{a_{42}} x_3^{a_{13}}$ . So

$$p(J_i) = p(J_{i+1}) - z^{\deg(q_i)} p(J_{i+1} : \langle q_i \rangle), \text{ for } 0 \leq i \leq 2.$$

In this case,  $J_1 : \langle q_0 \rangle = \langle x_2^{a_{42}}, x_3^{a_3}, x_4^{a_{34}} \rangle$ ,  $J_2 : \langle q_1 \rangle = \langle x_2^{a_{42}}, x_3^{a_{13}}, x_4^{a_{14}} \rangle$  and  $J_3 : \langle q_2 \rangle = \langle x_2^{a_{32}}, x_3^{a_3}, x_4^{a_4} \rangle$ . We have that  $p(J_1 : \langle q_0 \rangle) = (1 - z^{a_{42}})(1 - z^{a_3})(1 - z^{a_{34}})$ ,  $p(J_2 : \langle q_1 \rangle) = (1 - z^{a_{42}})(1 - z^{a_{13}})(1 - z^{a_{14}})$ ,  $p(J_3 : \langle q_2 \rangle) = (1 - z^{a_{32}})(1 - z^{a_3})(1 - z^{a_4})$  and  $p(J_3) = (1 - z^{a_2})(1 - z^{a_3+a_{13}})(1 - z^{a_4})$ . So

$$p(J_0) = (1 - z^{a_2})(1 - z^{a_3+a_{13}})(1 - z^{a_4}) - z^{a_{13}+a_{42}}(1 - z^{a_{32}})(1 - z^{a_3})(1 - z^{a_4}) - z^{a_{32}+a_{34}}(1 - z^{a_{42}})(1 - z^{a_{13}})(1 - z^{a_{14}}) - z^{a_{13}+a_{14}}(1 - z^{a_{42}})(1 - z^{a_3})(1 - z^{a_{34}}).$$

Thus  $p(J_0) = (1 - z)^3 h(z)$ , so  $\text{HS}(R/\text{LM}(I_C^*), z) = \frac{h(z)}{1-z}$ . □

The proof of the next proposition is similar to that of Proposition 2.7 and therefore it is omitted.

**Proposition 2.11** Suppose that  $I_C$  is given as in case 2(a). If  $a_{24} < a_{34}$  and  $a_{13} \leq a_{23}$ , then

$$G = \{x_1^{a_1} - x_2^{a_{12}} x_3^{a_{13}}, x_2^{a_2} - x_3^{a_{23}} x_4^{a_{24}}, x_3^{a_3} - x_1^{a_{31}} x_4^{a_{34}}, x_4^{a_4} - x_1^{a_{41}} x_2^{a_{42}}, x_1^{a_{41}} x_3^{a_{23}} - x_2^{a_{12}} x_4^{a_{34}}, x_2^{a_2+a_{12}} - x_1^{a_1} x_3^{a_{23}-a_{13}} x_4^{a_{24}}\}$$

is a standard basis for  $I_C$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$ .

**Theorem 2.12** Suppose that  $I_C$  is given as in case 2(a). If  $a_{24} < a_{34}$  and  $a_{13} \leq a_{23}$ , then the reduced Hilbert series of  $R/I_C^*$  is  $\text{HS}(R/I_C^*, z) = \frac{h(z)}{1-z}$  for

$$h(z) = \sum_{i=0}^{a_2+a_{12}-1} z^i \sum_{i=0}^{a_3-1} z^i \sum_{i=0}^{a_4-1} z^i - z^{a_{12}+a_{13}} \sum_{i=0}^{a_2-1} z^i \sum_{i=0}^{a_{23}-1} z^i \sum_{i=0}^{a_4-1} z^i - z^{a_{12}+a_{34}} \sum_{i=0}^{a_2-1} z^i \sum_{i=0}^{a_{13}-1} z^i \sum_{i=0}^{a_{34}-1} z^i - z^{a_{23}+a_{24}} \sum_{i=0}^{a_{12}-1} z^i \sum_{i=0}^{a_{13}-1} z^i \sum_{i=0}^{a_4-1} z^i.$$

**Proof.** By Proposition 2.11,  $G = \{x_1^{a_1} - x_2^{a_{12}} x_3^{a_{13}}, x_2^{a_2} - x_3^{a_{23}} x_4^{a_{24}}, x_3^{a_3} - x_1^{a_{31}} x_4^{a_{34}}, x_4^{a_4} - x_1^{a_{41}} x_2^{a_{42}}, x_1^{a_{41}} x_3^{a_{23}} - x_2^{a_{12}} x_4^{a_{34}}, x_2^{a_2+a_{12}} - x_1^{a_1} x_3^{a_{23}-a_{13}} x_4^{a_{24}}\}$  is a standard basis for  $I_C$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$ . Then  $I_C^*$  is generated by the least homogeneous summands of the elements in  $G$ . In addition,  $\text{LM}(I_C^*) = \langle x_3^{a_{23}} x_4^{a_{24}}, x_2^{a_2+a_{12}}, x_3^{a_3}, x_4^{a_4}, x_2^{a_{12}} x_3^{a_{13}}, x_2^{a_{12}} x_4^{a_{34}} \rangle$ . Let

$$J_0 = \text{LM}(I_C^*), J_1 = \langle x_2^{a_2+a_{12}}, x_3^{a_3}, x_4^{a_4}, x_2^{a_{12}} x_4^{a_{34}}, x_2^{a_{12}} x_3^{a_{13}} \rangle,$$

$$J_2 = \langle x_2^{a_2+a_{12}}, x_3^{a_3}, x_4^{a_4}, x_2^{a_{12}} x_3^{a_{13}} \rangle, J_3 = \langle x_2^{a_2+a_{12}}, x_3^{a_3}, x_4^{a_4} \rangle.$$

Then  $J_i = \langle J_{i+1}, q_i \rangle$ , where  $q_0 = x_3^{a_{23}} x_4^{a_{24}}$ ,  $q_1 = x_2^{a_{12}} x_4^{a_{34}}$  and  $q_2 = x_2^{a_{12}} x_3^{a_{13}}$ . So

$$p(J_i) = p(J_{i+1}) - z^{\deg(q_i)} p(J_{i+1} : \langle q_i \rangle), \text{ for } 0 \leq i \leq 2.$$

In this case,  $J_1 : \langle q_0 \rangle = \langle x_2^{a_{12}}, x_3^{a_{13}}, x_4^{a_{34}} \rangle$ ,  $J_2 : \langle q_1 \rangle = \langle x_2^{a_2}, x_3^{a_{13}}, x_4^{a_{24}} \rangle$  and  $J_3 : \langle q_2 \rangle = \langle x_2^{a_2}, x_3^{a_{23}}, x_4^{a_4} \rangle$ . We have that  $p(J_1 : \langle q_0 \rangle) = (1 - z^{a_{12}})(1 - z^{a_{13}})(1 - z^{a_{34}})$ ,  $p(J_2 : \langle q_1 \rangle) = (1 - z^{a_2})(1 - z^{a_{13}})(1 - z^{a_{24}})$ ,  $p(J_3 : \langle q_2 \rangle) = (1 - z^{a_2})(1 - z^{a_{23}})(1 - z^{a_4})$  and  $p(J_3) = (1 - z^{a_2+a_{12}})(1 - z^{a_3})(1 - z^{a_4})$ . Thus

$$p(J_0) = (1 - z^{a_2+a_{12}})(1 - z^{a_3})(1 - z^{a_4}) - z^{a_{12}+a_{13}}(1 - z^{a_2})(1 - z^{a_{23}})(1 - z^{a_4}) - z^{a_{12}+a_{34}}(1 - z^{a_2})(1 - z^{a_{13}})(1 - z^{a_{24}}) - z^{a_{23}+a_{24}}(1 - z^{a_{12}})(1 - z^{a_{13}})(1 - z^{a_{34}}).$$

Thus  $p(J_0) = (1 - z)^3 h(z)$ , and therefore  $\text{HS}(R/\text{LM}(I_C^*), z) = \frac{h(z)}{1-z}$ . □

The proof of the next proposition is similar to that of Proposition 2.9 and therefore it is omitted.

**Proposition 2.13** *Suppose that  $I_C$  is given as in case 2(a) and also that  $a_{34} \leq a_{24}$ . (1) If  $a_{23} < a_{13}$ , then*

$$G = \{x_1^{a_1} - x_2^{a_{12}} x_3^{a_{13}}, x_2^{a_2} - x_3^{a_{23}} x_4^{a_{24}}, x_3^{a_3} - x_1^{a_{31}} x_4^{a_{34}}, x_4^{a_4} - x_1^{a_{41}} x_2^{a_{42}}, x_1^{a_{41}} x_3^{a_{23}} - x_2^{a_{12}} x_4^{a_{34}}, x_2^{a_2+a_{12}} - x_1^{a_{41}} x_3^{2a_{23}} x_4^{a_{24}-a_{34}}\}$$

is a standard basis for  $I_C$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$ .

(2) If  $a_{13} \leq a_{23}$ , then

$$G = \{x_1^{a_1} - x_2^{a_{12}} x_3^{a_{13}}, x_2^{a_2} - x_3^{a_{23}} x_4^{a_{24}}, x_3^{a_3} - x_1^{a_{31}} x_4^{a_{34}}, x_4^{a_4} - x_1^{a_{41}} x_2^{a_{42}}, x_1^{a_{41}} x_3^{a_{23}} - x_2^{a_{12}} x_4^{a_{34}}, x_2^{a_2+a_{12}} - x_1^{a_1} x_3^{a_{23}-a_{13}} x_4^{a_{24}}\}$$

is a standard basis for  $I_C$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$ .

**Theorem 2.14** *Suppose that  $I_C$  is given as in case 2(a) and also that  $a_{34} \leq a_{24}$ . Then the reduced Hilbert series of  $R/I_C^*$  is  $\text{HS}(R/I_C^*, z) = \frac{h(z)}{1-z}$  for*

$$h(z) = \sum_{i=0}^{a_2+a_{12}-1} z^i \sum_{i=0}^{a_3-1} z^i \sum_{i=0}^{a_4-1} z^i - z^{a_{12}+a_{13}} \sum_{i=0}^{a_2-1} z^i \sum_{i=0}^{a_{23}-1} z^i \sum_{i=0}^{a_4-1} z^i - z^{a_{12}+a_{34}} \sum_{i=0}^{a_2-1} z^i \sum_{i=0}^{a_{13}-1} z^i \sum_{i=0}^{a_{24}-1} z^i - z^{a_{23}+a_{24}} \sum_{i=0}^{a_{12}-1} z^i \sum_{i=0}^{a_{13}-1} z^i \sum_{i=0}^{a_{34}-1} z^i.$$

**Proof.** By Proposition 2.13,  $\text{LM}(I_C^*)$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$  is written as  $\text{LM}(I_C^*) = \langle x_3^{a_{23}} x_4^{a_{24}}, x_2^{a_2+a_{12}}, x_3^{a_3}, x_4^{a_4}, x_2^{a_{12}} x_4^{a_{34}}, x_2^{a_{12}} x_3^{a_{13}} \rangle$ . Let

$$J_0 = \text{LM}(I_C^*), J_1 = \langle x_2^{a_2+a_{12}}, x_3^{a_3}, x_4^{a_4}, x_2^{a_{12}} x_4^{a_{34}}, x_2^{a_{12}} x_3^{a_{13}} \rangle,$$

$$J_2 = \langle x_2^{a_2+a_{12}}, x_3^{a_3}, x_4^{a_4}, x_2^{a_{12}} x_3^{a_{13}} \rangle, J_3 = \langle x_2^{a_2+a_{12}}, x_3^{a_3}, x_4^{a_4} \rangle.$$

Then  $J_i = \langle J_{i+1}, q_i \rangle$ , where  $q_0 = x_3^{a_{23}} x_4^{a_{24}}$ ,  $q_1 = x_2^{a_{12}} x_4^{a_{34}}$  and  $q_2 = x_2^{a_{12}} x_3^{a_{13}}$ . Thus

$$p(J_i) = p(J_{i+1}) - z^{\deg(q_i)} p(J_{i+1} : \langle q_i \rangle), \text{ for } 0 \leq i \leq 2.$$

In this case,  $J_1 : \langle q_0 \rangle = \langle x_2^{a_{12}}, x_3^{a_{13}}, x_4^{a_{34}} \rangle$ ,  $J_2 : \langle q_1 \rangle = \langle x_2^{a_2}, x_3^{a_{13}}, x_4^{a_{24}} \rangle$  and  $J_3 : \langle q_2 \rangle = \langle x_2^{a_2}, x_3^{a_{23}}, x_4^{a_4} \rangle$ . Thus  $p(J_0) = (1 - z^{a_2+a_{12}})(1 - z^{a_3})(1 - z^{a_4}) - z^{a_{12}+a_{13}}(1 - z^{a_2})(1 - z^{a_{23}})(1 - z^{a_4}) - z^{a_{12}+a_{34}}(1 - z^{a_2})(1 - z^{a_{13}})(1 - z^{a_{24}}) - z^{a_{23}+a_{24}}(1 - z^{a_{12}})(1 - z^{a_{13}})(1 - z^{a_{34}})$ . So  $p(J_0) = (1 - z)^3 h(z)$ , and therefore  $HS(R/LM(I_C^*), z) = \frac{h(z)}{1-z}$ .  $\square$

We continue with case 2(b). If  $a_3 \leq a_{32} + a_{34}$ , then we can use [2, Remark 2.9] to find the reduced Hilbert series of  $R/I_C^*$ .

**Theorem 2.15** *Suppose that  $I_C$  is given as in case 2(b) and also that  $a_3 \leq a_{32} + a_{34}$ . Then the reduced Hilbert series of  $R/I_C^*$  is  $HS(R/I_C^*, z) = \frac{h(z)}{1-z}$  for*

$$h(z) = \prod_{i=2}^4 (1 + z + z^2 + \dots + z^{a_i-1}) - z^{a_{12}+a_{13}} \sum_{i=0}^{a_{32}-1} z^i \sum_{i=0}^{a_{43}-1} z^i \sum_{i=0}^{a_4-1} z^i - z^{a_{13}+a_{24}} \sum_{i=0}^{a_{12}-1} z^i \sum_{i=0}^{a_{43}-1} z^i \sum_{i=0}^{a_{34}-1} z^i.$$

**Proof.** By [2, Remark 2.9],

$$G = \{x_1^{a_1} - x_2^{a_{12}} x_3^{a_{13}}, x_2^{a_2} - x_1^{a_{21}} x_4^{a_{24}}, x_3^{a_3} - x_2^{a_{32}} x_4^{a_{34}}, x_4^{a_4} - x_1^{a_{41}} x_3^{a_{43}}, x_1^{a_{41}} x_2^{a_{32}} - x_3^{a_{13}} x_4^{a_{24}}\}$$

is a standard basis for  $I_C$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$ . So  $LM(I_C^*) = \langle x_3^{a_{13}} x_4^{a_{24}}, x_2^{a_2}, x_3^{a_3}, x_4^{a_4}, x_2^{a_{12}} x_3^{a_{13}} \rangle$ . Let

$$J_0 = \langle LM(I_C^*) \rangle, J_1 = \langle x_2^{a_2}, x_3^{a_3}, x_4^{a_4}, x_2^{a_{12}} x_3^{a_{13}} \rangle, J_2 = \langle x_2^{a_2}, x_3^{a_3}, x_4^{a_4} \rangle.$$

Then  $J_i = \langle J_{i+1}, q_i \rangle$ , where  $q_0 = x_3^{a_{13}} x_4^{a_{24}}$  and  $q_1 = x_2^{a_{12}} x_3^{a_{13}}$ . Thus

$$p(J_i) = p(J_{i+1}) - z^{\deg(q_i)} p(J_{i+1} : \langle q_i \rangle), \text{ for } 0 \leq i \leq 1.$$

In this case,  $J_1 : \langle q_0 \rangle = \langle x_2^{a_{12}}, x_3^{a_{43}}, x_4^{a_{34}} \rangle$  and  $J_2 : \langle q_1 \rangle = \langle x_2^{a_{32}}, x_3^{a_{43}}, x_4^{a_4} \rangle$ . We have that  $p(J_1 : \langle q_0 \rangle) = (1 - z^{a_{12}})(1 - z^{a_{43}})(1 - z^{a_{34}})$ ,  $p(J_2 : \langle q_1 \rangle) = (1 - z^{a_{32}})(1 - z^{a_{43}})(1 - z^{a_4})$  and  $p(J_2) = (1 - z^{a_2})(1 - z^{a_3})(1 - z^{a_4})$ . So  $p(J_0) = (1 - z^{a_2})(1 - z^{a_3})(1 - z^{a_4}) - z^{a_{12}+a_{13}}(1 - z^{a_{32}})(1 - z^{a_{43}})(1 - z^{a_4}) - z^{a_{13}+a_{24}}(1 - z^{a_{12}})(1 - z^{a_{43}})(1 - z^{a_{34}})$ . So  $p(J_0) = (1 - z)^3 h(z)$ . Thus  $HS(R/LM(I_C^*), z) = \frac{h(z)}{1-z}$ .  $\square$

The proof of the next proposition is similar to that of Proposition 2.7 and therefore it is omitted.

**Proposition 2.16** *Suppose that  $I_C$  is given as in case 2(b) and also that  $a_3 > a_{32} + a_{34}$ . If  $a_{34} < a_{24}$  and  $a_{12} \leq a_{32}$ , then*

$$G = \{x_1^{a_1} - x_2^{a_{12}} x_3^{a_{13}}, x_2^{a_2} - x_1^{a_{21}} x_4^{a_{24}}, x_3^{a_3} - x_2^{a_{32}} x_4^{a_{34}}, x_4^{a_4} - x_1^{a_{41}} x_3^{a_{43}}, x_1^{a_{41}} x_2^{a_{32}} - x_3^{a_{13}} x_4^{a_{24}}, x_3^{a_3+a_{13}} - x_1^{a_1} x_2^{a_{32}-a_{12}} x_4^{a_{34}}\}$$

is a standard basis for  $I_C$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$ .

**Theorem 2.17** Suppose that  $I_C$  is given as in case 2(b) and also that  $a_3 > a_{32} + a_{34}$ . Assume that  $a_{34} < a_{24}$  and  $a_{12} \leq a_{32}$ . Then the reduced Hilbert series of  $R/I_C^*$  is  $HS(R/I_C^*, z) = \frac{h(z)}{1-z}$  for

$$h(z) = \sum_{i=0}^{a_2-1} z^i \sum_{i=0}^{a_3+a_{13}-1} z^i \sum_{i=0}^{a_4-1} z^i - z^{a_{12}+a_{13}} \sum_{i=0}^{a_{32}-1} z^i \sum_{i=0}^{a_3-1} z^i \sum_{i=0}^{a_4-1} z^i - z^{a_{32}+a_{34}} \sum_{i=0}^{a_{12}-1} z^i \sum_{i=0}^{a_{13}-1} z^i \sum_{i=0}^{a_{24}-1} z^i - z^{a_{13}+a_{24}} \sum_{i=0}^{a_{12}-1} z^i \sum_{i=0}^{a_3-1} z^i \sum_{i=0}^{a_{34}-1} z^i.$$

**Proof.** By Proposition 2.16,  $G = \{x_1^{a_1} - x_2^{a_{12}}x_3^{a_{13}}, x_2^{a_2} - x_1^{a_{21}}x_4^{a_{24}}, x_3^{a_3} - x_2^{a_{32}}x_4^{a_{34}}, x_4^{a_4} - x_1^{a_{41}}x_3^{a_{43}}, x_1^{a_{41}}x_2^{a_{32}} - x_3^{a_{13}}x_4^{a_{24}}, x_3^{a_3+a_{13}} - x_1^{a_1}x_2^{a_{32}-a_{12}}x_4^{a_{34}}\}$  is a standard basis for  $I_C$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$ . Then  $LM(I_C^*)$  with respect to the aforementioned order is written as  $LM(I_C^*) = \langle x_2^{a_2}, x_3^{a_3+a_{13}}, x_4^{a_4}, x_2^{a_{32}}x_4^{a_{34}}, x_2^{a_{42}}x_3^{a_{13}}, x_3^{a_{13}}x_4^{a_{24}} \rangle$ . Let

$$J_0 = LM(I_C^*), J_1 = \langle x_2^{a_2}, x_3^{a_3+a_{13}}, x_4^{a_4}, x_2^{a_{12}}x_3^{a_{13}}, x_2^{a_{32}}x_4^{a_{34}} \rangle,$$

$$J_2 = \langle x_2^{a_2}, x_3^{a_3+a_{13}}, x_4^{a_4}, x_2^{a_{12}}x_3^{a_{13}} \rangle, J_3 = \langle x_2^{a_2}, x_3^{a_3+a_{13}}, x_4^{a_4} \rangle.$$

Note that  $J_i = \langle J_{i+1}, q_i \rangle$ , where  $q_0 = x_3^{a_{13}}x_4^{a_{24}}$ ,  $q_1 = x_2^{a_{32}}x_4^{a_{34}}$  and  $q_2 = x_2^{a_{12}}x_3^{a_{13}}$ . So

$$p(J_i) = p(J_{i+1}) - z^{\deg(q_i)}p(J_{i+1} : \langle q_i \rangle), \text{ for } 0 \leq i \leq 2.$$

In this case,  $J_1 : \langle q_0 \rangle = \langle x_2^{a_{12}}, x_3^{a_3}, x_4^{a_{34}} \rangle$ ,  $J_2 : \langle q_1 \rangle = \langle x_2^{a_{12}}, x_3^{a_{13}}, x_4^{a_{24}} \rangle$  and  $J_3 : \langle q_2 \rangle = \langle x_2^{a_{32}}, x_3^{a_3}, x_4^{a_4} \rangle$ . We have that  $p(J_1 : \langle q_0 \rangle) = (1 - z^{a_{12}})(1 - z^{a_3})(1 - z^{a_{34}})$ ,  $p(J_2 : \langle q_1 \rangle) = (1 - z^{a_{12}})(1 - z^{a_{13}})(1 - z^{a_{24}})$ ,  $p(J_3 : \langle q_2 \rangle) = (1 - z^{a_{32}})(1 - z^{a_3})(1 - z^{a_4})$  and  $p(J_3) = (1 - z^{a_2})(1 - z^{a_3+a_{13}})(1 - z^{a_4})$ . So

$$p(J_0) = (1 - z^{a_2})(1 - z^{a_3+a_{13}})(1 - z^{a_4}) - z^{a_{12}+a_{13}}(1 - z^{a_{32}})(1 - z^{a_3})(1 - z^{a_4}) - z^{a_{32}+a_{34}}(1 - z^{a_{12}})(1 - z^{a_{13}})(1 - z^{a_{24}}) - z^{a_{13}+a_{24}}(1 - z^{a_{12}})(1 - z^{a_3})(1 - z^{a_{34}}).$$

Thus  $p(J_0) = (1 - z)^3h(z)$ . So  $HS(R/LM(I_C^*), z) = \frac{h(z)}{1-z}$ . □

The proof of the next proposition is similar to that of Proposition 2.9 and therefore it is omitted.

**Proposition 2.18** Suppose that  $I_C$  is given as in case 2(b) and also that  $a_3 > a_{32} + a_{34}$ . Assume that  $a_{24} \leq a_{34}$ . (1) If  $a_{32} < a_{12}$ , then

$$G = \{x_1^{a_1} - x_2^{a_{12}}x_3^{a_{13}}, x_2^{a_2} - x_1^{a_{21}}x_4^{a_{24}}, x_3^{a_3} - x_2^{a_{32}}x_4^{a_{34}}, x_4^{a_4} - x_1^{a_{41}}x_3^{a_{43}}, x_1^{a_{41}}x_2^{a_{32}} - x_3^{a_{13}}x_4^{a_{24}}, x_3^{a_3+a_{13}} - x_1^{a_{41}}x_2^{a_{32}}x_4^{a_{34}-a_{24}}\}$$

is a standard basis for  $I_C$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$ .

(2) If  $a_{12} \leq a_{32}$ , then

$$G = \{x_1^{a_1} - x_2^{a_{12}}x_3^{a_{13}}, x_2^{a_2} - x_1^{a_{21}}x_4^{a_{24}}, x_3^{a_3} - x_2^{a_{32}}x_4^{a_{34}}, x_4^{a_4} - x_1^{a_{41}}x_3^{a_{43}}, x_1^{a_{41}}x_2^{a_{32}} - x_3^{a_{13}}x_4^{a_{24}}, x_3^{a_3+a_{13}} - x_1^{a_1}x_2^{a_{32}-a_{12}}x_4^{a_{34}}\}$$

is a standard basis for  $I_C$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$ .

**Theorem 2.19** Suppose that  $I_C$  is given as in case 2(b) and also that  $a_3 > a_{32} + a_{34}$ . Assume that  $a_{24} \leq a_{34}$ . Then the reduced Hilbert series of  $R/I_C^*$  is  $\text{HS}(R/I_C^*, z) = \frac{h(z)}{1-z}$  for

$$h(z) = \sum_{i=0}^{a_2-1} z^i \sum_{i=0}^{a_3+a_{13}-1} z^i \sum_{i=0}^{a_4-1} z^i - z^{a_{13}+a_{24}} \sum_{i=0}^{a_2-1} z^i \sum_{i=0}^{a_3-1} z^i \sum_{i=0}^{a_{34}-1} z^i - z^{a_{12}+a_{13}} \sum_{i=0}^{a_{32}-1} z^i \sum_{i=0}^{a_3-1} z^i \sum_{i=0}^{a_{24}-1} z^i - z^{a_{32}+a_{34}} \sum_{i=0}^{a_{12}-1} z^i \sum_{i=0}^{a_{13}-1} z^i \sum_{i=0}^{a_{24}-1} z^i.$$

**Proof.** By Proposition 2.18,  $\text{LM}(I_C^*)$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$  is written as  $\text{LM}(I_C^*) = \langle x_2^{a_2}, x_3^{a_3+a_{13}}, x_4^{a_4}, x_3^{a_{13}} x_4^{a_{24}}, x_2^{a_{12}} x_3^{a_{13}}, x_2^{a_{32}} x_4^{a_{34}} \rangle$ . Let

$$J_0 = \text{LM}(I_C^*), J_1 = \langle x_2^{a_2}, x_3^{a_3+a_{13}}, x_4^{a_4}, x_3^{a_{13}} x_4^{a_{24}}, x_2^{a_{12}} x_3^{a_{13}} \rangle,$$

$$J_2 = \langle x_2^{a_2}, x_3^{a_3+a_{13}}, x_4^{a_4}, x_3^{a_{13}} x_4^{a_{24}} \rangle, J_3 = \langle x_2^{a_2}, x_3^{a_3+a_{13}}, x_4^{a_4} \rangle.$$

Then  $J_i = \langle J_{i+1}, q_i \rangle$ , where  $q_0 = x_2^{a_{32}} x_4^{a_{34}}$ ,  $q_1 = x_2^{a_{12}} x_3^{a_{13}}$  and  $q_2 = x_3^{a_{13}} x_4^{a_{24}}$ . So

$$p(J_i) = p(J_{i+1}) - z^{\deg(q_i)} p(J_{i+1} : \langle q_i \rangle), \text{ for } 0 \leq i \leq 2.$$

In this case,  $J_1 : \langle q_0 \rangle = \langle x_2^{a_{12}}, x_3^{a_{13}}, x_4^{a_{24}} \rangle$ ,  $J_2 : \langle q_1 \rangle = \langle x_2^{a_{32}}, x_3^{a_3}, x_4^{a_{24}} \rangle$  and  $J_3 : \langle q_2 \rangle = \langle x_2^{a_2}, x_3^{a_3}, x_4^{a_{34}} \rangle$ . We have that  $p(J_1 : \langle q_0 \rangle) = (1 - z^{a_{12}})(1 - z^{a_{13}})(1 - z^{a_{24}})$ ,  $p(J_2 : \langle q_1 \rangle) = (1 - z^{a_{32}})(1 - z^{a_3})(1 - z^{a_{24}})$ ,  $p(J_3 : \langle q_2 \rangle) = (1 - z^{a_2})(1 - z^{a_3})(1 - z^{a_{34}})$  and  $p(J_3) = (1 - z^{a_2})(1 - z^{a_3+a_{13}})(1 - z^{a_4})$ . Thus

$$p(J_0) = (1 - z^{a_2})(1 - z^{a_3+a_{13}})(1 - z^{a_4}) - z^{a_{13}+a_{24}}(1 - z^{a_2})(1 - z^{a_3})(1 - z^{a_{34}}) - z^{a_{12}+a_{13}}(1 - z^{a_{32}})(1 - z^{a_3})(1 - z^{a_{24}}) - z^{a_{32}+a_{34}}(1 - z^{a_{12}})(1 - z^{a_{13}})(1 - z^{a_{24}}).$$

So  $p(J_0) = (1 - z)^3 h(z)$ , and therefore  $\text{HS}(R/\text{LM}(I_C^*), z) = \frac{h(z)}{1-z}$ . □

**Theorem 2.20** Suppose that  $I_C$  is given as in case 3(a). Then the reduced Hilbert series of  $R/I_C^*$  is  $\text{HS}(R/I_C^*, z) = \frac{h(z)}{1-z}$  for

$$h(z) = \prod_{i=2}^4 (1 + z + z^2 + \dots + z^{a_i-1}) - z^{a_{12}+a_{14}} \sum_{i=0}^{a_{42}-1} z^i \sum_{i=0}^{a_3-1} z^i \sum_{i=0}^{a_{34}-1} z^i - z^{a_{14}+a_{23}} \sum_{i=0}^{a_{12}-1} z^i \sum_{i=0}^{a_{43}-1} z^i \sum_{i=0}^{a_{34}-1} z^i.$$

**Proof.** By [2, Remark 2.9],

$$G = \{x_1^{a_1} - x_2^{a_{12}} x_4^{a_{14}}, x_2^{a_2} - x_1^{a_{21}} x_3^{a_{23}}, x_3^{a_3} - x_1^{a_{31}} x_4^{a_{34}}, x_4^{a_4} - x_2^{a_{42}} x_3^{a_{43}}, x_3^{a_{23}} x_4^{a_{14}} - x_1^{a_{31}} x_2^{a_{42}}\}$$

is a standard basis for  $I_C$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$ . Then  $\text{LM}(I_C^*) = \langle x_2^{a_2}, x_3^{a_3}, x_4^{a_4}, x_2^{a_{12}} x_4^{a_{14}}, x_3^{a_{23}} x_4^{a_{14}} \rangle$ . Let

$$J_0 = \text{LM}(I_C^*), J_1 = \langle x_2^{a_2}, x_3^{a_3}, x_4^{a_4}, x_2^{a_{12}} x_4^{a_{14}} \rangle, J_2 = \langle x_2^{a_2}, x_3^{a_3}, x_4^{a_4} \rangle.$$

Note that  $J_i = \langle J_{i+1}, q_i \rangle$ , where  $q_0 = x_3^{a_{23}} x_4^{a_{14}}$  and  $q_1 = x_2^{a_{12}} x_4^{a_{14}}$ . So

$$p(J_i) = p(J_{i+1}) - z^{\deg(q_i)} p(J_{i+1} : \langle q_i \rangle), \text{ for } 0 \leq i \leq 1.$$

In this case,  $J_1 : \langle q_0 \rangle = \langle x_2^{a_{12}}, x_3^{a_{43}}, x_4^{a_{34}} \rangle$  and  $J_2 : \langle q_1 \rangle = \langle x_2^{a_{42}}, x_3^{a_{43}}, x_4^{a_{34}} \rangle$ . We have that  $p(J_1 : \langle q_0 \rangle) = (1 - z^{a_{12}})(1 - z^{a_{43}})(1 - z^{a_{34}})$ ,  $p(J_2 : \langle q_1 \rangle) = (1 - z^{a_{42}})(1 - z^{a_{43}})(1 - z^{a_{34}})$  and  $p(J_2) = (1 - z^{a_2})(1 - z^{a_3})(1 - z^{a_4})$ . So  $p(J_0) = (1 - z^{a_2})(1 - z^{a_3})(1 - z^{a_4}) - z^{a_{12}+a_{14}}(1 - z^{a_{42}})(1 - z^{a_{43}})(1 - z^{a_{34}}) - z^{a_{14}+a_{23}}(1 - z^{a_{12}})(1 - z^{a_{43}})(1 - z^{a_{34}})$ . So  $p(J_0) = (1 - z)^3 h(z)$ , and therefore  $\text{HS}(R/\text{LM}(I_C^*), z) = \frac{h(z)}{1-z}$ .  $\square$

The proof of the next proposition is similar to that of Proposition 2.7 and therefore it is omitted.

**Proposition 2.21** *Suppose that  $I_C$  is given as in case 3(b). If  $a_{23} < a_{43}$  and  $a_{14} \leq a_{24}$ , then*

$$G = \{x_1^{a_1} - x_2^{a_{12}} x_4^{a_{14}}, x_2^{a_2} - x_3^{a_{23}} x_4^{a_{24}}, x_3^{a_3} - x_1^{a_{31}} x_2^{a_{32}}, \\ x_4^{a_4} - x_1^{a_{41}} x_3^{a_{43}}, x_1^{a_{31}} x_4^{a_{24}} - x_2^{a_{12}} x_3^{a_{43}}, x_2^{a_2+a_{12}} - x_1^{a_1} x_3^{a_{23}} x_4^{a_{24}-a_{14}}\}$$

is a standard basis for  $I_C$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$ .

**Theorem 2.22** *Suppose that  $I_C$  is given as in case 3(b). If  $a_{23} < a_{43}$  and  $a_{14} \leq a_{24}$ , then the reduced Hilbert series of  $R/I_C^*$  is  $\text{HS}(R/I_C^*, z) = \frac{h(z)}{1-z}$  for*

$$h(z) = \sum_{i=0}^{a_2+a_{12}-1} z^i \sum_{i=0}^{a_3-1} z^i \sum_{i=0}^{a_4-1} z^i - z^{a_{12}+a_{43}} \sum_{i=0}^{a_2-1} z^i \sum_{i=0}^{a_{23}-1} z^i \sum_{i=0}^{a_4-1} z^i - \\ z^{a_{12}+a_{14}} \sum_{i=0}^{a_2-1} z^i \sum_{i=0}^{a_{43}-1} z^i \sum_{i=0}^{a_{24}-1} z^i - z^{a_{23}+a_{24}} \sum_{i=0}^{a_{12}-1} z^i \sum_{i=0}^{a_{43}-1} z^i \sum_{i=0}^{a_{14}-1} z^i.$$

**Proof.** By Proposition 2.21,  $G = \{x_1^{a_1} - x_2^{a_{12}} x_4^{a_{14}}, x_2^{a_2} - x_3^{a_{23}} x_4^{a_{24}}, x_3^{a_3} - x_1^{a_{31}} x_2^{a_{32}}, x_4^{a_4} - x_1^{a_{41}} x_3^{a_{43}}, x_1^{a_{31}} x_4^{a_{24}} - x_2^{a_{12}} x_3^{a_{43}}, x_2^{a_2+a_{12}} - x_1^{a_1} x_3^{a_{23}} x_4^{a_{24}-a_{14}}\}$  is a standard basis for  $I_C$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$ . Then

$$\text{LM}(I_C^*) = \langle x_2^{a_2+a_{12}}, x_3^{a_3}, x_4^{a_4}, x_2^{a_{12}} x_4^{a_{14}}, x_2^{a_{12}} x_3^{a_{43}}, x_3^{a_{23}} x_4^{a_{24}} \rangle.$$

Let

$$J_0 = \text{LM}(I_C^*), J_1 = \langle x_2^{a_2+a_{12}}, x_3^{a_3}, x_4^{a_4}, x_2^{a_{12}} x_3^{a_{43}}, x_2^{a_{12}} x_4^{a_{14}} \rangle, \\ J_2 = \langle x_2^{a_2+a_{12}}, x_3^{a_3}, x_4^{a_4}, x_2^{a_{12}} x_3^{a_{43}} \rangle, J_3 = \langle x_2^{a_2+a_{12}}, x_3^{a_3}, x_4^{a_4} \rangle.$$

Note that  $J_i = \langle J_{i+1}, q_i \rangle$ , where  $q_0 = x_3^{a_{23}} x_4^{a_{24}}$ ,  $q_1 = x_2^{a_{12}} x_4^{a_{14}}$  and  $q_2 = x_2^{a_{12}} x_3^{a_{43}}$ . So

$$p(J_i) = p(J_{i+1}) - t^{\deg(q_i)} p(J_{i+1} : \langle q_i \rangle), \text{ for } 0 \leq i \leq 2.$$

In this case,  $J_1 : \langle q_0 \rangle = \langle x_2^{a_{12}}, x_3^{a_{43}}, x_4^{a_{14}} \rangle$ ,  $J_2 : \langle q_1 \rangle = \langle x_2^{a_2}, x_3^{a_{43}}, x_4^{a_{24}} \rangle$  and  $J_3 : \langle q_2 \rangle = \langle x_2^{a_2}, x_3^{a_{23}}, x_4^{a_4} \rangle$ . We have that  $p(J_1 : \langle q_0 \rangle) = (1 - z^{a_{12}})(1 - z^{a_{43}})(1 - z^{a_{14}})$ ,  $p(J_2 : \langle q_1 \rangle) = (1 - z^{a_2})(1 - z^{a_{43}})(1 - z^{a_{24}})$ ,  $p(J_3 : \langle q_2 \rangle) = (1 - z^{a_2})(1 - z^{a_{23}})(1 - z^{a_4})$  and  $p(J_3) = (1 - z^{a_2+a_{12}})(1 - z^{a_3})(1 - z^{a_4})$ . So

$$p(J_0) = (1 - z^{a_2+a_{12}})(1 - z^{a_3})(1 - z^{a_4}) - z^{a_{12}+a_{43}}(1 - z^{a_2})(1 - z^{a_{23}})(1 - z^{a_4}) -$$



$$z^{a_{12}+a_{14}}(1-z^{a_2})(1-z^{a_{43}})(1-z^{a_{24}}) - z^{a_{23}+a_{24}}(1-z^{a_{12}})(1-z^{a_{43}})(1-z^{a_{14}}).$$

Thus  $p(J_0) = (1-z)^3 h(z)$ , and therefore  $\text{HS}(R/\text{LM}(I_C^*), z) = \frac{h(z)}{1-z}$ . □

The proof of the next proposition is similar to that of Proposition 2.9 and therefore it is omitted.

**Proposition 2.23** *Suppose that  $I_C$  is given as in case 3(b) and also that  $a_{43} \leq a_{23}$ . (1) If  $a_{24} < a_{14}$ , then*

$$G = \{x_1^{a_1} - x_2^{a_{12}}x_4^{a_{14}}, x_2^{a_2} - x_3^{a_{23}}x_4^{a_{24}}, x_3^{a_3} - x_1^{a_{31}}x_2^{a_{32}}, \\ x_4^{a_4} - x_1^{a_{41}}x_3^{a_{43}}, x_1^{a_{31}}x_4^{a_{24}} - x_2^{a_{12}}x_3^{a_{43}}, x_2^{a_2+a_{12}} - x_1^{a_{31}}x_3^{a_{23}-a_{43}}x_4^{a_{24}}\}$$

is a standard basis for  $I_C$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$ .

(2) If  $a_{14} \leq a_{24}$ , then

$$G = \{x_1^{a_1} - x_2^{a_{12}}x_4^{a_{14}}, x_2^{a_2} - x_3^{a_{23}}x_4^{a_{24}}, x_3^{a_3} - x_1^{a_{31}}x_2^{a_{32}}, \\ x_4^{a_4} - x_1^{a_{41}}x_3^{a_{43}}, x_1^{a_{31}}x_4^{a_{24}} - x_2^{a_{12}}x_3^{a_{43}}, x_2^{a_2+a_{12}} - x_1^{a_1}x_3^{a_{23}}x_4^{a_{24}-a_{14}}\}$$

is a standard basis for  $I_C$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$ .

**Theorem 2.24** *Suppose that  $I_C$  is given as in case 3(b) and also that  $a_{43} \leq a_{23}$ . Then the reduced Hilbert series of  $R/I_C^*$  is  $\text{HS}(R/I_C^*, z) = \frac{h(z)}{1-z}$  for*

$$h(z) = \sum_{i=0}^{a_2+a_{12}-1} z^i \sum_{i=0}^{a_3-1} z^i \sum_{i=0}^{a_4-1} z^i - z^{a_{12}+a_{43}} \sum_{i=0}^{a_2-1} z^i \sum_{i=0}^{a_{23}-1} z^i \sum_{i=0}^{a_4-1} z^i - \\ z^{a_{12}+a_{14}} \sum_{i=0}^{a_2-1} z^i \sum_{i=0}^{a_{43}-1} z^i \sum_{i=0}^{a_{24}-1} z^i - z^{a_{23}+a_{24}} \sum_{i=0}^{a_{12}-1} z^i \sum_{i=0}^{a_{43}-1} z^i \sum_{i=0}^{a_{14}-1} z^i.$$

**Proof.** By Proposition 2.23,  $\text{LM}(I_C^*)$  with respect to the negative degree reverse lexicographic term ordering with  $x_4 > x_3 > x_2 > x_1$  is written as  $\text{LM}(I_C^*) = \langle x_2^{a_2+a_{12}}, x_3^{a_3}, x_4^{a_4}, x_2^{a_{12}}x_3^{a_{43}}, x_2^{a_{12}}x_4^{a_{14}}, x_3^{a_{23}}x_4^{a_{24}} \rangle$ . Let

$$J_0 = \text{LM}(I_C^*), J_1 = \langle x_2^{a_2+a_{12}}, x_3^{a_3}, x_4^{a_4}, x_2^{a_{12}}x_3^{a_{43}}, x_2^{a_{12}}x_4^{a_{14}} \rangle,$$

$$J_2 = \langle x_2^{a_2+a_{12}}, x_3^{a_3}, x_4^{a_4}, x_2^{a_{12}}x_3^{a_{43}} \rangle, J_3 = \langle x_2^{a_2+a_{12}}, x_3^{a_3}, x_4^{a_4} \rangle.$$

Then  $J_i = \langle J_{i+1}, q_i \rangle$ , where  $q_0 = x_3^{a_{23}}x_4^{a_{24}}$ ,  $q_1 = x_2^{a_{12}}x_4^{a_{14}}$  and  $q_2 = x_2^{a_{12}}x_3^{a_{43}}$ . Thus

$$p(J_i) = p(J_{i+1}) - z^{\deg(q_i)} p(J_{i+1} : \langle q_i \rangle), \text{ for } 0 \leq i \leq 2.$$

In this case,  $J_1 : \langle q_0 \rangle = \langle x_2^{a_{12}}, x_3^{a_{43}}, x_4^{a_{14}} \rangle$ ,  $J_2 : \langle q_1 \rangle = \langle x_2^{a_2}, x_3^{a_{43}}, x_4^{a_{24}} \rangle$  and  $J_3 : \langle q_2 \rangle = \langle x_2^{a_2}, x_3^{a_{23}}, x_4^{a_4} \rangle$ . So

$$p(J_0) = (1-z^{a_2+a_{12}})(1-z^{a_3})(1-z^{a_4}) - z^{a_{12}+a_{43}}(1-z^{a_2})(1-z^{a_{23}})(1-z^{a_4}) - \\ z^{a_{12}+a_{14}}(1-z^{a_2})(1-z^{a_{43}})(1-z^{a_{24}}) - z^{a_{23}+a_{24}}(1-z^{a_{12}})(1-z^{a_{43}})(1-z^{a_{14}}).$$

Now we have that  $p(J_0) = (1-z)^3 h(z)$ . Therefore  $\text{HS}(R/\text{LM}(I_C^*), z) = \frac{h(z)}{1-z}$ . □

**3. Examples**

In this section, we give some examples showing how the formulas in the previous section can be used to compute  $HS(R/I_C^*, z)$ .

**Example 3.1** Let  $m \geq 5$  be an integer such that  $\gcd(51, 13m + 1, 14m + 5, 32m + 26) = 1$ . Consider the monomial curve  $C(m)$  defined parametrically by  $x_1 = t^{51}$ ,  $x_2 = t^{13m+1}$ ,  $x_3 = t^{14m+5}$  and  $x_4 = t^{32m+26}$ . The ideal  $I_{C(m)}$  is minimally generated by  $x_1^{2m+1} - x_3^5 x_4$ ,  $x_2^5 - x_1^m x_3$ ,  $x_3^6 - x_2^4 x_4$ ,  $x_4^2 - x_1^{m+1} x_2$  and  $x_1^m x_4 - x_2 x_3^5$ . Thus we are in case 1(b) of Remark 2.2 and from Theorem 2.3 we deduce that  $R/I_{C(m)}^*$  is Cohen–Macaulay. By Theorem 2.10, the numerator of the reduced Hilbert series of  $R/I_{C(m)}^*$  is

$$h(z) = (1 + z) \sum_{i=0}^4 z^i \sum_{i=0}^{10} z^i - z^6(1 + z) \sum_{i=0}^3 z^i \sum_{i=0}^5 z^i - z^5 \sum_{i=0}^4 z^i - z^6 \sum_{i=0}^5 z^i.$$

We have that

$$(1 + z) \sum_{i=0}^4 z^i \sum_{i=0}^{10} z^i = 1 + 3z + 5z^2 + 7z^3 + 9z^4 + \sum_{i=5}^{10} 10z^i + 9z^{11} + 7z^{12} + 5z^{13} + 3z^{14} + z^{15}$$

and also

$$z^6(1 + z) \sum_{i=0}^3 z^i \sum_{i=0}^5 z^i = z^6 + 3z^7 + 5z^8 + 7z^9 + 8z^{10} + 8z^{11} + 7z^{12} + 5z^{13} + 3z^{14} + z^{15}.$$

So  $h(z) = \sum_{i=0}^4 (2i + 1)z^i + \sum_{i=5}^9 (19 - 2i)z^i + z^{10}$ , and therefore

$$HS(R/I_{C(m)}^*, z) = \frac{\sum_{i=0}^4 (2i + 1)z^i + \sum_{i=5}^9 (19 - 2i)z^i + z^{10}}{1 - z}.$$

**Example 3.2** Let  $m \geq 4$  be an integer such that  $\gcd(3m + 19, 13m + 36, 3m^2 + 15m + 21, 4m^2 + 17m + 9) = 1$ . Consider the monomial curve  $C(m)$  defined parametrically by  $x_1 = t^{3m+19}$ ,  $x_2 = t^{13m+36}$ ,  $x_3 = t^{3m^2+15m+21}$  and  $x_4 = t^{4m^2+17m+9}$ . The ideal  $I_{C(m)}$  is minimally generated by  $x_1^{m+3} - x_2 x_3$ ,  $x_2^{m+2} - x_3^3 x_4$ ,  $x_3^4 - x_1^3 x_4^3$ ,  $x_4^4 - x_1^m x_2^{m+1}$  and  $x_1^m x_3^3 - x_2 x_4^3$ . Thus we are in case 2(a) of Remark 2.2 and from Theorem 2.3 we deduce that  $R/I_{C(m)}^*$  is Cohen–Macaulay. By Theorem 2.12, the numerator of the reduced Hilbert series of  $R/I_{C(m)}^*$  is

$$h(z) = \sum_{i=0}^{m+2} z^i \left( \sum_{i=0}^3 z^i \right)^2 - z^2 \sum_{i=0}^{m+1} z^i \sum_{i=0}^2 z^i \sum_{i=0}^3 z^i - z^4 \sum_{i=0}^{m+1} z^i - z^4 \sum_{i=0}^2 z^i.$$

We have that  $\left( \sum_{i=0}^3 z^i \right)^2 = 1 + 2z + 3z^2 + 4z^3 + 3z^4 + 2z^5 + z^6$  and also

$$\sum_{i=0}^{m+2} z^i \left( \sum_{i=0}^3 z^i \right)^2 = 1 + 3z + 6z^2 + 10z^3 + 13z^4 + 15z^5 + \sum_{i=6}^{m+2} 16z^i +$$

$$15z^{m+3} + 13z^{m+4} + 10z^{m+5} + 6z^{m+6} + 3z^{m+7} + z^{m+8}.$$

In addition,

$$\begin{aligned} \sum_{i=0}^{m+1} z^i \sum_{i=0}^2 z^i \sum_{i=0}^3 z^i &= \sum_{i=0}^{m+1} z^i (1 + 2z + 3z^2 + 3z^3 + 2z^4 + z^5) = 1 + 3z + 6z^2 + \\ &9z^3 + 11z^4 + \sum_{i=5}^{m+1} 12z^i + 11z^{m+2} + 9z^{m+3} + 6z^{m+4} + 3z^{m+5} + z^{m+6}, \end{aligned}$$

so

$$\begin{aligned} z^2 \sum_{i=0}^{m+1} z^i \sum_{i=0}^2 z^i \sum_{i=0}^3 z^i &= z^2 + 3z^3 + 6z^4 + 9z^5 + 11z^6 + \sum_{i=7}^{m+3} 12z^i + 11z^{m+4} + \\ &9z^{m+5} + 6z^{m+6} + 3z^{m+7} + z^{m+8}. \end{aligned}$$

Thus  $h(z) = \sum_{i=0}^3 (2i + 1)z^i + 5z^4 + 4z^5 + \sum_{i=6}^{m+2} 3z^i + 2z^{m+3} + z^{m+4}$ , and therefore

$$\text{HS}(R/I_{C(m)}^*, z) = \frac{\sum_{i=0}^3 (2i + 1)z^i + 5z^4 + 4z^5 + \sum_{i=6}^{m+2} 3z^i + 2z^{m+3} + z^{m+4}}{1 - z}.$$

**Example 3.3** Let  $m \geq 7$  be an integer such that  $\text{gcd}(2m + 21, 14m + 5, 2m^2 + m + 3, 2m^2 + 7m - 5) = 1$ . Consider the monomial curve  $C(m)$  defined parametrically by  $x_1 = t^{2m+21}$ ,  $x_2 = t^{14m+5}$ ,  $x_3 = t^{2m^2+m+3}$  and  $x_4 = t^{2m^2+7m-5}$ . The ideal  $I_{C(m)}$  is minimally generated by  $x_1^m - x_2x_4$ ,  $x_2^{m+1} - x_3^5x_4^2$ ,  $x_3^7 - x_1x_2^m$ ,  $x_4^3 - x_1^{m-1}x_3^2$  and  $x_1x_4^2 - x_2x_3^2$ . Thus we are in case 3(b) of Remark 2.2 and from Theorem 2.3 we deduce that  $R/I_{C(m)}^*$  is Cohen–Macaulay. By Theorem 2.24, the numerator of the reduced Hilbert series of  $R/I_{C(m)}^*$  is

$$h(z) = \sum_{i=0}^{m+1} z^i \sum_{i=0}^6 z^i \sum_{i=0}^2 z^i - z^3 \sum_{i=0}^m z^i \sum_{i=0}^4 z^i \sum_{i=0}^2 z^i - z^2(1+z)^2 \sum_{i=0}^m z^i - z^7(1+z).$$

We have that

$$\sum_{i=0}^6 z^i \sum_{i=0}^2 z^i = 1 + 2z + \sum_{i=2}^6 3z^i + 2z^7 + z^8$$

and also

$$\begin{aligned} \sum_{i=0}^{m+1} z^i \sum_{i=0}^6 z^i \sum_{i=0}^2 z^i &= 1 + 3z + 6z^2 + 9z^3 + 12z^4 + 15z^5 + 18z^6 + 20z^7 + \sum_{i=8}^{m+1} 21z^i + \\ &20z^{m+2} + 18z^{m+3} + 15z^{m+4} + 12z^{m+5} + 9z^{m+6} + 6z^{m+7} + 3z^{m+8} + z^{m+9}. \end{aligned}$$

In addition,

$$z^3 \sum_{i=0}^4 z^i \sum_{i=0}^2 z^i = z^3 + 2z^4 + \sum_{i=5}^7 3z^i + 2z^8 + z^9$$

and also

$$z^2 \sum_{i=0}^m z^i \sum_{i=0}^4 z^i \sum_{i=0}^2 z^i = z^3 + 3z^4 + 6z^5 + 9z^6 + 12z^7 + 14z^8 + \sum_{i=9}^{m+3} 15z^i +$$

$$14z^{m+4} + 12z^{m+5} + 9z^{m+6} + 6z^{m+7} + 3z^{m+8} + z^{m+9}.$$

Moreover

$$z^2(1+z)^2 \sum_{i=0}^m z^i = (z^2 + 2z^3 + z^4) \sum_{i=0}^m z^i = z^2 + 3z^3 + \sum_{i=4}^{m+2} 4z^i + 3z^{m+3} + z^{m+4}.$$

Thus  $h(z) = 1 + 3z + \sum_{i=2}^6 5z^i + 3z^7 + \sum_{i=8}^{m+1} 2z^i + z^{m+2}$ , and therefore

$$\text{HS}(R/I_{C(m)}^*, z) = \frac{1 + 3z + \sum_{i=2}^6 5z^i + 3z^7 + \sum_{i=8}^{m+1} 2z^i + z^{m+2}}{1 - z}.$$

### References

- [1] Arslan F, Katsabekis A, Nalbandiyan M. On the Cohen-Macaulayness of tangent cones of monomial curves in  $\mathbb{A}^4(K)$ . Turkish Journal of Mathematics 2019; 43: 1425-1446.
- [2] Arslan F, Mete P. Hilbert functions of Gorenstein monomial curves. Proceedings of the American Mathematical Society 2007; 135: 1993-2002.
- [3] Bayer D, Stillman M. Computation of Hilbert functions. Journal of Symbolic Computation 1992; 14: 31-50.
- [4] Bresinsky H. Symmetric semigroups of integers generated by 4 elements. Manuscripta Mathematica 1975; 17 (3): 205-219.
- [5] Greuel G-M, Pfister G. A Singular Introduction to Commutative Algebra. Berlin, Germany: Springer-Verlag, 2002.
- [6] Kunz E. The value semigroup of one dimensional Gorenstein ring. Proceedings of the American Mathematical Society 1970; 25: 748-751.
- [7] Sturmfels B. Gröbner Bases and Convex Polytopes. University Lecture Series Vol. 8. Providence, RI, USA: American Mathematical Society, 1996.