

## Inverse nodal problems for Dirac type integro differential system with a nonlocal boundary condition

Baki KESKİN\* 

Department of Mathematics, Faculty of Science, Sivas Cumhuriyet University, Sivas, Turkey

Received: 18.04.2022

Accepted/Published Online: 21.05.2022

Final Version: 04.07.2022

**Abstract:** In this paper, the Dirac type integro differential system with a nonlocal integral boundary condition is considered. First, we derive the asymptotic expressions for the solutions and large eigenvalues. Second, we provide asymptotic expressions for the nodal points and prove that a dense subset of nodal points uniquely determines the boundary condition parameter and the potential function of the considered differential system. We also provide an effective procedure for solving the inverse nodal problem.

**Key words:** Dirac operator, integro-differential operators, inverse nodal problems, nonlocal boundary conditions, boundary value problem

### 1. Introduction

In recent years, boundary value problems with nonlocal conditions are one of the fastest developing current topics in mathematical physics. Nonlocal boundary conditions appear when we cannot measure data directly at the boundary. Problems of this type arise in various fields of mathematical physics, biology, biotechnology, mechanics, geophysics and other branches of natural sciences [2], [3], [4], [5], [6]. Such problems were studied firstly by Samarskii and Bitsadze. They formulated and investigated the nonlocal boundary problem for an elliptic equation [1].

Inverse spectral problems consist in recovering operators from their spectral characteristics. Results of the inverse problem for various nonlocal operators can be found in [7], [8], [9], [10], [41], [42].

In a classical inverse spectral problem, the potential and the coefficients of the operators are to be determined from spectral data (e.g., two sets of eigenvalues, or one set of eigenvalues and norming constants). This is called the inverse eigenvalue problem. An alternative is to take as data the zeros (nodes) of the eigenfunctions of the considered operators, which are just as experimentally observable as eigenvalues in some situations. This is generally referred to as the "Inverse Nodal Problem".

In 1988, McLaughlin raised and solved the inverse nodal problem for Sturm-Liouville problems for the first time [31]. She showed that knowledge of a dense subset of zeros (nodes) of the eigenfunctions alone can determine the potential function of the Sturm-Liouville problem up to a constant. Hald and McLaughlin [23] provided some numerical schemes for the reconstruction of the potential for more general boundary conditions. In 1997, Yang suggested a constructive procedure for reconstructing the potential and the boundary condition of the Sturm-Liouville problem from nodes of its eigenfunctions [39]. Inverse nodal problems have been addressed

2010 *AMS Mathematics Subject Classification:* 26A33, 34A55, 34L05, 34L20, 34K29, 34K10, 47G20.

by various researchers in several papers for different operators with different kinds of boundary conditions ([15], [18], [19], [21], [22], [33], [34], [38], [37], [40], [35], [43], [44], [45] and references therein).

In recent years, integro-differential operators attracted much attention of mathematicians. Such operators have important applications in many fields of science (see monographs [14], [32] and references therein). Therefore, many researchers are currently working on inverse problems for these operators ([11], [12], [13], [16], [17], [20], [24], [25], [30], [36] and [46]). The inverse nodal problem for Dirac type integro-differential operators with Robin boundary conditions was first studied by [26]. This operator with parameter dependent boundary conditions linearly and nonlinearly were studied by [27] and [28], respectively. In their studies, the authors considered  $p(x)$  and  $r(x)$ , which are the components of the potential function  $Q(x)$ , as a special case such that  $p(x) - r(x) = \text{const}$ .

In this study, the perturbation of the Dirac differential system by a Volterra type operator on a finite interval with one classical boundary condition and another nonlocal integral boundary condition is considered. We also consider  $p(x)$  and  $r(x)$  as two independent functions and investigate a more general case. In this way, we have the opportunity to determine  $p(x)$  and  $r(x)$  separately. It is shown that the coefficients of the differential part of the operator and the boundary condition parameter can be determined by using a dense subset of the nodes. For this problem, we also give a constructive procedure for determining the coefficients as well as the useful asymptotics regarding the solution, eigenvalues and nodes.

## 2. Main results

Consider the nonlocal boundary value problem of the Dirac type integro-differential system

$$BY' + Q(x)Y + \int_0^x K(x, \varsigma)Y d\varsigma = \lambda Y, \quad x \in (0, \pi), \tag{2.1}$$

with one classical boundary condition

$$y_1(0) \sin \theta + y_2(0) \cos \theta = 0 \tag{2.2}$$

and nonlocal integral boundary condition

$$y_1(\pi) - \int_0^\pi y_1(x)\omega(x)dx = 0, \tag{2.3}$$

where  $0 < \theta < \pi$  are real numbers,  $\lambda$  is the spectral parameter,

$B = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ ,  $Q(x) = \begin{pmatrix} p(x) & 0 \\ 0 & r(x) \end{pmatrix}$ ,  $K(x, \varsigma) = \begin{pmatrix} K_{11}(x, \varsigma) & K_{12}(x, \varsigma) \\ K_{21}(x, \varsigma) & K_{22}(x, \varsigma) \end{pmatrix}$ ,  $Y = \begin{pmatrix} y_1 \\ y_2 \end{pmatrix}$ ,  $p(x)$ ,  $r(x)$ ,  $\omega(x)$  and  $K_{ij}(x, \varsigma)$ , ( $i, j = 1, 2$ ) are real-valued functions in  $W_2^1(0, \pi)$ .

Let  $\varphi(x, \lambda) = (\varphi_1(x, \lambda), \varphi_2(x, \lambda))^T$  be the solution of (2.1) satisfying the initial condition  $\varphi(0, \lambda) =$

$(\cos \theta, -\sin \theta)^T \cdot \varphi(x, \lambda)$  is an entire function of  $\lambda$  and satisfies:

$$\begin{aligned} \varphi_1(x, \lambda) &= \cos \theta \cos \lambda x + \sin \theta \sin \lambda x \\ &+ \int_0^x \sin \lambda(x - \varsigma) p(\varsigma) \varphi_1(\varsigma, \lambda) d\varsigma + \int_0^x \cos \lambda(x - \varsigma) r(\varsigma) \varphi_2(\varsigma, \lambda) d\varsigma \\ &+ \int_0^x \int_0^\varsigma \sin \lambda(x - \varsigma) \{K_{11}(\varsigma, \xi) \varphi_1(\lambda, \xi) + K_{12}(\varsigma, \xi) \varphi_2(\lambda, \xi)\} d\xi d\varsigma \\ &+ \int_0^x \int_0^\varsigma \cos \lambda(x - \varsigma) \{K_{21}(\varsigma, \xi) \varphi_1(\lambda, \xi) + K_{22}(\varsigma, \xi) \varphi_2(\lambda, \xi)\} d\xi d\varsigma \end{aligned} \tag{2.4}$$

$$\begin{aligned} \varphi_2(x, \lambda) &= \cos \theta \sin \lambda x - \sin \theta \cos \lambda x \\ &- \int_0^x \cos \lambda(x - \varsigma) p(\varsigma) \varphi_1(\varsigma, \lambda) d\varsigma + \int_0^x \sin \lambda(x - \varsigma) r(\varsigma) \varphi_2(\varsigma, \lambda) d\varsigma \\ &- \int_0^x \int_0^\varsigma \cos \lambda(x - \varsigma) \{K_{11}(\varsigma, \xi) \varphi_1(\lambda, \xi) + K_{12}(\varsigma, \xi) \varphi_2(\lambda, \xi)\} d\xi d\varsigma \\ &+ \int_0^x \int_0^\varsigma \sin \lambda(x - \varsigma) \{K_{21}(\varsigma, \xi) \varphi_1(\lambda, \xi) + K_{22}(\varsigma, \xi) \varphi_2(\lambda, \xi)\} d\xi d\varsigma. \end{aligned}$$

**Theorem 2.1** ([25], the case:  $\alpha = 1$ ) For  $|\lambda| \rightarrow \infty$ , the following asymptotic formulae are valid:

$$\begin{aligned} \varphi_1(x, \lambda) &= \cos(\lambda x - \sigma(x) - \theta) + \frac{1}{2\lambda} \eta(x) \cos(\lambda x - \sigma(x) - \theta) \\ &- \frac{1}{2\lambda} \eta(0) \cos(\lambda x - \sigma(x) + \theta) + \frac{1}{2\lambda} \sin(\lambda x - \sigma(x) - \theta) \int_0^x \eta^2(\varsigma) d\varsigma \\ &- \frac{1}{2\lambda} K_+(x) \cos(\lambda x - \sigma(x) - \theta) - \frac{1}{2\lambda} K_-(x) \sin(\lambda x - \sigma(x) - \theta) \\ &+ o\left(\frac{1}{\lambda} \exp(|\tau| x)\right), \end{aligned} \tag{2.5}$$

$$\begin{aligned} \varphi_2(x, \lambda) &= \sin(\lambda x - \sigma(x) - \theta) - \frac{1}{2\lambda} \eta(x) \sin(\lambda x - \sigma(x) - \theta) \\ &- \frac{1}{2\lambda} \eta(0) \sin(\lambda x - \sigma(x) + \theta) - \frac{1}{2\lambda} \cos(\lambda x - \sigma(x) - \theta) \int_0^x \eta^2(\varsigma) d\varsigma \\ &- \frac{1}{2\lambda} K_+(x) \sin(\lambda x - \sigma(x) - \theta) + \frac{1}{2\lambda} K_-(x) \cos(\lambda x - \sigma(x) - \theta) \\ &+ o\left(\frac{1}{\lambda} \exp(|\tau| x)\right), \end{aligned} \tag{2.6}$$

uniformly in  $x \in [0, \pi]$ , where  $\sigma(x) = \frac{1}{2} \int_0^x (p(\varsigma) + r(\varsigma)) d\varsigma$ ,  $\eta(x) = \frac{1}{2} (p(x) - r(x))$ ,

$K_+(x) = \int_0^x (K_{11}(\varsigma, \varsigma) + K_{22}(\varsigma, \varsigma)) d\varsigma$ ,  $K_-(x) = \int_0^x (K_{12}(\varsigma, \varsigma) - K_{21}(\varsigma, \varsigma)) d\varsigma$

and  $\tau = \text{Im}\lambda$ .

The zero-sequences  $\{\lambda_n\}_{n \in \mathbb{Z}}$  of the entire function

$$\Lambda(\lambda) := \varphi_1(\pi, \lambda) - \int_0^\pi \varphi_1(x, \lambda)\omega(x)dx = 0$$

are the spectra of the boundary value problem (2.1)–(2.3) and  $\Lambda$  satisfies

$$\begin{aligned} \Lambda(\lambda) &= \cos(\lambda\pi - \sigma(\pi) - \theta) + \frac{1}{2\lambda}\eta(\pi)\cos(\lambda\pi - \sigma(\pi) - \theta) \\ &\quad - \frac{1}{2\lambda}\eta(0)\cos(\lambda\pi - \sigma(\pi) + \theta) + \frac{1}{2\lambda}\sin(\lambda\pi - \sigma(\pi) - \theta)\int_0^\pi \eta^2(\varsigma)d\varsigma \\ &\quad - \frac{1}{2\lambda}K_+(\pi)\cos(\lambda\pi - \sigma(\pi) - \theta) - \frac{1}{2\lambda}K_-(\pi)\sin(\lambda\pi - \sigma(\pi) - \theta) \\ &\quad - \frac{1}{\lambda}\sin(\lambda\pi - \sigma(\pi) - \theta)\omega(\pi) + o\left(\frac{1}{\lambda}\exp(|\tau|\pi)\right), \end{aligned} \tag{2.7}$$

for sufficiently large  $|\lambda|$ . Since the eigenvalues of the problem (2.1)–(2.3) are the roots of  $\Lambda(\lambda_n) = 0$ , we can write the following equation for them:

$$\begin{aligned} &\left(1 + \frac{1}{2\lambda_n}\eta(\pi) - \frac{1}{2\lambda_n}\eta(0)\cos 2\theta - \frac{1}{2\lambda_n}K_+(\pi)\right)\tan(\lambda_n\pi - \sigma(\pi) - \frac{\pi}{2} - \theta) = \\ &\frac{1}{2\lambda_n}\eta(0)\sin 2\theta + \frac{1}{2\lambda_n}\int_0^\pi \eta^2(\varsigma)d\varsigma - \frac{1}{2\lambda_n}K_-(\pi) - \frac{1}{\lambda_n}\omega(\pi) + o\left(\frac{1}{\lambda_n}\right) \end{aligned}$$

from this expression, for sufficiently large  $|n|$ ,

$$\begin{aligned} \lambda_n &= \left(n + \frac{1}{2}\right) + \frac{\theta + \sigma(\pi)}{\pi} \\ &\quad + \frac{1}{2n\pi}\left(\eta(0)\sin 2\theta + \int_0^\pi \eta^2(\varsigma)d\varsigma - K_-(\pi) - 2\omega(\pi)\right) + o\left(\frac{1}{n}\right), \quad n \geq 1, \end{aligned} \tag{2.8}$$

and similarly

$$\begin{aligned} \lambda_n &= \left(n - \frac{1}{2}\right) + \frac{\theta + \sigma(\pi)}{\pi} \\ &\quad - \frac{1}{2n\pi}\left(\eta(0)\sin 2\theta + \int_0^\pi \eta^2(\varsigma)d\varsigma - K_-(\pi) - 2\omega(\pi)\right) + o\left(\frac{1}{n}\right), \quad n \leq -1. \end{aligned} \tag{2.9}$$

**Lemma 2.2** For sufficiently large  $n$ ,  $\varphi_1(x, \lambda_n)$  has exactly  $n$  nodes  $\{x_n^k : k = 0, 1, \dots, n - 1\}$  in the interval

$(0, \pi)$ :  $0 < x_n^0 < x_n^1 < \dots < x_n^{n-1} < \pi$ . Moreover,

$$\begin{aligned}
 x_n^k &= \frac{(k + 1/2)\pi}{n} + \frac{\sigma(x_n^k) + \theta}{n} \\
 &\quad - \frac{(k + 1/2)\pi}{n} \frac{(\pi + 2\theta + 2\sigma(\pi))}{2n\pi} - (\sigma(x_n^k) + \theta) \frac{(\pi + 2\theta + 2\sigma(\pi))}{2n^2\pi} \\
 &\quad - \frac{(k + 1/2)\pi}{n} \frac{1}{2n^2\pi} \left( \eta(0) \sin 2\theta + \int_0^\pi \eta^2(\zeta) d\zeta - K_-(\pi) - 2\omega(\pi) \right) \\
 &\quad + \frac{(k + 1/2)\pi}{n} \frac{(\pi + 2\theta + 2\sigma(\pi))^2}{4n^2\pi^2} + \frac{1}{2n^2} \left( \eta(0) \sin 2\theta + \int_0^x \eta^2(\zeta) d\zeta - K_-(x) \right) \\
 &\quad - \frac{\sigma(x_n^k) + \theta}{2n^3\pi} \left( \eta(0) \sin 2\theta + \int_0^\pi \eta^2(\zeta) d\zeta - K_-(\pi) - 2\omega(\pi) \right) - \\
 &\quad + (\sigma(x_n^k) + \theta) \frac{(\pi + 2\theta + 2\sigma(\pi))^2}{4n^3\pi^2} + O\left(\frac{1}{n^4}\right).
 \end{aligned} \tag{2.10}$$

uniformly with respect to  $k \in \mathbb{Z}^+$ .

**Proof** The first component  $\varphi_1(x, \lambda_n)$  of the eigenfunction  $\varphi(x, \lambda_n)$  has the following asymptotic formula:

$$\begin{aligned}
 \varphi_1(x, \lambda_n) &= \cos(\lambda_n x - \sigma(x) - \theta) + \frac{1}{2\lambda_n} \eta(x) \cos(\lambda_n x - \sigma(x) - \theta) \\
 &\quad - \frac{1}{2\lambda_n} \eta(0) \cos(\lambda_n x - \sigma(x) + \theta) + \frac{1}{2\lambda_n} \sin(\lambda_n x - \sigma(x) - \theta) \int_0^x \eta^2(\zeta) d\zeta \\
 &\quad - \frac{1}{2\lambda_n} K_+(x) \cos(\lambda x - \sigma(x) - \theta) - \frac{1}{2\lambda_n} K_-(x) \sin(\lambda_n x - \sigma(x) - \theta) \\
 &\quad + o\left(\frac{1}{\lambda_n} \exp(|\tau|x)\right),
 \end{aligned} \tag{2.11}$$

for  $n \rightarrow \infty$  uniformly in  $x$ . Then for the nodal points  $x_n^k$  of  $\varphi_1(x, \lambda_n)$ , from  $\varphi_1(x_n^k, \lambda_n) = 0$ , we obtain

$$\begin{aligned}
 &\left( 1 + \frac{1}{2\lambda_n} \eta(x_n^k) - \frac{1}{2\lambda_n} \eta(0) \cos 2\theta - \frac{1}{2\lambda_n} K_+(x_n^k) \right) \tan\left(\lambda_n x - \sigma(x_n^k) - \theta - \frac{\pi}{2}\right) = \\
 &\frac{1}{2\lambda_n} \eta(0) \sin 2\theta + \frac{1}{2\lambda_n} \int_0^{x_n^k} \eta^2(\zeta) d\zeta - \frac{1}{2\lambda_n} K_-(x_n^k) + o\left(\frac{1}{\lambda_n}\right).
 \end{aligned}$$

Taking into account Taylor’s expansions, we get

$$\lambda_n x_n^k - \sigma(x_n^k) - \theta - \frac{\pi}{2} = k\pi + \frac{1}{2\lambda_n} \left( \eta(0) \sin 2\theta + \int_0^{x_n^k} \eta^2(\zeta) d\zeta - K_-(x_n^k) \right) + o\left(\frac{1}{\lambda_n}\right).$$

It follows from the last equality

$$x_n^k = \frac{(k + \frac{1}{2})\pi + \sigma(x_n^k) + \theta}{\lambda_n} + \frac{1}{2\lambda_n^2} \left( \eta(0) \sin 2\theta + \int_0^{x_n^k} \eta^2(\varsigma) d\varsigma - K_-(x_n^k) \right) + o\left(\frac{1}{\lambda_n^2}\right).$$

The relation (2.10) is proven by using the asymptotic formula

$$\begin{aligned} \lambda_n^{-1} &= \frac{1}{n} \left( 1 - \frac{\pi + 2\theta + 2\sigma(\pi)}{2n\pi} - \frac{(\eta(0) \sin 2\theta + \int_0^\pi \eta^2(\varsigma) d\varsigma - K_-(\pi) - 2\omega(\pi))}{2n^2\pi} \right) \\ &\quad \left( \frac{\pi + 2\theta + 2\sigma(\pi)}{2n\pi} \right)^2 + o\left(\frac{1}{n^2}\right) \end{aligned}$$

as  $n \rightarrow \infty$  uniformly in  $k \in \mathbb{Z}^+$ . □

**Theorem 2.3** *Let  $X$  be the set of nodal points. Fix  $x \in (0, \pi)$ . Let a sequence  $\{x_n^k\} \subset X$  be chosen such that  $x_n^k$  converges to  $x$  as  $n \rightarrow \infty$ . Then the following limits exist and finite and corresponding equalities hold:*

$$\lim_{|n| \rightarrow \infty} n \left( x_n^k - \frac{(k + 1/2)\pi}{n} \right) = \sigma(x) + \theta - x \frac{\pi + 2\theta + 2\sigma(\pi)}{2\pi} \triangleq f(x), \tag{2.12}$$

$$\begin{aligned} &\lim_{|n| \rightarrow \infty} 2n^2\pi \left( x_n^k - \frac{(k + 1/2)\pi}{n} + \frac{\sigma(x_n^k) + \theta}{n} \right) \\ &+ \frac{(k + 1/2)\pi (\pi + 2\theta + 2\sigma(\pi))}{2n\pi} \Big) = -(\sigma(x) + \theta) (\pi + 2\theta + 2\sigma(\pi)) \\ &- x \left( \eta(0) \sin 2\theta + \int_0^\pi \eta^2(\varsigma) d\varsigma - K_-(\pi) - 2\omega(\pi) \right) + x \frac{(\pi + 2\theta + 2\sigma(\pi))^2}{2\pi} \\ &+ \pi \left( \eta(0) \sin 2\theta + \int_0^x \eta^2(\varsigma) d\varsigma - K_-(x) \right) \triangleq g(x), \end{aligned} \tag{2.13}$$

and

$$\begin{aligned} &\lim_{|n| \rightarrow \infty} 2n^3\pi \left( x_n^k - \frac{(k + 1/2)\pi}{n} + \frac{\sigma(x_n^k) + \theta}{n} \right) \\ &+ \frac{(k + 1/2)\pi (\pi + 2\theta + 2\sigma(\pi))}{2n\pi} + (\sigma(x_n^k) + \theta) \frac{(\pi + 2\theta + 2\sigma(\pi))}{2n^2\pi} \\ &+ \frac{(k + 1/2)\pi}{n} \frac{1}{2n^2\pi} \left( \eta(0) \sin 2\theta + \int_0^\pi \eta^2(\varsigma) d\varsigma - K_-(\pi) - 2\omega(\pi) \right) \\ &- \frac{(k + 1/2)\pi (\pi + 2\theta + 2\sigma(\pi))^2}{4n^2\pi^2} - \frac{1}{2n^2} \left( \eta(0) \sin 2\theta + \int_0^x \eta^2(\varsigma) d\varsigma - K_-(x) \right) \\ &= (\sigma(x_n^k) + \theta) \left( \eta(0) \sin 2\theta + \int_0^\pi \eta^2(\varsigma) d\varsigma - K_-(\pi) - 2\omega(\pi) \right) \\ &+ (\sigma(x) + \theta) \frac{(\pi + 2\theta + 2\sigma(\pi))^2}{2\pi} \triangleq h(x). \end{aligned} \tag{2.14}$$

Therefore, proof of the following theorem is clear.

**Theorem 2.4** Let  $\sigma(\pi) = 0$ . The given dense subset of nodal points  $X$  uniquely determines the coefficients  $\theta$  and  $\omega(\pi)$  of the boundary conditions and if  $K_-(x)$  is known,  $X$  also uniquely determines the potential  $Q(x)$  a.e. on  $(0, \pi)$ . Moreover,  $Q(x), \omega(\pi)$  and  $\theta$  can be determined via the following algorithm:

(1) For each  $x \in (0, \pi)$  choose a sequence  $\{x_n^k\} \subset X$  such that  $\lim_{|n| \rightarrow \infty} x_n^k = x$ ;

(2) Find the function  $f(x)$  from (2.12) and calculate

$$\begin{aligned} \theta &= f(0) \\ 2\sigma'(x) &= p(x) + r(x) = 2f'(x) + \frac{\pi + 2f(0)}{\pi} \end{aligned}$$

(3) Find the function  $g(x)$  from (2.13) and calculate

$$\begin{aligned} \eta(0) &= \frac{g(0) - \theta\pi - 2\theta^2}{\pi \sin 2\theta} \\ \omega(\pi) &= \frac{g(\pi) - 3\theta\pi - 4\theta^2 - \pi^2/2}{2\pi} \end{aligned}$$

(4) If  $K'_-(x)$  is known then from (2.12)-(2.14) calculate

$$\begin{aligned} p(x) &= f'(x) + \frac{\pi + 2f(0)}{2\pi} + \rho(x) \\ r(x) &= f'(x) + \frac{\pi + 2f(0)}{2\pi} - \rho(x), \end{aligned}$$

where

$$\rho^2(x) = \frac{1}{\pi} \left( g'(x) + \left( f'(x) + f(0) + \frac{\pi}{2} \right) (\pi + 2\theta) + \frac{h(0)}{\theta} + K'_-(x) \right).$$

Thus, we have shown that we can reconstruct the potential function and obtain the coefficients of the boundary conditions using only dense subset of a nodal points. Our reconstruction formulae also directly imply the uniqueness of this inverse problem.

### 3. Conclusion

In this work, we study inverse nodal problems for Dirac type integro-differential systems with one classical boundary condition and another nonlocal integral boundary condition. We get useful asymptotics regarding the solution, eigenvalues, and nodes. And we present a constructive procedure to solve the inverse nodal problems. It makes sense for the theoretical integrity of the inverse nodal problem with nonlocal integral conditions.

### Acknowledgment

The author would like to express his gratitude to the editor and anonymous referees for their helpful comments which significantly improved the quality of the paper.

## References

- [1] Bitsadze AV, Samarskii AA. Some elementary generalizations of linear elliptic boundary value problems. *Doklady Akademii Nauk SSSR* 1969; 185: 739-740.
- [2] Day WA. Extensions of a property of the heat equation to linear thermoelasticity and order theories. *Quarterly of Applied Mathematics* 1982; 40: 319-330.
- [3] Gordeziani N. On some nonlocal problems of the theory of elasticity. *Bulletin of TICMI* 2000; 4: 43-46.
- [4] Yin YF. On nonlinear parabolic equations with nonlocal boundary conditions. *Journal of Mathematical Analysis and Applications* 1994; 185: 161-174. doi: doi.org/10.1006/jmaa.1994.1239
- [5] Nakhushiev AM. *Equations of mathematical biology*. Moscow: Vysshaya Shkola, 1995 (in Russian).
- [6] Schuegerl K. *Bioreaction Engineering. Reactions involving microorganisms and cells*. vol. 1, John Wiley and Sons, 1987.
- [7] Albeverio S, Hryniv R, Nizhnik LP. Inverse spectral problems for nonlocal Sturm-Liouville operators. *Inverse Problems* 2007; 23: 523-535. doi: doi.org/10.1088/0266-5611/23/2/005
- [8] Freiling G, Yurko VA. Inverse problems for differential operators with a constant delay. *Applied Mathematics Letters* 2012; 25: 1999–2004. doi: 10.1016/j.aml.2012.03.026
- [9] Kravchenko KV. On differential operators with nonlocal boundary conditions. *Differentsialnye Uravneniya* 2000; 36: 464–469. (English transl. in *Differential Equations* 2000; 3: 517–523).
- [10] Nizhnik LP. Inverse nonlocal Sturm–Liouville problem. *Inverse Problems* 2010; 26: 125006. doi: 10.1088/0266-5611/26/12/125006
- [11] Bondarenko NP. An inverse problem for an integro-differential operator on a star-shaped graph. *Mathematical Methods in the Applied Sciences* 2018; 41: 1697-1702. doi: 10.1002/mma.4698
- [12] Bondarenko NP. An inverse problem for the integro-differential Dirac system with partial information given on the convolution kernel. *Journal of Inverse and Ill-posed Problems* 2019; 27 (2): 151-157. 10.1515/jiip-2017-0058
- [13] Bondarenko NP, Buterin SA. An inverse spectral problem for integro-differential Dirac operators with general convolution kernels. *Applicable Analysis* 2020; 99: 700-716.
- [14] Bažant ZP, Jirásek M. Nonlocal integral formulation of plasticity and damage: survey of progress, american society of civil engineers. *Journal of Engineering Mechanics* 2002; 1119-1149.
- [15] Buterin SA, Shieh CT. Inverse nodal problem for differential pencils. *Applied Mathematics Letters* 2009; 22: 1240-1247.
- [16] Buterin SA, On an inverse spectral problem for a convolution integro-differential operator. *Results in Mathematics* 2007; 50: 173-181.
- [17] Buterin SA, Yurko VA. Inverse problems for second order integral and integro-differential operators. *Analysis and Mathematical Physics* 2018; 4: 1-10.
- [18] Cheng YH, Law CK, Tsay J. Remarks on a new inverse nodal problem. *Journal of Mathematical Analysis and Applications* 2000; 248: 145-155.
- [19] Çakmak Y. Inverse nodal problem for a conformable fractional diffusion operator. *Inverse Problems in Science and Engineering* 2021; 29 (9): 1308-1322. doi: 10.1080/17415977.2020.1847103
- [20] Freiling G, Yurko VA. *Inverse Sturm-Liouville problems and their applications*. New York: Nova Science, 2001.
- [21] Guo Y, Wei Y. Inverse nodal problem for Dirac equations with boundary conditions polynomially dependent on the spectral parameter. *Results in Mathematics* 2015; 67: 95-110.
- [22] Guo Y, Wei Y, Yao R. Uniqueness theorems for the Dirac operator with eigenparameter boundary conditions and transmission conditions. *Applicable Analysis* 2020; 99 (9): 1564-1578. doi: 10.1080/00036811.2018.1540039



- [23] Hald OH, McLaughlin JR. Solutions of inverse nodal problems. *Inverse Problems* 1989; 5: 307-347.
- [24] Hu YT, Bondarenko NP, Shieh CT, Yang CF. Traces and inverse nodal problems for Dirac-type integro-differential operators on a graph. *Applied Mathematics and Computation* 2019; 363: 124606. doi: 10.1016/j.amc.2019.124606
- [25] Keskin B. Inverse problems for one dimensional conformable fractional Dirac type integro differential system. *Inverse Problems* 2020; 363: 065001. doi: 10.1088/1361-6420/ab7e03
- [26] Keskin B, Ozkan AS. Inverse nodal problems for Dirac-type integro-differential operators. *Journal of Differential Equations* 2017; 263: 8838-8847. doi: 10.1016/j.jde.2017.08.068
- [27] Keskin B, Tel HD. Reconstruction of the Dirac-type integro-differential operator from nodal data. *Numerical Functional Analysis and Optimization* 2018; 39: 1208-1220. doi: 10.1080/01630563.2018.1470097
- [28] Keskin B, Tel HD. Inverse nodal problems for Dirac-type integro-differential system with boundary conditions polynomially dependent on the spectral parameter. *Cumhuriyet Science Journal* 2019; 40 (4): 875-885. doi: 10.17776/csj.620668
- [29] Kuryshova YV, Shieh CT. An Inverse nodal problem for integro-differential operators. *Journal of Inverse and III-posed Problems* 2010; 18: 357-369.
- [30] Kuryshova YV. Inverse spectral problem for integro-differential operators. *Mathematical Notes* 2007; 81 (6): 767-777.
- [31] McLaughlin JR. Inverse spectral theory using nodal points as data – a uniqueness result. *Journal of Differential Equations* 1988; 73: 354-362.
- [32] Lakshmikantham V, Rama Mi Mohana R. *Theory of integro-differential equations Stability and Control: Theory, Methods and Applications*. v.1, Singapore: Gordon and Breach Science Publishers, 1995.
- [33] Law CK, Shen CL, Yang, CF. The Inverse nodal problem on the smoothness of the potential function. *Inverse Problems* 1999; 15 (1): 253-263. (Erratum, *Inverse Problems* 2001; 17: 361-363).
- [34] Şen E. Traces and inverse nodal problems for a class of delay Sturm–Liouville operators. *Turkish Journal of Mathematics* 2021; 45: 305-318. doi: 10.3906/mat-2005-55
- [35] Qin X, Gao Y, Yang C. Inverse nodal problems for the Sturm-Liouville Operator with some nonlocal integral conditions. *Journal of Applied Mathematics and Physics* 2019; 7: 111-122.
- [36] Wang YP, Koyunbakan H, Yang CF. Trace formula for integro-differential operators on the finite interval. *Acta Mathematicae Applicatae Sinica* 2017; 33 (1): 141-146. doi: 10.1007/s10255-017-0644-7
- [37] Yurko VA. An inverse problem for integro-differential operators. *Matematicheskie Zametki* 1991; 50: 134-146. (English transl. in *Mathematical Notes* 1991; 50: 1188-1197).
- [38] Shieh CT, Yurko VA. Inverse nodal and inverse spectral problems for discontinuous boundary value problems. *Journal of Mathematical Analysis and Applications* 2008; 347: 266-272. doi: 10.1016/j.jmaa.2008.05.097
- [39] Yang XF. A solution of the nodal problem. *Inverse Problems* 1997; 13: 203-213.
- [40] Yang CF. Inverse nodal problem for a class of nonlocal Sturm-Liouville operator. *Mathematical Modelling and Analysis* 2010; 15 (3): 383-392. doi: 10.3846/1392-6292.2010.15.383-392
- [41] Yang CF. Trace and inverse problem of a discontinuous Sturm-Liouville operator with retarded argument. *Journal of Mathematical Analysis and Applications* 2012; 395: 30-41.
- [42] Yang CF, Yurko VA. Recovering Dirac operator with nonlocal boundary conditions. *Journal of Mathematical Analysis and Applications* 2016; 440: 155-166.
- [43] Yang CF, Huang ZY. Reconstruction of the Dirac operator from nodal data. *Integral Equation and Operator Theory* 2010; 66: 539-551. doi:10.1007/s00020-010-1763-1
- [44] Yang CF, Pivovarchik VN. Inverse nodal problem for Dirac system with spectral parameter in boundary conditions. *Complex Analysis and Operator Theory* 2013; 7: 1211-1230. doi: 10.1007/s11785-011-0202-x

- [45] Yılmaz E, Koyunbakan H. On the Lipschitz stability of inverse nodal problem for Dirac system. Communications Faculty of Sciences University of Ankara Series A1: Mathematics and Statistics 2021; 70 (1): 341-356. doi: 10.31801/cfsuasmas.733215
- [46] Yılmaz E. Inverse nodal problem for an integro-differential operator. Cankaya University Journal of Science and Engineering 2015; 12 (1): 014-019.