

Szabó's algorithm and applications

Michel Bertrand DJIADEU NGAHA , Salomon Joseph MBATAKOU ,
Romain NIMPA PEFOUKEU* 

Department of Mathematics, Faculty of Science, University of Yaounde 1, Yaounde, Cameroon

Received: 22.09.2022

Accepted/Published Online: 27.02.2023

Final Version: 16.05.2023

Abstract: In this paper, Szabó's algorithm is used as the main tool to find locally symmetric left invariant Riemannian metrics on some 4-dimensional Lie groups. Locally symmetric left invariant Riemannian Lie groups constitute an important subclass of Riemannian Lie groups with zero-divergence Weyl-tensor the so-called C-manifolds. Some properties of the curvature operator of these 4-dimensional C-manifolds are studied.

Key words: Lie algebra, Lie group, locally symmetric metric, Z -decomposition, Weyl tensor

1. Introduction

A left invariant Riemannian metric on a connected Riemannian manifold is locally symmetric if the covariant derivative of its Riemannian curvature tensor vanishes. It is well known that the class of locally symmetric Riemannian metrics contains Riemannian metrics with constant sectional curvature, product of locally symmetric Riemannian metrics, the left invariant Riemannian metrics induced by the opposite of the Killing form on a connected compact semisimple Lie groups and so on. A manifold endowed with a locally symmetric Riemannian metric is a locally symmetric Riemannian space. Cartan in [3, 4] observes that irreducible globally symmetric spaces are homogeneous spaces G/H where G is a connected simple Lie group, H a compact subgroup of G and the Riemannian symmetric metric is the G -invariant Riemannian metric induced by the Killing form of the Lie algebra \mathfrak{g} of G . He used this insight to classify Riemannian symmetric spaces in connection with the theory of Lie groups [9].

A Lie group G together with a left invariant Riemannian metric g is called a Riemannian Lie group. A left invariant Riemannian metric g on G induces an inner product $g(e) = \langle \cdot, \cdot \rangle$ on the Lie algebra \mathfrak{g} of G and conversely, any inner product on \mathfrak{g} gives rise to a unique left invariant Riemannian metric on G .

Let (G, g) be a connected n -dimensional Riemannian Lie group, endowed with Levi-Civita connection ∇ , with Lie algebra \mathfrak{g} . A \mathbb{R} -bilinear function L is called Levi-civita product if

$$\forall u, v \in \mathfrak{g} \quad L_u v := (\nabla_{u^l} v^l)(e),$$

where u^l and v^l are left invariant vector fields associated to u and v . We defined by K the Riemannian curvature tensor at the identity element e by

$$\forall u, v, w \in \mathfrak{g} \quad K(u, v)w = L_{[u, v]}w - L_u L_v w + L_v L_u w.$$

*Correspondence: romain.nimpa@facsociences-uy1.cm

2010 AMS Mathematics Subject Classification: 53C20, 53C35, 53C30

The algebraic condition of local symmetricity on (G, g) is therefore equivalent to

$$[L_{e_m}, K(e_i, e_j)] = K(L_{e_m} e_i, e_j) + K(e_i, L_{e_m} e_j), \tag{1.1}$$

where (e_1, e_2, \dots, e_n) is an $g(e)$ -orthonormal basis of the Lie algebra \mathfrak{g} with respect to the inner product and $[\]$ is the commutator.

The above equations induce a system of $\frac{n^3(n-1)^2}{4}$ polynomial equations on the structure constants of the Lie algebra \mathfrak{g} .

For connected Lie groups of dimension 1 or 2, it follows from a direct computation that every left invariant Riemannian metric is locally symmetric.

For connected Lie groups of dimension 3, locally symmetric left invariant Riemannian metrics are well known, see [14].

For 4-dimensional Riemannian Lie groups, we have from Equation (1.1) a system of 144 polynomial equations for each of the 12 nonisomorphic associated Euclidean Lie algebra. Those systems are not easy to handle. In order to avoid the algebraic condition of local symmetricity for these Lie groups, we use the Szabó's algorithm as the main tool to split any left Riemannian metric of \mathcal{C} -spaces [2] as a direct product of left invariant metrics on Lie groups of lower dimension. In this paper, we are interested in finding locally symmetric left invariant Riemannian metrics on some 4-dimensional connected Lie groups.

The outline of this paper is as follows: Section 2 is devoted to some basic knowledge on the curvature tensor. In Section 3, we describe Szabó's algorithm and recall some main results on locally symmetric Riemannian Lie groups of dimension 3. The Szabó's algorithm is applied to some 4-dimensional Euclidean Lie algebra. In Section 4, we prove Theorem 4.1.

2. Preliminaries

Let (G, g) be an n -dimensional connected Riemannian Lie group and denote by \mathcal{H}^{inf} and \mathcal{H} the infinitesimal holonomy group and the primitive holonomy group at the identity element e of G , respectively. Their Lie algebras are denoted by \mathfrak{h}^{inf} and \mathfrak{h} , respectively. For more details about the holonomy group and his subgroups, see [12].

2.1. The linear curvature operator

Let $x, y, v, w \in \mathfrak{g}$. We recall that on the set $\wedge^2 \mathfrak{g}$ of bivectors of \mathfrak{g} , the inner product denoted $\langle \cdot, \cdot \rangle_{\wedge^2 \mathfrak{g}}$ is defined by

$$\langle x \wedge y, v \wedge w \rangle_{\wedge^2 \mathfrak{g}} = \langle x, v \rangle \langle y, w \rangle - \langle x, w \rangle \langle y, v \rangle. \tag{2.1}$$

Let \overline{K} be the self-adjoint operator called the curvature operator, associated to the symmetric bilinear form K (Riemannian curvature) on the euclidian space $(\wedge^2 \mathfrak{g}, \langle \cdot, \cdot \rangle_{\wedge^2 \mathfrak{g}})$:

$$\langle \overline{K}(x \wedge y), v \wedge w \rangle_{\wedge^2 \mathfrak{g}} = K(x \wedge y, v \wedge w) := \langle K(x, y)v, w \rangle = K(x, y, v, w). \tag{2.2}$$

The relation (2.1) induces a linear isomorphism from the set $\wedge^2 \mathfrak{g}$ of bivectors to the set $\mathfrak{o}(\mathfrak{g})$ of skew-symmetric endomorphisms, such that for $v \wedge w \in \wedge^2 \mathfrak{g}$ and $x \in \mathfrak{g}$,

$$\widetilde{v \wedge w}(x) = \langle v, x \rangle w - \langle w, x \rangle v. \tag{2.3}$$

In the rest of this paper, the skew-symmetric endomorphism associated to a bivector h will be denoted \tilde{h} .

Definition 2.1 [16] Let h be any arbitrary bivector of \mathfrak{g} .

1. The decomposition

$$\mathfrak{g} = U_0 \oplus U_1 \oplus \cdots \oplus U_q, \tag{2.4}$$

where U_0 is the kernel of \tilde{h} and $U_k, 1 \leq k \leq q$ are 2-dimensional invariant real planes of the real operator \tilde{h} , is the Jordan decomposition of the Lie algebra \mathfrak{g} with respect to h .

2. The number q in the Jordan decomposition is the rank of h .

Remark 2.2 $q = \sum m_k$ where m_k is the multiplicity of the nonzero eigenvalue $\lambda_k = i\nu_k, \nu_k > 0$ of \tilde{h} .

Definition 2.3 [16] Let h be any arbitrary bivector. The decomposition

$$h = \sum_{k=1}^q v_k \wedge w_k, \tag{2.5}$$

where $v_k, w_k \in U_k$ and U_k are the 2-dimensional subspaces of the Jordan decomposition is the Darboux decomposition or the Darboux normal form of the bivector h .

Remark 2.4 [16] If the multiplicity m_k of nonzero eigenvalue $\lambda_k = i\nu_k$ of \tilde{h} is greater than one, the Darboux decomposition is not unique.

Definition 2.5 [16] The eigenvector $h \in \wedge^2 \mathfrak{g}$ of the curvature operator \overline{K} is said to be irreducible if any Darboux normal form of h does not split into two nontrivial summands such that they are also eigenvectors of \overline{K} .

Let V_0 be the subspace of \mathfrak{g} such that the action of \mathcal{H} on V_0 is trivial. From [2, 16] the subspace V_0^\perp decomposes into orthogonal, invariant and irreducible subspaces $V_i, i > 1$ under the action of \mathcal{H} such that

$$\mathfrak{g} = V_0 \oplus V_1 \oplus V_2 \oplus \cdots \oplus V_k. \tag{2.6}$$

Here, irreducible means that there are no nontrivial invariant subspace of V_i .

Definition 2.6 [16] The V -decomposition of the Lie algebra \mathfrak{g} of a connected Riemannian Lie group (G, g) is an $g(e)$ -orthogonal and irreducible decomposition of \mathfrak{g} with respect to the primitive holonomy group at $e \in G$.

2.2. The \mathcal{C} -spaces Riemannian Lie groups

Definition 2.7 [7, 15, 17] A Riemannian Lie group (G, g) of dimension ≥ 4 is called a \mathcal{C} -space if its Weyl tensor W satisfies $\text{div}W = 0$.

In the literature, there are some classifications of the metrics Lie algebras of 4-dimensional \mathcal{C} -spaces (see [7, 15, 17] for more details). Table 1 below gives the Lie algebras and the structure constants in an orthonormal basis (e_1, e_2, e_3, e_4) , for 4-dimensional Riemannian Lie groups with respect to their structure constants and the spectrum of the curvature operator.

Table 1. Lie Algebras of 4-dimensional C -spaces riemannian Lie groups.

Lie algebras	Structure constants	$\text{Spec}(\bar{K}) = \{\sigma_{12}, \sigma_{13}, \sigma_{14}, \sigma_{23}, \sigma_{24}, \sigma_{34}\}$
$4A_1$	$C_{i,j}^k = 0, \forall i, j, k,$	$\{0, 0, 0, 0, 0, 0\}$
$A_{3,6} \oplus A_1 c > 0$	$C_{2,3}^1 = c, C_{1,3}^2 = -c$	$\{0, 0, 0, 0, 0, 0\}$
$A_{3,9} \oplus A_1$ $a > 0$	$C_{1,3}^2 = a\sqrt{1+m^2}$ $C_{1,2}^3 = C_{2,3}^1 = -a\sqrt{1+m^2}$ $C_{2,4}^1 = -C_{1,4}^2 = am\sqrt{1+m^2}$	$\left\{ \frac{a^2(1+m^2)}{4}, \frac{a^2(1+m^2)}{4}, 0, \frac{a^2(1+m^2)}{4}, 0, 0 \right\}$
$A_{3,3} \oplus A_1$ $a > 0$	$C_{1,3}^1 = C_{2,3}^2 = a$	$\{-a^2, -a^2, 0, -a^2, 0, 0\}$
$A_{3,7} \oplus A_1$ $a > 0, \alpha > 0$	$C_{1,3}^1 = C_{2,3}^2 = \alpha a$ $C_{2,3}^1 = -C_{1,3}^2 = a$	$\{-\alpha^2 a^2, -\alpha^2 a^2, 0, -\alpha^2 a^2, 0, 0\}$
$A_{4,5}^{\alpha,\beta}$ $a > 0$ $\alpha = \beta = 1$	$C_{1,4}^1 = C_{2,4}^2 = C_{3,4}^3 = a$	$\{-a^2, -a^2, -a^2, -a^2, -a^2, -a^2\}$
$A_{4,6}^{\alpha,\beta}$ $\alpha = \beta,$ $a > 0, \alpha > 0, W=0$	$C_{1,4}^1 = C_{2,4}^2 = C_{3,4}^3 = \alpha a$ $C_{3,4}^2 = -C_{2,4}^3 = a$	$\{-\alpha^2 a^2, -\alpha^2 a^2, -\alpha^2 a^2, -\alpha^2 a^2, -\alpha^2 a^2, -\alpha^2 a^2\}$
$A_{4,12}$ $a > 0, d > 0$	$C_{1,3}^1 = C_{2,3}^2 = \sqrt{a^2 + b^2}$ $C_{2,4}^1 = -C_{1,4}^2 = \frac{ad}{\sqrt{a^2 + b^2}}$ $C_{2,3}^1 = -C_{1,3}^2 = \frac{bd}{\sqrt{a^2 + b^2}}$	$\{-(a^2 + b^2), -(a^2 + b^2), 0, -(a^2 + b^2), 0, 0\}$
$A_2 \oplus 2A_1, a > 0$	$C_{1,2}^2 = a$	$\{-a^2, 0, 0, 0, 0, 0\}$
$2A_2, a > 0, b > 0$	$C_{1,2}^2 = a, C_{3,4}^3 = b$	$\{-a^2, 0, 0, 0, -b^2\}$
$A_{4,5}^{\alpha,\beta}, \alpha\beta \neq 0, -1 \leq \alpha \leq \beta \leq 1, a > 0$	$C_{1,4}^1 = C_{2,4}^2 = C_{3,4}^3 = a$	$\{-a^2, -a^2, -a^2, -a^2, -a^2, -a^2\}$
$A_{4,6}^{\alpha,\beta}, \alpha \neq 0, \beta \geq 0, \alpha \neq 1, a > 0, W \neq 0, \text{div}W = 0$	$C_{1,4}^1 = \alpha a, C_{2,4}^3 = -C_{3,4}^2 = -a$	$\{0, 0, -(\alpha a)^2, 0, 0, 0\}$
$A_{4,6}^{\alpha,\beta}, \alpha \neq 0, \beta > 0, a > 0, W \neq 0, \text{div}W = 0$	$C_{1,4}^1 = C_{2,4}^2 = C_{3,4}^3 = \beta a, C_{2,4}^3 = -C_{3,4}^2 = -a$	$\{-(\beta a)^2, -(\beta a)^2, -(\beta a)^2, -(\beta a)^2, -(\beta a)^2, -(\beta a)^2\}$
$A_{4,9}^{\beta}, -1 < \beta \leq 1, a > 0$	$C_{1,4}^1 = C_{2,3}^2 = 2a, C_{2,4}^3 = C_{3,4}^2 = a$	$\{0, 0, -6a^2, -2a^2, -2a^2, -2a^2\}$
$A_{4,11}^{\alpha}, \alpha > 0, a > 0$	$C_{1,4}^1 = C_{2,3}^2 = 2\alpha a,$ $C_{2,4}^3 = C_{3,4}^2 = \alpha a,$ $C_{3,4}^2 = -C_{2,4}^3 = -a$	$\{0, 0, -6a^2\alpha^2, -2a^2\alpha^2, -2a^2\alpha^2, -2a^2\alpha^2\}$

2.3. Some main results on Riemannian Lie groups of dimension $n \leq 3$

An Euclidian 4-dimensional Lie algebra on which Szabó's algorithm is effective, splits into a direct sum (as linear spaces), of nontrivial Lie algebras both of dimension 2 or dimension 1 and 3. Riemannian Lie groups of dimension 1 and 2 are locally symmetric Riemannian manifolds. The structure constants of the Lie algebra of a 3-dimensional locally symmetric Riemannian Lie group satisfy the following conditions:

Proposition 2.8 [14] *Let G be a connected 3-dimensional real unimodular Lie group with left-invariant Riemannian metric. (G, g) is a locally symmetric Riemannian Lie group if and only if in the Lie algebra \mathfrak{g} of G , there exists an $g(e) = \langle , \rangle$ -orthonormal basis (e_1, e_2, e_3) in which the brackets of the Lie algebra are presented in Table 2.*

Table 2. 3-dimensional unimodular case.

Lie algebra	Structure constants	Restrictions
\mathbb{R}^3	$[e_1, e_2] = [e_2, e_3] = [e_1, e_3] = 0$	
$\mathbb{R}^2 \rtimes \mathfrak{so}(2)$	$[e_1, e_2] = ae_3, [e_3, e_1] = ae_2$	$a > 0$
$\mathfrak{su}(2)$	$[e_1, e_2] = ae_3, [e_2, e_3] = ae_1, [e_3, e_1] = ae_2$	$a > 0$

Proposition 2.9 [14] *Let G be a connected 3-dimensional real nonunimodular Lie group with left-invariant Riemannian metric. (G, g) is a locally symmetric Riemannian Lie group if and only if in the Lie algebra \mathfrak{g} of G , there exist an $g(e) = \langle , \rangle$ -orthonormal basis (e_1, e_2, e_3) in which the brackets of the Lie algebra are presented in Table 3.*

Table 3. 3-dimensional nonunimodular case.

Lie algebra	Structure constants	Restrictions
\mathfrak{g}_I	$[e_1, e_2] = ae_2, [e_1, e_3] = ae_3$	$a > 0$
\mathfrak{g}_D	$[e_1, e_2] = ae_2 + be_3, [e_1, e_3] = -be_2 + ae_3$	$a > 0, b > 0, D \neq 0$
\mathfrak{g}_0	$[e_1, e_2] = ae_2$	$a > 0$

For more details on locally symmetric left invariant 3-dimensional riemannian Lie groups, see [14].

3. Szabó's algorithm

3.1. Szabó's algorithm proof

Let h be an irreducible eigenvector of \overline{K} with nonnull eigenvalue. \tilde{h} leaves the invariant and irreducible subspaces of the orthogonal V -decomposition invariant. Thus, we have the following proposition:

Proposition 3.1 *Let h be an irreducible eigenvector of a nonzero eigenvalue of \overline{K} , the nontrivial invariant subspace*

$$H^1 = U_1 \oplus U_2 \oplus \dots \oplus U_q$$

of \tilde{h} is contained in a unique invariant and irreducible subspace of the V -decomposition.

Proof Let $\mathfrak{g} = U_0 \oplus U_1 \oplus \dots \oplus U_q$ be the Jordan decomposition of \mathfrak{g} with respect to \tilde{h} and $\mathfrak{g} = V_0 \oplus V_1 \oplus \dots \oplus V_k$ a V -decomposition of \mathfrak{g} such that $U_1 \subset V_1$. Suppose that for $j > 1, U_j$ is a subset of one of the subspaces V_i , for some $i > 1$. The equation $\lambda h = \overline{K}(h)$ is equivalent to

$$\lambda v_1 \wedge w_1 - \overline{K}(v_1 \wedge w_1) = -\lambda \sum_{j=2}^q v_j \wedge w_j + \sum_{j=2}^q \overline{K}(v_j \wedge w_j). \tag{3.1}$$

The skew-symmetric linear operator associated to the left hand side and the right hand side of Equation (3.1) satisfies the relation

$$\lambda \widetilde{v_1 \wedge w_1} - K(v_1, w_1) = -\lambda \sum_{j=2}^q \widetilde{v_j \wedge w_j} + \sum_{j=2}^q K(v_j, w_j). \tag{3.2}$$

Let $u = \sum_{s=0}^k u_s$ be the decomposition of u with respect to the V -decomposition. Since $K(u_i, w_i)$ and $\widetilde{u_i \wedge w_i}$ are linear operators on V_i and $K(u_i, w_i)(v_j) = 0$ for all $u_i, w_i \in V_i$ and $v_j \in V_j$ with $i \neq j$, we have

$$K(v_1, w_1)(u) = K(v_1, w_1)(u_1) \in V_1, \quad \lambda \widetilde{v_1 \wedge w_1}(u) \in V_1,$$

$$\lambda \sum_{j=2}^q \widetilde{v_j \wedge w_j}(u) = \lambda \sum_{j=2}^q (\langle v_j, u \rangle w_j - \langle w_j, u \rangle v_j) \in \bigoplus_{j=2}^q V_j$$

$$\text{and } \sum_{j=2}^q K(v_j, w_j)(u) = \sum_{j=2}^q K(v_j, w_j)(u_j) \in \bigoplus_{j=2}^q V_j.$$

For all $u \in \mathfrak{g}$,

$$\left\{ \begin{array}{l} K(v_1, w_1)(u) - \lambda \widetilde{v_1 \wedge w_1}(u) \in V_1 \cap \bigoplus_{j=2}^q V_j = \{0\} \\ \sum_{j=2}^q K(v_j, w_j)(u) - \lambda \sum_{j=2}^q \widetilde{v_j \wedge w_j}(u) \in V_1 \cap \bigoplus_{j=2}^q V_j = \{0\} \end{array} \right. ,$$

thus

$$\left\{ \begin{array}{l} \overline{K}(v_1 \wedge w_1) = \lambda v_1 \wedge w_1 \\ \overline{K}(\sum_{j=2}^q v_j \wedge w_j) = \lambda \sum_{j=2}^q v_j \wedge w_j \end{array} \right. . \tag{3.3}$$

It follows that the eigenvector h splits into two nontrivial sum such that they are also eigenvectors of \overline{K} . This contradicts the irreducibility of h . □

Let $\{h_1, h_2, \dots, h_\rho, h_{\rho+1}, \dots, h_{\frac{n(n-1)}{2}}\}$ be a system of linearly independent irreducible eigenvectors of \overline{K} which form a basis of $\wedge^2 \mathfrak{g}$ and assume that just the first ρ vectors are corresponding to nonzero eigenvalues. Therefore, \mathfrak{h} is the free Lie algebra on $\{\tilde{h}_1, \tilde{h}_2, \dots, \tilde{h}_\rho\}$.

For irreducible bivectors $h_k, 1 \leq k \leq \rho$, let us consider the Jordan decomposition

$$\mathfrak{g} = U_{k0} \oplus U_{k1} \oplus U_{k2} \oplus \dots \oplus U_{kN_k},$$

where U_{k_0} is the kernel of \tilde{h}_k , $U_{kl}, l \in \{1, 2, \dots, N_k\} \subset \mathbb{N}$ the real \tilde{h}_k invariant 2-plan in the Jordan decomposition and N_k the rank of h_k . Let

$$H_k^0 = U_{k_0} \quad \text{and} \quad H_k^1 = U_{k_1} \oplus U_{k_2} \oplus \dots \oplus U_{k_{N_k}}, \tag{3.4}$$

We have

$$\tilde{h}_k H_k^0 = 0, \quad \text{and} \quad \tilde{h}_k H_k^1 \subset H_k^1.$$

Let us choose an arbitrary vector h_{k_1} , $1 \leq k_1 \leq \rho$, and consider its subspaces $H_{k_1}^0$ and $H_{k_1}^1$ constructed above. $H_{k_1}^1$ is a nontrivial linear subspace of a unique V_i , $i > 0$ of the V -decomposition.

Proposition 3.2 *If for any h_i , $i \neq k_1$ the relation $H_{k_1}^1 \subseteq H_i^0$ holds, then $H_{k_1}^1 = V_i$.*

Proof For all real number t , $\exp\left(t \sum_{i=1}^{\rho} \alpha_i \tilde{h}_i\right) = Id + \sum_{i=1}^{\infty} \frac{t^k}{k!} \left(\sum_{i=1}^{\rho} \alpha_i \tilde{h}_i\right)^k \in \mathcal{H}_e$.

By the above, the (\cdot) -action of \mathcal{H} on H_k^1 is completely determined by the action of the generators $\tilde{h}_1, \tilde{h}_2, \dots, \tilde{h}_\rho$ of the free Lie algebra \mathfrak{h}_e on H_k^1 . The image of H_k^1 by $\sum_{i=1}^{\rho} \alpha_i \tilde{h}_i$ is a linear subspace of H_k^1 , written

$$\alpha_1 \tilde{h}_1 \cdot H_k^1 + \alpha_2 \tilde{h}_2 \cdot H_k^1 + \dots + \alpha_k \tilde{h}_k \cdot H_k^1 + \dots + \alpha_\rho \tilde{h}_\rho \cdot H_k^1 \subset H_k^1.$$

Suppose that $\alpha_j \tilde{h}_j \cdot H_k^1$ is trivial with $j \neq k$, i.e. $H_k^1 \subset H_j^0$ for $j \in \{1, 2, \dots, \rho\}$ and $j \neq k$. Then H_k^1 is one of the subspace V_i , $i > 0$ of the V -decomposition. □

Proposition 3.3 *Let h_1 and h_2 be two irreducible eigenvectors with nonzero eigenvalue of \overline{K} . If $H_1^1 \not\subseteq H_2^0$, then $H_1^1 + H_2^1$ is contained in a unique invariant and irreducible subspace of the V -decomposition.*

Proof Let $h_1 = \sum_{i=1}^{N_1} v_i \wedge w_i$ and $h_2 = \sum_{s=1}^{N_2} v'_s \wedge w'_s$ be the Darboux normal form of h_1 and h_2 , respectively.

We suppose that

$$H_1^1 \subset V_k \quad \text{and} \quad H_2^1 \subset V_l \quad \text{with} \quad k \neq l.$$

Let $x = \sum_{i=1}^{N_1} (\alpha_i v_i + \beta_i w_i) \in H_1^1$, x is an element of V_k and we have:

$$\begin{aligned} \tilde{h}_2(x) &= \sum_{s=1}^{N_2} \sum_{i=1}^{N_1} \alpha_i v'_s \wedge w'_s(v_i) + \sum_{s=1}^{N_2} \sum_{i=1}^{N_1} \beta_i v'_s \wedge w'_s(w_i) \\ &= \sum_{s=1}^{N_2} \sum_{i=1}^{N_1} \alpha_i (\langle v'_s, v_i \rangle w'_s - \langle w'_s, v_i \rangle v'_s) + \sum_{s=1}^{N_2} \sum_{i=1}^{N_1} \beta_i (\langle v'_s, w_i \rangle w'_s - \langle w'_s, w_i \rangle v'_s) \\ &= 0. \end{aligned}$$

In fact, $\langle u, v \rangle = 0$ for all $u \in V_k$, $v \in V_l$, since the V -decomposition is an orthogonal decomposition. Thus, $x \in \ker \tilde{h}_2$ and $H_1^1 \subset \ker \tilde{h}_2 = H_2^0$. This contradicts the fact that $H_1^1 \not\subseteq H_2^0$. Therefore $V_k = V_l$. □

Remark 3.4 *The distribution (V_i) of \mathfrak{g} is not always involutive.*

Let $V_i, i > 0$ be a linear subspace given by the V -decomposition. Let us consider for $i > 0$, the subspaces

$$Z_i = \text{Span}\{u_1, L_{u_1}u_2, L_{u_1}L_{u_2}u_3, \dots, L_{u_1}L_{u_2} \dots L_{u_l}u_{l+1}, u_i \in V_i, l \in \mathbb{N} \setminus \{0\}\}$$

of \mathfrak{g} and Z_0 the complete subspace in \mathfrak{g} which is totally orthogonal to the space $Z_1 + Z_2 + \dots + Z_k$.

Remark 3.5 $Z_0 \subset V_0, \quad V_i \subset Z_i \quad \text{and} \quad \mathcal{H}^{inf} Z_i \subset Z_i$.

We have the orthogonal splitting

$$\mathfrak{g} = Z_0 \oplus Z_1 \oplus Z_2 \oplus \dots \oplus Z_k. \tag{3.5}$$

See [16]. It is a decomposition of the Lie algebra \mathfrak{g} into invariant and irreducible subspaces with respect to the infinitesimal holonomy group at e .

Definition 3.6 [16]

The splitting (3.5) is called a Z -decomposition of the Lie algebra \mathfrak{g} of a connected Riemannian Lie group (G, g) .

Proposition 3.7 [16] *The subspaces Z_i induce on G a totally parallel distribution. Thus they are involutive, and the integral manifolds are totally geodesic.*

Remark 3.8

1. *If (G, g) is locally symmetric Riemannian Lie group, then $V_i = Z_i$, since $\mathcal{H}^{inf} = \mathcal{H}$ and V_i is a nonnull \mathcal{H} -invariant subspace of \mathcal{H} -irreducible subspace Z_i ;*
2. *$Z_j, j \geq 0$ is a Lie subalgebra of \mathfrak{g} .*

Proposition 3.9 [1, 5, 10, 12] *If an Euclidean Lie algebra \mathfrak{g} admits a Z -decomposition, then the associated connected Riemannian Lie group (G, g) splits into a direct product of Riemannian immersed Lie subgroups of G .*

3.2. Steps of Szabó's algorithm

Step 1: V -decomposition of the Euclidean Lie algebra \mathfrak{g}

1. i) $\mathcal{H}H_{k_1}^1 \subseteq H_{k_1}^1$ so that $H_{k_1}^1$ is a nontrivial \mathcal{H} -invariant subspace of V_k ;
- ii) \mathcal{H} acts irreducibly on V_k .

Therefore $H_{k_1}^1 = V_k, k > 0$ is one of the subspaces of the V -decomposition.

2. If there exist vectors $h_{k_2}, h_{k_3}, \dots, h_{k_l}$ with $1 \leq k_i \leq \rho$, such that $H_{k_1}^1 \not\subseteq H_{k_i}^0$ holds for each k_i ,

(a) either, if for any $h_i, i \notin \{k_2, k_3, \dots, k_l\}$, the relation $H_{k_1}^1 + H_{k_2}^1 + \dots + H_{k_l}^1 \subseteq H_i^0$ holds. Then

$$H_{k_1}^1 + H_{k_2}^1 + \dots + H_{k_l}^1 \text{ is the subspace } V_k \text{ of the } V\text{-decomposition,}$$

(b) or, there exist $h_i, i \notin \{k_2, k_3, \dots, k_l\}$ such that $H_{k_1}^1 + H_{k_2}^1 + \dots + H_{k_l}^1 \not\subseteq H_i^0$. Therefore, we extend the system $H_{k_1}^1 + H_{k_2}^1 + \dots + H_{k_l}^1$ with the elements H_i^1 .

This process can be repeated for $H_{k_1}^1 + H_{k_2}^1 + \dots + H_{k_l}^1 + H_i^1$ until we obtain the *maximal system* $h_{k_1}, h_{k_2}, \dots, h_{k_j}$ such that:

- For every index $i, 1 \leq i \leq j$, there is another index $i', 1 \leq i' < i$ with $H_{k_{i'}}^1 \not\subseteq H_{k_i}^0$.
- For every index $i \notin \{k_1, k_2, \dots, k_j\}$, we get the relation $H_{k_s}^1 \subseteq H_i^0 \quad s \in \{1, 2, \dots, j\}$.

Then $V_k = H_{k_1}^1 + H_{k_2}^1 + \dots + H_{k_j}^1, k > 0$.

3. By continuing the procedure we can construct all other invariant subspaces $V_k, k > 0$.

Step 2: Z-decomposition of the Euclidean Lie algebra \mathfrak{g}

Construct all the invariant subspaces $Z_k, k \geq 0$.

4. Applications of Szabó’s algorithm

We apply Szabó’s algorithm on the euclidian Lie algebras of 4-dimensional \mathcal{C} - spaces Riemannian Lie groups. Precisely we have the following.

Theorem 4.1 *Let (G, g) be a 4-dimensional connected Riemannian Lie group. If the metric Lie algebra $(\mathfrak{g}, g(e) = \langle, \rangle)$ belongs to Table 4 below, then g is locally symmetric.*

Proof According to the classification table of 4-dimensional Riemannian Lie groups with nonnull harmonic Weyl tensor, see Table 1, we compute the matrix of the curvature tensor in the basis $(e_1 \wedge e_2, e_1 \wedge e_3, e_1 \wedge e_4, e_2 \wedge e_3, e_2 \wedge e_4, e_3 \wedge e_4)$, where (e_1, e_2, e_3, e_4) is an orthonormal basis of \mathfrak{g} .

1. The Lie algebra $\mathbb{A}_2 \oplus 2\mathbb{A}_1$.

For this Lie algebra, the nonnull components of the curvature tensors are:

$$\begin{aligned} K(e_1, e_2)e_1 &= -(a^2 + \frac{3}{4}b^2)e_2, & K(e_1, e_4)e_1 &= \frac{1}{4}b^2e_4, & K(e_2, e_4)e_2 &= \frac{1}{4}b^2e_4, \\ K(e_1, e_2)e_2 &= (a^2 + \frac{3}{4}b^2)e_1, & K(e_1, e_4)e_4 &= -\frac{1}{4}b^2e_1, & K(e_2, e_4)e_4 &= -\frac{1}{4}b^2e_2 \end{aligned} .$$

Linear curvature operator K is $\text{diag}(- (a^2 + \frac{3}{4}b^2), 0, \frac{1}{4}b^2, 0, \frac{1}{4}b^2, 0)$.

Case I : $b \neq 0$

The eigenvectors $h_1 = e_1 \wedge e_2, h_2 = e_1 \wedge e_4, h_3 = e_2 \wedge e_4$ associated respectively to nonnull eigenvalues $-(a^2 + \frac{3}{4}b^2), \frac{1}{4}b^2, \frac{1}{4}b^2$ are irreducible. $H_1^0 = \text{Span}\{e_3, e_4\}, H_1^1 = \text{Span}\{e_1, e_2\}, H_2^0 = \text{Span}\{e_2, e_3\}, H_2^1 = \text{Span}\{e_1, e_4\}, H_3^0 = \text{Span}\{e_1, e_3\}$ and $H_3^1 = \text{Span}\{e_2, e_4\}$.

$H_1^1 \not\subseteq H_2^0$ and $H_1^1 \not\subseteq H_3^0$. The subspace $V_1 = H_1^1 + H_2^1 + H_3^1 = \text{Span}\{e_1, e_2, e_4\}$ is one of subspaces $V_j, j > 0$ and the V -decomposition of $\mathbb{A}_2 \oplus 2\mathbb{A}_1$ is $\mathbb{A}_2 \oplus 2\mathbb{A}_1 = V_0 \oplus V_1$, where $V_0 = \text{Span}\{e_3\}$. The Z -decomposition of $\mathbb{A}_2 \oplus 2\mathbb{A}_1$ with $b \neq 0$ is

$$\mathbb{A}_2 \oplus 2\mathbb{A}_1 = Z_0 \oplus Z_1,$$

Table 4. Szabó's decomposition Lie Algebra.

Lie algebras	Structure constants	Results obtained
$\mathbb{A}_2 \oplus 2\mathbb{A}_1$	$C_{1,2}^2 = a, \quad a > 0$	Szabó's algorithm effective locally symmetric
$2\mathbb{A}_2$	$C_{1,2}^2 = a, \quad C_{3,4}^4 = b$ $a > 0, \quad b > 0$	Szabó's algorithm effective locally symmetric
$\mathbb{A}_{4,6}^{\alpha,\beta}$ Weyl tensor $W \neq 0$ $\alpha \neq 0, \beta \geq 0, \alpha \neq 1, a > 0$	$C_{1,4}^1 = \alpha a, \quad C_{2,4}^3 = -C_{3,4}^2 = -a$	Szabó's algorithm effective locally symmetric
$\mathbb{A}_{3,9} \oplus \mathbb{A}_1$ $a > 0$	$C_{1,3}^2 = a\sqrt{1+m^2}$ $C_{1,2}^3 = C_{2,3}^1 = -a\sqrt{1+m^2}$ $C_{2,4}^1 = -C_{1,4}^2 = am\sqrt{1+m^2}$	Szabó's algorithm effective locally symmetric
$\mathbb{A}_{3,3} \oplus \mathbb{A}_1$ $a > 0$	$C_{1,3}^1 = C_{2,3}^2 = a$	Szabó's algorithm effective locally symmetric
$\mathbb{A}_{3,7}^\alpha \oplus \mathbb{A}_1$ $a > 0, \alpha > 0$	$C_{1,3}^1 = C_{2,3}^2 = \alpha a$ $C_{2,3}^1 = -C_{1,3}^2 = a$	Szabó's algorithm effective locally symmetric
$\mathbb{A}_{4,12}$ $a > 0, d > 0$	$C_{1,3}^1 = C_{2,3}^2 = \sqrt{a^2 + b^2}$ $C_{2,4}^1 = -C_{1,4}^2 = \frac{ad}{\sqrt{a^2 + b^2}}$ $C_{2,3}^1 = -C_{1,3}^2 = \frac{bd}{\sqrt{a^2 + b^2}}$	Szabó's algorithm effective locally symmetric

where $Z_1 = \text{Span}\{e_1, e_2, e_4\}$ and $Z_0 = \text{Span}\{e_3\}$.

For the Lie algebra Z_1 , the Lie brackets are

$$[e_2, e_1] = -ae_1 - be_4, \quad [e_2, e_4] = 0, \quad [e_1, e_4] = 0.$$

Moreover $[Z_0, Z_1] = \{0\}$.

For this decomposition, left invariant Riemannian metric induced by the restriction of the inner product \langle , \rangle on the 3-dimensional Lie algebra Z_1 is not locally symmetric by proposition 2.8 or 2.9.

Therefore if $b \neq 0$, then $\mathbb{A}_2 \oplus 2\mathbb{A}_1$ does not admit locally symmetric metric.

Case II : $b = 0$.

The eigenvector $h_1 = e_1 \wedge e_2$ associated to the nonnull eigenvalue $-a^2$ is irreducible. $H_1^0 = \text{Span}\{e_3, e_4\}$ and $H_1^1 = \text{Span}\{e_1, e_2\}$. The subspace $V_1 = \text{Span}\{e_1, e_2\}$ is one of the subspaces $V_j, j > 0$. The V -decomposition is $\mathbb{A}_2 \oplus 2\mathbb{A}_1 = V_0 \oplus V_1$, where $V_0 = \text{Span}\{e_3, e_4\}$. The Z -decomposition of $\mathbb{A}_2 \oplus 2\mathbb{A}_1$ with $b = 0$ is

$$\mathbb{A}_2 \oplus 2\mathbb{A}_1 = Z_0 \oplus Z_1,$$

where $Z_1 = \text{Span}\{e_1, e_2\}$ and $Z_0 = \text{Span}\{e_3, e_4\}$.

In addition, Z_0 is a 2-dimensional abelian Lie algebra and $[Z_0, Z_1] = \{0\}$.

For this decomposition, left invariant Riemannian metrics induced by the restriction of the inner product \langle , \rangle on the 2-dimensional Lie algebras Z_0 and Z_1 are locally

Therefore if $b = 0$, then $\mathbb{A}_2 \oplus 2\mathbb{A}_1$ admit locally symmetric metric.

2. The Lie algebra $2\mathbb{A}_2$.

The matrix of the curvature operator is $[\overline{K}] = \text{diag}(-a^2, 0, 0, 0, 0, -b^2)$. For the nonnull eigenvalue $-a^2$ and $-b^2$ of $[\overline{K}]$, the associated and irreducible eigenvectors are $h_1 = e_1 \wedge e_2$ and $h_2 = e_3 \wedge e_4$. Therefore $H_1^0 = \text{Span}\{e_3, e_4\}$, $H_1^1 = \text{Span}\{e_1, e_2\}$, $H_2^0 = \text{Span}\{e_1, e_2\}$, and $H_2^1 = \text{Span}\{e_3, e_4\}$.

- i) $H_1^1 \subset H_2^0$; therefore, H_1^1 is one of the subspaces V_j , $j > 0$ in the V -decomposition.
- ii) $H_2^1 \subset H_1^0$; hence, H_2^1 is one of the subspaces V_i , $i > 0$ in the V -decomposition.

Setting $V_1 = H_1^1$ and $V_2 = H_2^1$, the V -decomposition of $2\mathbb{A}_2$ is $2\mathbb{A}_2 = V_1 \oplus V_2$ and the Z -decomposition is

$$2\mathbb{A}_2 = Z_1 \oplus Z_2,$$

where $Z_1 = \text{Span}\{e_1, e_2\}$ and $Z_2 = \text{Span}\{e_3, e_4\}$.

The components Z_1 and Z_2 of the Z -decomposition satisfy $[Z_1, Z_2] = \{0\}$. Moreover, the left invariant Riemannian metrics induced by the restriction on $Z_i, i \in \{1, 2\}$ of the inner product \langle, \rangle are locally symmetric since $\dim Z_1 = \dim Z_2 = 2$. Thus, the metric g is locally symmetric.

3. The Lie algebra $\mathbb{A}_{4,6}^{\alpha,\beta}$, $\alpha \neq 1$.

The nonnull structure constants are $C_{1,4}^1 = \alpha a$, $C_{2,4}^3 = -a$, $C_{3,4}^2 = a$, $a > 0$ and the matrix of the curvature tensor linear operator is $[\overline{K}] = \text{diag}(0, 0, -(\alpha a)^2, 0, 0, 0)$. For the nonnull eigenvalue $-(\alpha a)^2$ of $[\overline{K}]$, the associated irreducible eigenvector is $h_1 = e_1 \wedge e_4$. The Z -decomposition is

$$\mathfrak{g} = Z_0 \oplus Z_1,$$

where $Z_1 = \text{Span}\{e_1, e_4\}$ and $Z_0 = \text{Span}\{e_2, e_3\}$. On the other hand, the left invariant Riemannian metrics induces by the restriction on $Z_i, i \in \{0, 1\}$ of the inner product \langle, \rangle are locally symmetric, since $\dim Z_0 = \dim Z_1 = 2$. Therefore, the metric g is locally symmetric. Also $[Z_0, Z_1] = Z_0$ and Z_0 is an abelian Lie algebra.

4. The Lie algebra $\mathbb{A}_{3,3} \oplus \mathbb{A}_1$:

The matrix of the linear curvature operator \overline{K} is $\text{diag}(-a^2, -a^2, 0, -a^2, 0, 0)$. The eigenvectors $h_1 = e_1 \wedge e_2$, $h_2 = e_1 \wedge e_3$, $h_3 = e_2 \wedge e_3$ associated respectively to nonzero eigenvalues $-a^2$, $-a^2$, $-a^2$ are irreducible. By direct computation, the Z -decomposition of $\mathbb{A}_{3,3} \oplus \mathbb{A}_1$ is

$$\mathbb{A}_{3,3} \oplus \mathbb{A}_1 = Z_0 \oplus Z_1,$$

where $Z_1 = \text{Span}\{e_1, e_2, e_3\}$ $Z_0 = \text{Span}\{e_4\}$.

For the Lie algebra Z_1 , the Lie brackets are

$$[e_3, e_1] = -ae_1 + 0e_2, \quad [e_3, e_2] = 0e_1 - ae_2, \quad \text{and} \quad [e_1, e_2] = 0.$$

Furthermore, $[Z_0, Z_1] = \{0\}$.

The restriction of the inner product \langle , \rangle on Z_1 induces a locally symmetric left invariant Riemannian metric by proposition 2.8. Therefore the inner product \langle , \rangle on $\mathbb{A}_{3,3} \oplus \mathbb{A}_1$ is locally symmetric as product of locally symmetric left invariant metrics.

For other Lie algebras, $\mathbb{A}_{3,9} \oplus \mathbb{A}_1$, $\mathbb{A}_{3,7}^\alpha \oplus \mathbb{A}_1$, and $\mathbb{A}_{4,12}$, the irreducible eigenvectors associated to nonnull eigenvalues are $h_1 = e_1 \wedge e_2$, $h_2 = e_1 \wedge e_3$ and $h_3 = e_2 \wedge e_3$. By direct computation, the Z -decomposition is

$$\mathfrak{g} = Z_0 \oplus Z_1,$$

where $Z_0 = \{e_4\}$ and $Z_1 = \text{Span}\{e_1, e_2, e_3\}$. The Lie brackets on the 3-dimensional components Z_1 of the above Lie algebras are as follows:

(a) For $\mathbb{A}_{3,9} \oplus \mathbb{A}_1$,

$$[e_1, e_2] = -a\sqrt{1+m^2} e_3, \quad [e_2, e_3] = -a\sqrt{1+m^2} e_1, \quad [e_3, e_1] = -a\sqrt{1+m^2} e_2.$$

Moreover, $[Z_0, Z_1] \subset Z_1$.

(b) For $\mathbb{A}_{3,7}^\alpha \oplus \mathbb{A}_1$,

$$[e_3, e_1] = -\alpha a e_1 + a e_2, \quad [e_3, e_2] = -a e_1 - \alpha a e_2, \quad [e_1, e_2] = 0.$$

and it holds that $[Z_0, Z_1] = \{0\}$.

(c) For $\mathbb{A}_{4,12}$,

$$[e_3, e_1] = -\sqrt{a^2 + b^2} e_1 + \frac{bd}{\sqrt{a^2 + b^2}} e_2, \quad [e_3, e_2] = -\frac{bd}{\sqrt{a^2 + b^2}} e_1 - \sqrt{a^2 + b^2} e_2,$$

$$[e_1, e_2] = 0.$$

We also have $[Z_0, Z_1] \subset Z_1$.

For each of these Lie algebras, the left invariant Riemannian metrics induced by the restriction of the inner product \langle , \rangle on Z_1 are locally symmetric by Propositions 2.8 and 2.9. Therefore, the metrics g are locally symmetric.

□

5. Conclusion

We point out a rich and complete important class of examples of 4-dimensionnal locally symmetric Riemannian manifolds: the 4-dimensional locally symmetric Riemannian Lie groups which are decomposable with respect to the infinitesimal holonomy group into a product of locally symmetric Riemannian Lie groups of lower dimension. For 4-dimensionnal metric Lie algebras $\mathbb{A}_{4,5}^{\alpha,\beta}$ ($\alpha\beta \neq 0$, $-1 \leq \alpha \leq \beta \leq 1$), $\mathbb{A}_{4,6}^{\alpha,\beta}$ ($\alpha \neq 0$, $\beta > 0$ and $W \neq 0$), $\mathbb{A}_{3,6} \oplus \mathbb{A}_1$ and $4\mathbb{A}_1$ of C -spaces Riemannian Lie groups, the Szabó's algorithm is not effective. But, the associated left invariant Riemannian metrics are locally symmetric since their sectional curvatures are constants. For the remaining metric Lie algebras $\mathbb{A}_{4,9}^\beta$, $\mathbb{A}_{4,11}^\alpha$ ($\alpha > 0$) and $\mathbb{A}_{3,3} \oplus \mathbb{A}_1$ ($b \neq 0$) of C -spaces Riemannian Lie groups, only the direct computation of algebraic condition of locally symmetricity seems affordable. But this method is very tedious.

References

- [1] Atceken M, Keleş S. On the product Riemannian manifolds. *Differential Geometry - Dynamical Systems*, Balkan Society of Geometers, Geometry Balkan Press 2003; 5 (1): 1-8.
- [2] Besse AL. *Einstein Manifolds*. Springer-Verlag Berlin, Heidelberg Berlin, 1987.
- [3] Cartan E. Sur les classes remarquables d'espaces Riemanniens. *Bulletin de la Société Mathématique Française* 1926; 54 : 214-264 (in French).
- [4] Cartan E. Sur les classes remarquables d'espaces Riemanniens II. *Bulletin de la Société Mathématique Française* 1927; 55: 114-134 (in French).
- [5] Di Scala AJ. Autoparallel Distribution and Splitting Theorems. *TSUKUBA Journal of Mathematics* 2003; 27 (1): 99-102.
- [6] Gladunova OP, Rodionov ED, Slavskii VV. On the Spectrum of Curvature Operator of Conformally Flat Riemannian Manifold. *Doklady Mathematics* 2013; 87 (3): 279-281. <https://doi.org/10.1134/S1064562413030149>.
- [7] Gladunova OP, Slavskii VV. Harmonicity of the Weyl Tensor of Left-Invariant Riemannian Metric on Four-Dimensional Unimodular Lie Groups. *Siberian advances in Mathematics* 2013; 23 (1): 32-46. <https://doi.org/10.3103/S1055134413010033>
- [8] Gladunova OP, Oskorbin DN. An Application of Symbolic Computation Packages to the Investigation of the Curvature Operator Spectrum on the Metric Lie Groups. *The news of Altai State University* 2013; 1: 19-23.
- [9] Helgason S. *Differential Geometry and Symmetric Spaces*. Pure and Applied Mathematics Academic press, New York-London, 1962.
- [10] Kirillov A. *An introduction to Lie groups and Lie algebras*. Cambridge University Press, 2008.
- [11] Klepikov PN, Oskorbin DN. Milnor's Generalized Bases for Some Four-dimensional Metric Lie Algebras. *The news of Altai State University* 2015; (in Russian with an abstract in English) [https://doi.org/10.14258/izvasu\(2015\)1.1-13](https://doi.org/10.14258/izvasu(2015)1.1-13).
- [12] Kobayashi S, Nomizu K. *Foundations of Differential Geometry*. Vol I, Wiley, New York, 1963.
- [13] Milnor J. Curvatures of Left Invariant Metrics on Lie Groups. *Advances in Mathematics* 1976; 21: 293-329.
- [14] Nimpa RP, Djiadeu Ngaha MB, Wouafo Kamga J. Locally symmetric left invariant Riemannian metrics on 3-dimensional Lie groups. *Mathematische Nachrichten*. 2017; 290: 2341-2355. <https://doi.org/10.1002/mana.201600332>
- [15] Rodionov ED, Slavskii VV, Khromova OP. On the Curvature Operator Spectrum of (Half)Conformally Flat Riemannian Metrics. *The news of Altai State University* 2015; 85 (1): 107-112. [https://doi.org/10.14258/izvasu\(2015\)1.1-19](https://doi.org/10.14258/izvasu(2015)1.1-19)
- [16] Szabó ZI. Structure theorems on Riemannian spaces satisfying $R(X, Y).R = 0$. I. the local version. *Journal of Differential Geometry* 1982; 17 (4): 531-582. <https://doi.org/10.4310/jdg/1214437486>
- [17] Voronov DS, Rodionov ED. Left-Invariant Riemannian Metrics on Four-Dimensional Nonunimodular Lie Groups with Zero-Divergence Weyl Tensor. *Doklady Mathematics* 2010; 81 (3): 392-394. <https://doi.org/10.1134/S1064562410030154>