

## Dimensions of water contamination in the subprovinces of Adana province, Turkey

Ferdi TANIR<sup>1</sup>, Beril AKÇİMEN<sup>2</sup>, Fatih KÖKSAL<sup>2</sup>, Necdet AYTAÇ<sup>1</sup>, İbrahim CENGİZLER<sup>3</sup>

**Aim:** To determine the dimension of the contamination of the drinking and utility water in the touristic Bahçe district of the Karataş subprovince of Adana province, and to share our views on how to raise the quality of the water in the district.

**Materials and methods:** In 2007 and 2008, 80 samples were taken 8 times, on a seasonal basis, from the drinking water and sea water of the Bahçe district. Eleven organic and inorganic parameters were measured and presented following a comparison with criteria values

**Results:** Ni, Zn, Cu, Pb, NO<sub>3</sub>-N, COD, and phenol contamination parameters were found to be high in the stations that provide drinking water and utility water for the Bahçe district.

**Conclusion:** This level of contamination was attributed to the extensive use of natural and artificial fertilizers and pesticide, agricultural contamination, and the fact that polluters were not at a safe distance from the stations and were particularly hazardous types of polluters. It is of the utmost importance that training be provided in order to ensure use of fertilizers and pesticides in standard amounts and at standard times. The training should be supported with warnings and inspections.

**Key words:** Water pollution, heavy metals, pesticides, contamination

### Adana ili kırsalında su kirliliğinin boyutları

**Amaç:** Bu araştırmanın amacı, Adana ili Karataş ilçesi turistik Bahçe beldesindeki içme ve kullanma sularındaki kirliliğin boyutlarını saptamak ve su kalitesini artırma konusundaki görüşlerimizi paylaşmaktır.

**Yöntem ve gereç:** 2007 ve 2008 yıllarında sezonluk dönemlerle sekiz kezde içme suyu ve denizden alınan 80 su örneği alındı. Ölçülen organik ve inorganik 11 parametrenin ölçümleri standart değerlerle karşılaştırıldı.

**Bulgular:** Ni, Zn, Cu, Pb, NO<sub>3</sub>-N, kimyasal oksijen ihtiyacı ile fenol değerleri, içme ve kullanma sularında yüksek bulundu.

**Sonuç:** Bu yüksek değerler atıklara ve tarımsal ilaçlara atfedildi. Belli miktarlarda ve zamanlarda zirai ilaçların kullanılması konusunda eğitim en önemli hedefdir. Bu eğitimler, gözlem ve uyarılarla desteklenmelidir.

**Anahtar sözcükler:** Su kirliliği, ağır metaller, tarım ilaçları, bulaşma

Received: 29.12.2009 – Accepted: 02.07.2010

<sup>1</sup> Department of Public Health, Faculty of Medicine, Çukurova University, Adana - TURKEY

<sup>2</sup> Department of Microbiology, Faculty of Medicine, Çukurova University, Adana - TURKEY

<sup>3</sup> Faculty of Fisheries, Çukurova University, Adana - TURKEY

**Correspondence:** Ferdi TANIR, Department of Public Health, Faculty of Medicine, Çukurova University, Adana - TURKEY

E-mail: ftanir@cu.edu.tr

## Introduction

Adana province is one of the most important provinces of the ancient Cilicia region. Many species that are about to become extinct in other parts of the world still exist in marshy places, fish garths, and lagoons on the Mediterranean coast of Adana (approximately 80 km), a region that has rich nature with endemic plant varieties. The primary source of income in the Bahçe district of Karataş subprovince, 47 km from Adana, is tourism. There is 1 hotel with 110 beds and a public resort for 5000 people, and there are summer houses with 1654 residents. The permanent population of Bahçe district is 2800, and the temporary population is approximately 10,000 people (1-3).

Drinking water is obtained by collecting the water extracted from deep bores of 20-40 m, chlorinating it in a depot, and distributing it through networks. In the center of the district, the water extracted from 3 active wells is stored in 1 depot and transported to the utility areas through network chlorination. While 79% of the population uses water from this network, approximately 15% lack sufficient water (4). Only 20% of the community has treated water, and 80% lacks healthy drinking water (5,6).

This study aimed to determine the dimension of contamination of the drinking and utility water in the touristic Bahçe district of Karataş subprovince in Adana province and to share suggestions on raising the quality of the water in the district.

## Materials and methods

Drinking water samples were taken from the summer houses, the hotel, and the settlements in Bahçe district (DWS-1, DWS-2, DWS-3), and seawater samples were taken from 2 stations on the touristic coast (SS-1, SS-2). Samples were taken 80 times in the course of 2 years, 10 times a month, twice each in January, April, July, and October in 2007 and 2008. The usage of pesticides and fertilizers was most commonly observed from April to September. The middle month (i.e. the second month) of each season was selected to determine the change in water pollution in relation to seasons and agricultural activities. This periodic sampling was intended to demonstrate the seasonal and agricultural differences among the months (5,7). These samples were analyzed

and recorded. Chemical water and bacteriological water samples, which were taken in accordance with the regulations, were sent to the analysis center within 2 h. The samples were analyzed with a UV-Vis spectrophotometer for organic and inorganic contamination at the Work and Occupational Health Laboratory of the Department of Public Health in the Medical Faculty of Çukurova University.

Furthermore, bacteriological contamination was tested by using cultures in the Bacteriology Laboratories of the Department of Microbiology. Only 2 organic parameters of water contamination and 9 inorganic parameters were tested; other organic and inorganic parameters could not be measured due to the insufficient amount of chemicals (4).

Chemical oxygen demand (COD), phenol, inorganic water contamination, nickel (Ni), cyanide (CN), zinc (Zn), copper (Cu), lead (Pb), total chromium (T-Cr), cadmium (Cd), nitrate-nitrogen ( $\text{NO}_3\text{-N}$ ), and total phosphorus (T-P), which are among the criteria of organic water contamination, were measured (mg/L). Contamination measurements were conducted by using cultures of fecal coliforms, EMS/100 mL, and other reproducing bacteria (8). A team, comprising 2 physicians and 3 technicians, collected and analyzed the samples. The measurements were evaluated on the basis of the criteria set out in "Water Contamination Control Regulation" and "Turkish Standards Institute-TSI" (6). The findings were analyzed with variance analyses through repetitive measurements via the SPSS statistical program and presented.

## Results

- It was found that the measurements of Ni, Cu,  $\text{NO}_3\text{-N}$ , T-P, phenol, and COD were higher than the normal values for both years on the basis of the annual average values (Table 1, Figures 1-10).
- It was found that the Pb values measured at the SS-1 station were normal for both years and higher than the standard values at the other stations (Table 1).
- It was found that all measurements of seawater from SS-1 and SS-2 were higher than normal values, although the Zn measurements at DWS-1, DWS-2, and DWS-3 were at normal levels for both years (Table 1).




Figure 1. Seasonal distribution of Ni measurements according to the Wells in 2007.




Figure 2. Seasonal distribution of Ni measurements according to the Wells in 2008.




Figure 3. Seasonal distribution of NO<sub>3</sub>-N measurements according to the Wells in 2007.




Figure 4. Seasonal distribution of NO<sub>3</sub>-N measurements according to the Wells in 2008.




Figure 5. Seasonal distribution of T-P measurements according to the Wells in 2007.




Figure 6. Seasonal distribution of T-P measurements according to the Wells in 2008.




Figure 7. Seasonal distribution of Phenol measurements according to the Wells in 2007.




Figure 8. Seasonal distribution of Phenol measurements according to the Wells in 2008.




Figure 9. Seasonal distribution of COD measurements according to the Well in 2007.




Figure 10. Seasonal distribution of COD measurements according to the Well in 2008.

- During the T-Cr measurements, only the average value of 2008 at the DWS-2 station was higher than the normal values, and the average values at the other stations were normal for both years (Table 1).
- Measurements of CN and Cd, which are inorganic chemicals, were at normal levels for both years at all stations. These chemicals were not subject to statistical analyses as they were at constant levels (Table 1).
- In the seasonal distribution of measurements of the samples from the stations, it was seen that the Zn level was normal at DWS-1, DWS-2, and DWS-

- 3, and above normal at SS-1 and SS-2; T-Cr was normal at all stations; and other chemicals (Ni, Cu, Pb, NO<sub>3</sub>-N, T-P, phenol, and COD) were above normal at all stations (Table 2).
- In the spring, T-Cr was above normal only at DWS-2 and at normal levels at all other stations; other chemicals (Ni, Zn, Cu, Pb, NO<sub>3</sub>-N, T-P, phenol, and COD) were above normal at all other stations (Table 2).
- In the summer, T-Cr was above normal at DWS-2 and normal at all other stations, Zn was normal only at DWS-1, and all other chemicals (Ni, Cu,

Table 1. Annual average measures of chemicals in water, by stations.

Stations	Years	Measurements of chemicals (mg/L) and P-values										
		Ni P = 0.1	CN	Zn P < 0.001	Cu P < 0.001	Pb P < 0.001	T-Cr P < 0.001	Cd	NO <sub>3</sub> -N P < 0.001	T-P P < 0.001	Phenol P < 0.001	COD P < 0.001
DWS-1	2007	<b>0.12</b>	<0.01	0.32	<b>0.51</b>	<b>0.15</b>	0.02	<0.01	<b>0.35</b>	<b>1.35</b>	<b>0.12</b>	<b>123</b>
	2008	<b>0.14</b>	<0.01	0.36	<b>0.92</b>	<b>0.13</b>	0.02	<0.01	<b>0.55</b>	<b>1.52</b>	<b>0.22</b>	<b>154</b>
	Av.	<b>0.13</b>	<0.01	0.34	<b>0.72</b>	<b>0.14</b>	0.02	<0.01	<b>0.45</b>	<b>1.44</b>	<b>0.17</b>	<b>138</b>
DWS-2	2007	<b>0.16</b>	<0.01	0.40	<b>0.69</b>	<b>0.14</b>	0.02	<0.01	<b>0.47</b>	<b>1.62</b>	<b>2.20</b>	<b>156</b>
	2008	<b>0.19</b>	<0.01	0.38	<b>0.91</b>	<b>0.16</b>	<b>0.12</b>	<0.01	<b>0.48</b>	<b>2.20</b>	<b>2.80</b>	<b>194</b>
	Av.	<b>0.18</b>	<0.01	0.39	<b>0.80</b>	<b>0.15</b>	0.07	<0.01	<b>0.48</b>	<b>1.96</b>	<b>2.50</b>	<b>175</b>
DWS-3	2007	<b>0.18</b>	<0.01	0.35	<b>0.66</b>	<b>0.23</b>	0.02	<0.01	<b>0.47</b>	<b>3.90</b>	<b>1.55</b>	<b>199</b>
	2008	<b>0.20</b>	<0.01	0.37	<b>0.98</b>	<b>0.34</b>	0.03	<0.01	<b>0.68</b>	<b>4.30</b>	<b>1.78</b>	<b>235</b>
	Av.	<b>0.19</b>	<0.01	0.36	<b>0.82</b>	<b>0.28</b>	0.03	<0.01	<b>0.58</b>	<b>4.10</b>	<b>1.67</b>	<b>217</b>
SS-1	2007	<b>1.72</b>	<0.01	<b>3.03</b>	<b>0.39</b>	0.07	0.03	<0.01	<b>0.19</b>	<b>2.86</b>	<b>4.12</b>	<b>3524</b>
	2008	<b>1.34</b>	<0.01	<b>2.68</b>	<b>0.42</b>	0.08	0.03	<0.01	<b>0.24</b>	<b>2.65</b>	<b>4.16</b>	<b>4622</b>
	Av.	<b>1.53</b>	<0.01	<b>2.86</b>	<b>0.41</b>	0.08	0.03	<0.01	<b>0.22</b>	<b>2.76</b>	<b>4.14</b>	<b>4073</b>
SS-2	2007	<b>1.94</b>	<0.01	<b>2.31</b>	<b>1.01</b>	<b>0.26</b>	0.04	<0.01	<b>0.18</b>	<b>3.31</b>	<b>5.07</b>	<b>4432</b>
	2008	<b>2.05</b>	<0.01	<b>3.30</b>	<b>0.96</b>	<b>0.27</b>	0.05	<0.01	<b>0.19</b>	<b>3.42</b>	<b>6.11</b>	<b>4615</b>
	Av.	<b>2.00</b>	<0.01	<b>2.81</b>	<b>0.99</b>	<b>0.27</b>	0.05	<0.01	<b>0.19</b>	<b>3.37</b>	<b>5.60</b>	<b>4524</b>

**Abbreviations:** Drinking water station: DWS, sea station: SS, nickel: Ni, cyanide: CN, zinc: Zn, copper: Cu, lead: Pb, total chromium: T-Cr, cadmium: Cd, nitrogen-nitrate: NO<sub>3</sub>-N, total phosphorus: T-P, chemical oxygen demand: COD, average: Av.

**Note:** Values above standard were highlighted in bold. As all measurements of CN and Cd were the same, analysis could not be performed.

- Pb, NO<sub>3</sub>-N, T-P, phenol, and COD) were above normal at all other stations (Table 2).
- In the autumn, Zn was normal at DWS-1 and DWS-3 and above normal at other stations, T-Cr was above normal at DWS-2, and other chemicals (Ni, Cu, Pb, NO<sub>3</sub>-N, T-P, phenol, and COD) were above normal at all other stations (Table 2).
  - During the analysis of all organic and inorganic chemicals in the water samples according to seasons, a significant difference was found between measurements. This difference was more obvious in the increase in the summer months, and it was at the lowest level in the winter months (Table 2, P < 0.001).
  - The differences between the measurements of organic and inorganic chemicals in the water samples on the basis of the seasons were found to be significant (Table 2, P < 0.001).
  - The differences in the water samples between the years on the basis of seasons were found to be significant, at P < 0.001 for Ni, T-Cr, NO<sub>3</sub>-N, and COD, and at P = 0.016 for T-P. It was found to be insignificant for Zn (P = 0.48), Pb (P = 0.72), and phenol (P = 0.313) (Table 2).
  - The difference between the stations in terms of seasons within a year was significant at P < 0.001 for Ni, Pb, T-Cr, NO<sub>3</sub>-N, T-P, phenol, and COD, and insignificant for Zn (P = 0.052) and Cu (P = 0.108) (Table 2).
  - In analysis of the differences between stations irrespective of the year and season, no significant difference was found in the water samples taken from DWS-1, DWS-2, and DWS-3 for Ni and Zn (P = 1.0). However, the difference between the drinking water (DWS-1, DWS-2, DWS-3) and seawater (SS-1 and SS-2) was found to be

Table 2. Annual average measures of chemicals in water, by season.

Seasons	Stations	Measurements of chemicals (mg/L)										
		Ni P < 0.001	CN	Zn P < 0.001	Cu P = 0.01	Pb P < 0.001	T-Cr P < 0.001	Cd	NO <sub>3</sub> -N P < 0.001	T-P P < 0.001	Phenol P < 0.001	COD P < 0.001
Winter	DWS-1	0.11	<0.01	0.28	0.52	0.13	0.02	<0.01	0.38	1.33	0.12	122
	DWS-2	0.16	<0.01	0.75	0.73	0.15	0.05	<0.01	0.41	1.81	2.08	152
	DWS-3	0.18	<0.01	0.79	0.81	0.25	0.02	<0.01	0.54	3.95	1.51	204
	SS-1	1.61	<0.01	2.78	0.31	0.05	0.02	<0.01	0.18	2.66	4.09	3656
	SS-2	1.90	<0.01	2.71	1.01	0.17	0.04	<0.01	0.16	3.17	5.06	4330
Spring	DWS-1	0.13	<0.01	0.32	0.74	0.12	0.02	<0.01	0.42	1.41	0.13	135
	DWS-2	0.18	<0.01	0.85	0.84	0.13	0.07	<0.01	0.46	1.90	2.56	166
	DWS-3	0.19	<0.01	0.79	0.79	0.30	0.04	<0.01	0.55	3.99	1.62	213
	SS-1	1.49	<0.01	2.84	0.38	0.06	0.04	<0.01	0.24	2.70	4.16	3874
	SS-2	1.95	<0.01	2.82	1.02	0.28	0.05	<0.01	0.18	3.35	5.61	4435
Summer	DWS-1	0.14	<0.01	0.38	0.85	0.16	0.02	<0.01	0.56	1.57	0.23	156
	DWS-2	0.19	<0.01	0.80	0.98	0.17	0.08	<0.01	0.54	2.09	2.87	199
	DWS-3	0.20	<0.01	0.42	0.84	0.35	0.04	<0.01	0.61	4.35	1.85	229
	SS-1	1.54	<0.01	2.96	0.48	0.12	0.03	<0.01	0.25	2.87	4.20	4657
	SS-2	2.12	<0.01	2.90	1.01	0.32	0.05	<0.01	0.22	3.52	6.01	4842
Autumn	DWS-1	0.14	<0.01	0.36	0.76	0.14	0.02	<0.01	0.44	1.45	0.21	139
	DWS-2	0.19	<0.01	0.79	0.65	0.16	0.07	<0.01	0.51	2.03	2.48	184
	DWS-3	0.20	<0.01	0.32	0.82	0.22	0.02	<0.01	0.60	4.12	1.70	222
	SS-1	1.48	<0.01	2.85	0.46	0.09	0.03	<0.01	0.21	2.80	4.12	4105
	SS-2	2.02	<0.01	2.82	0.91	0.31	0.05	<0.01	0.19	3.44	5.73	4490

**Abbreviations:** Drinking water station: DWS, sea station: SS, nickel: Ni, cyanide: CN, zinc: Zn, copper: Cu, lead: Pb, total chromium: T-Cr, cadmium: Cd, nitrogen-nitrate: NO<sub>3</sub>-N, total phosphorus: T-P, chemical oxygen demand: COD, average: Av.

**Note:** Values above standard were highlighted in bold. As all measurements of CN and Cd were the same, analysis could not be performed.

significant ( $P < 0.001$ ). Measurement differences were found to be significant ( $P < 0.001$ ) in terms of Pb, Cu, NO<sub>3</sub>-N, T-P, phenol, and COD. It was also found that the difference ( $P = 0.328$ ) between the T-Cr values measured at DWS-1 and DWS-3 and

the difference ( $P = 0.428$ ) between the Cu values at DWS-2 and DWS-3 were insignificant, while the difference between the values measured at the other stations was significant ( $P < 0.001$ , Tables 1 and 2).

Table 3. Annual average bacteriological measures in water, according to stations.

Bacteriological Parameter (mg/L)	DWS-1* 2007	DWS-1* 2008	DWS-2* 2007	DWS-2* 2008	DWS-3* 2007	DWS-3* 2008	SS-1* 2007	SS-1* 2008	SS-2* 2007	SS-2* 2008
Fecal Coliform (EMS/100 mL)	0	0	0	0	0	0	600	600	600	600
<i>Pseudomonas aeruginosa</i> (in 250 mL)	<b>1500</b>	0	<b>1500</b>	0	<b>1500</b>	0	0	<b>2500</b>	0	<b>2500</b>

\*Water contamination control regulation (5).

Note: Values above standard were highlighted in bold.

- The difference between the measured averages of 2007 and 2008 irrespective of station and season was insignificant only for Ni ( $P = 0.1$ ) and significant for all other chemicals. The value in 2008 was the highest one ( $P < 0.001$ , Table 1 and Figures 1 and 2).
- During the analysis of the bacteriological contamination in the samples, it was detected that the fecal coliform values measured in 2007 were at normal levels for DWS-1, DWS-2, and DWS-3, and at the level of contamination at the other stations and at other times. *Pseudomonas aeruginosa* (in 250 mL) was detected in the drinking water samples taken from DWS-1, DWS-2, and DWS-3 in 2007 and the seawater taken from SS-1 and SS-2 in 2008. This contamination was considered to be attributable to the location of the sewerage system, which was nearer to the stations than the normal distance (Table 3).

## Discussion

Ammonium nitrate ( $\text{NH}_4\text{NO}_3$ ), phosphorus pentoxide ( $\text{P}_4\text{O}_{10}$ ), and potassium sulfate ( $\text{K}_2\text{SO}_4$ ) are the most commonly used fertilizer components in the Adana region of Turkey, which is also the area most exposed to pesticides in Turkey, at a rate of 40%. The most preferred pesticides are methamidophos, parathion-methyl, dichlorvos (DDVP), azinphos-methyl, chlorpyrifos-ethyl, and carbaryl. This high usage rate of both fertilizers and pesticides in the agricultural areas of Adana province has been reported to cause water, soil, air, and food contamination (5,7,9,10).

The contamination at the drinking water stations, very clear at DWS-2 and particularly in the agricultural areas in our study, is in line with previous studies, and thus previous studies support our findings (7,9,11-13).

The contamination level, which was above the contamination parameters in the district, can be attributed to the unconscious use of natural and/or artificial fertilizers in the subprovince and the location of water wells nearer to the sources of pollution than the distances set out in regulations. The findings of studies conducted in Bornova in 1981, Bursa in 1981, and Çukurova in 1979, which reported fertilizer and pesticide contamination in underground waters, and the similar findings of some other studies in the literature are supportive of our findings (10,14-16).

Detection of herbicide-type pesticides used for cultivated fields on the surface and in underground water, such as garden soil and water and river water, was found in a study conducted in the Salamanca and Zamora regions of Spain (17). Detection of contamination of insecticide-type pesticides with organophosphate in the water of rivers and creeks in a study conducted on water louse in Tokyo (18), detection of phosphorus and nitrogen in the watershed of Pennsylvania (19), and the reporting of high levels of Zn, Cu, and Pb concentrations in the riverbed of the Guadiamar in Spain (20) all support our findings, but the high level of Cd in the same place was defined as a different finding. In a contamination study conducted on *Cyperus alternifolius* and *Villarsia exaltata* plants living in water, Zn and Pb values similar to ours were found. However, that study was different than ours as the presence of Al and Cd was also reported (21).

## Conclusions

The distance of one drinking water station (33.3%) from a stable and the distance of one (33.3%) from fields and/or gardens meant that 2 drinking water stations (66.6%) were below the criteria values. It was reported that the use of natural and/or artificial fertilizers according to the types of soil in the district was much higher than the recommended level. It was found that the content of fertilizers and pesticides included inorganic substances with values surpassing the criteria values.

In light of these observations, it was concluded that the high level of organic and inorganic parameters in the drinking and utility water of Bahçe district is attributable to the excessive use of

natural and artificial fertilizers, causing agricultural contamination, and the fact that the distance of the pollution sources from the water stations is less than is allowed, as well as the structure of these pollutants (4,8,9,14,22). It is of the utmost importance that people be trained in order to ensure that fertilizers and pesticides are used in the amounts and at the periods and places advised by the relevant authorities. Furthermore, warnings should be given and inspections should be conducted in order to ensure compliance. Also needed is coordination among formal and informal education and training institutions, health organizations, and agricultural institutions to raise environmental consciousness in each section of the public.

## References

1. Culture and Tourism Directorate of Adana Province (Adana Valiliği İl Turizm Müdürlüğü), Adana Province, Adana, Turkey (in Turkish); 2007.
2. Environment and Forest Directorate of Adana Province (Adana Çevre ve Orman Müdürlüğü). Provincial environment report of Adana Province (İl çevre durum raporu), Adana (in Turkish); 2007.
3. Data of 2006, Turkish Statistical Institute of the Prime Ministry of the Republic of Turkey (T.C. Başbakanlık TÜİK - 2006 Verileri), Ankara (in Turkish); 2007.
4. Güler Ç, Çobanoğlu Z. Water pollution (Su kirliliği). The Ministry of Health of Turkey, Environmental Health Publication (T.C. Sağlık Bakanlığı Çevre Sağlığı Yayını), No. 12, (1), Ankara (in Turkish); 1994.
5. Regulations for control of water pollution of the Republic of Turkey's Ministry of Environment and Forestry (T.C. Çevre Bakanlığı Mevzuatı. Su Kirliliği Kontrolü Yönetmeliği), Ankara (in Turkish); 2004.
6. Turkish Standards Institution. Waters intended for human consumption (İnsani tüketim amaçlı sular), TS 266, ICS 13.060.20, Ankara (in Turkish); 2005.
7. Water pollution with heavy metals (Ağır Metallerle Su Kirliliği) (in Turkish). www.yunus.hacettepe.edu.tr/water pollution/22; 2009 (accessed 12 February 2009).
8. Meinhardt PL. Water quality management and water-borne disease trends. In: Maxcy-Rosenau-Last Public Health and Preventive Medicine. 15th ed. 2007. p.863-900.
9. Uslu O. Water pollution: Environmental problems of Turkey '95 (Su Kirliliği: Türkiye'nin Çevre Sorunları '95). The Environment Foundation of Turkey (Türkiye Çevre Vakfı), No. 100, Ankara (in Turkish); 1995. p.91-301.
10. Environmental problems of the water in Dicle river basin (Dicle Havzasında Su İle İlgili Çevre Sorunları), www.gap.gov.tr/diclebasin.2; 2008 (accessed 10 October 2008).
11. Tanır F, Göksu Lugal MZ, Akbaba M. Chemical and bacteriological evaluation of drinking water in Adana province and its districts. Asian Journal of Chemistry 2005; 17: 2443-50.
12. Halkman AK, Atamer M, Ertaş AH. Relationships of industry and environment (Endüstri ve Çevre İlişkileri). II. The Chamber of Agricultural Engineers (Ziraat Müh. Od.). Turkey, Fifth Technical Congress Book (5. Teknik Kongresi Bildiri Kitabı), Ankara (in Turkish); 2001. p.102-108.
13. Buğdaycı R. Bacteriological, physical and chemical examinations of well waters in Doğanekent region (Doğanekent S.E.A. Bölgesinde Kuyu Sularının Fiziksel, Kimyasal ve Bakteriolojik İncelemesi). Expertise thesis (Uzmanlık Tezi), Adana, (in Turkish); 1992. p.77.
14. Moses M. Pesticides. In: Maxcy-Rosenau-Last Public Health and Preventive Medicine, 15th ed. 2007. p.707-724.
15. Akbaba M, Tanır F, Karaömerlioğlu Ö. Chemical evaluations of drinking water in Doğanekent region of Adana province (Adana Doğanekent Beldesi Su Kuyularında Bazı Kimyasal Parametrelerin Ölçüm Değerleri). Fifth National Public Health Congress Book (V. Ulusal Halk Sağlığı Kongresi Bildiri Kitabı), İstanbul (in Turkish); 1996. p.465.
16. Akbaba M, Apan E, Tanır F. Chemical evaluations of drinking water of Seyhan region of Adana province (Adana Seyhan İlçesi Su Kuyularında Kirlilik Göstergesi Olan Bazı Kimyasal Parametrelerin Ölçüm Değerleri). Erciyes Medical Journal 1998; 20: 154-59 (in Turkish).



17. Martinez RC, Gonzalo ER, Laespada EF, San Roman FJS. Evaluation of surface- and ground-water pollution due to herbicides in agricultural areas of Zamora and Salamanca (Spain). *Journal of Chromatography A* 2000; 869: 471-480.
18. Kikuchi M, Sasaki Y, Wakabayashi M. Screening of organophosphate insecticide pollution in water by using *Daphnia magna*. *Ecotoxicology and Environmental Safety* 2000; 47: 239-245.
19. Heathwaite L, Sharpley A, Gburek W. A conceptual approach for integrating phosphorus and nitrogen management at watershed scales. *J Environ Qual* 2000; 29: 158-166.
20. Alonso E, Santos A, Callejon M, Jimenez JC. Speciation as a screening tool for the determination of heavy metal surface water pollution in the Guadiana river basin. *Chemosphere* 2004; 56: 561-570.
21. Shuiping CS, Grosse W, Karrenbrock F, Thoennessen M. Efficiency of constructed wetlands in decontamination of water polluted by heavy metals. *Ecological Engineering* 2002; 18: 317-325.
22. Ayaz SC, Akca L. Treatment of wastewater by natural systems. *Environment International* 2001; 263: 189-195.