

Cholesterol efflux from sperm: approaches and applications

Zahid NASEER, Ejaz AHMAD, Melih AKSOY*

Department of Reproduction and Artificial Insemination, Faculty of Veterinary Medicine, Adnan Menderes University, Aydın, Turkey

Received: 14.04.2014

Accepted: 16.05.2014

Published Online: 24.10.2014

Printed: 21.11.2014

Abstract: Following ejaculation, sperm appear morphologically mature, but functionally they are inactive. To attain functional capability or capacitation *in vivo*, sperm must reside in the female reproductive tract for a certain period of time before fertilization. In the female reproductive tract, cholesterol efflux from the sperm plasma membrane is considered vital for the initiation of capacitation. A number of *in vitro* studies have used biological and nonphysiological means to understand the phenomenon of cholesterol efflux and capacitation. In this review, we present a short overview of the role of cholesterol efflux in regulating sperm capacitation using various agents. This review provides insight to researchers designing new procedures for sperm preparation to be used in assisted reproductive technology.

Key words: Cholesterol efflux, sperm, capacitation, *in vitro* fertilization

1. Introduction

Sperm are very peculiar cells because they undergo several essential maturational changes throughout their life span to accomplish specific functions. The rates of these changes or modifications are highly variable at different subcellular regions (acrosome, post acrosome, midpiece, tail, etc.) of sperm. In the epididymis, epithelial secretions and the constituents of luminal fluids provide an environment for sperm to attain motility and fertilizing ability. Though freshly ejaculated sperm are motile and have fertilizing ability, they are unable to fuse with the oocyte immediately after ejaculation. During their migration through the female reproductive tract, sperm undergo a series of controlled biochemical and membranous changes known as capacitation (1). During capacitation, sperm experience a number of events such as an increase in respiration and motility patterns (2,3), removal of cholesterol from the plasma membrane (4), destabilization of membranes (5,6), an increase in intracellular pH and calcium (7,8), activation of second messenger systems (9), and increased tyrosine phosphorylation of sperm proteins (10). Subsequently, capacitated sperm undergo the acrosome reaction, which is a morphological alteration consisting of a series of point fusions between the outer acrosomal membrane and the overlying plasma membrane (11,12). These functional changes are not one event but are a combination of sequential and concomitant processes involving modifications at the molecular level in both the head (i.e. preparation for the acrosome reaction) and the tail (i.e. motility changes).

Existing research suggests that cholesterol depletion from the acrosomal region is a prerequisite for the acrosome reaction and sperm-egg binding. Similarly, changes in the cholesterol concentration in the plasma membrane over the midpiece and tail region increase lateral mobility of the membrane components, which in turn support the hyperactive motility of sperm (13). The presence of several cholesterol acceptors such as albumin and high-density lipoprotein (HDL) in the follicular and oviductal fluid mediates cholesterol efflux from the sperm membrane *in vivo* (14,15). However, *in vitro*, cholesterol efflux from the sperm plasma membranes has been facilitated by cholesterol acceptors such as bovine serum albumin (BSA) (6,16–18), HDL (14,19), and β -cyclodextrin (20). Likewise, seminal proteins and Percoll gradient centrifugation have been used to induce cholesterol efflux from sperm to promote the acrosome reaction (21–23). This review summarizes studies conducted on the approaches used for cholesterol efflux and their application in sperm capacitation, acrosome reaction, and fertilization ability.

2. The role of cholesterol in sperm membranes

The cellular membranes of sperm consist of phospholipids and sphingolipids in large proportions, along with sterols, which have a role in membrane composition for different regulatory functions (24). Cholesterol is the major essential sterol required for viability and cell proliferation. In addition, cholesterol stabilizes the membrane, reduces membrane permeability, facilitates morphological mem-

* Correspondence: aksoym@tnn.net

brane characteristics and enables cell-to-cell interactions, influences membrane phase transition, provides suitable microenvironments for membrane-associated proteins, and serves as a membrane antioxidant (25). The interaction of cholesterol with membrane phospholipids and sphingolipids influences membrane fluidity. The cholesterol/phospholipid ratio is an important determinant of membrane fluidity and stability. The distribution of cholesterol, which is incorporated during sperm maturation in the male genital tract, is variable at different regions of the sperm. Usually, the sperm head has a high amount of cholesterol, which regulates the membrane fluidity and plays an important role during the capacitation process (26).

3. Why is cholesterol efflux necessary for sperm capacitation?

Mammalian sperm are unable to fertilize an egg until they undergo capacitation, either during their passage through the female reproductive tract or in a suitable *in vitro* medium. During capacitation, spermatozoa become hyperactive and their acrosome becomes destabilized to initiate the acrosome reaction. The removal of seminal plasma depletes some sterols from the plasma membrane and capacitates the sperm. Cholesterol depletion from the acrosomal and postacrosomal regions is vital to capacitation and promotes the acrosome reaction and the sperm-egg binding process. The acrosomal region has a very tight membrane that is composed of sulfoconjugates of cholesterol (27). The presence of large charged and hydrated polar sulfate groups of these cholesterol sulfates provides stability and prevents disruption of the nonbilayer organization of the acrosomal membrane compared to other regions of sperm (28). Cholesterol efflux, after desulfation, is thought to increase membrane fluidity and allow greater lateral movements in integral membrane proteins. The increase in membrane fluidity enhances calcium influx in order to mediate the acrosome reaction (29). Similarly, changes in cholesterol concentrations in the plasma membrane over the midpiece and tail may increase lateral mobility of the membrane components, supporting hyperactivated motility for penetrating an egg.

4. Agents used for cholesterol efflux

Cholesterol depletion is a time-dependent process and can be accomplished *in vitro* by using biological and nonphysiological cholesterol acceptors (30). A wide variety of such cholesterol acceptors are used to modify the membrane cholesterol content in sperm and somatic cells. Following are some physiological and nonphysiological cholesterol acceptors that have been tested *in vitro*.

4.1. Bovine serum albumin

Serum albumin is a protein derived from human and bovine serum and is widely used as a carrier molecule for

a variety of substances (e.g., cholesterol, hormones, ions, fatty acids, amino acids). It also serves as a pH buffer, an osmotic pressure regulator, and a heavy metal ion chelator in different biological processes (31,32). In 1971 Toyoda et al. (33) reported successful *in vitro* fertilization in mice by using a chemically defined medium supplemented with BSA. Later, the role of BSA in capacitation media was tested for mouse, rat, and human sperm capacitation (7,16–19,22,34). Likewise, BSA has been used in most media to maintain sperm motility and to promote membrane cholesterol efflux (18–19,34). The mechanism of cholesterol removal by albumin is not clear, but BSA presumably solubilizes or hydrolyzes a small amount of cholesterol from the cell surface because it does not have specific binding sites for cholesterol (2). Removing cholesterol from the sperm plasma membrane markedly increased concentrations of the phospholipids when sperm were incubated in medium supplemented with BSA. The increase in phospholipids leads to increased membrane fluidity, which is the first landmark for the initiation of capacitation (35).

The role of BSA as a cholesterol acceptor has been compared with other compounds such as beta-cyclodextrin and HDL in *in vitro* studies. The detailed results of those studies regarding the influence of BSA on sperm capacitation and fertilization are presented in the Table. A considerable decrease (20%–60%) in cholesterol concentration occurred when rat or mouse epididymal sperm were incubated with different concentrations (4–20 mg/mL) of BSA (16,18). Similar observations were reported for human sperm incubated in the presence of BSA for cholesterol efflux and induction of the acrosome reaction (29,36). However, the effect of BSA on cholesterol efflux, the acrosome reaction, and the fertilization rate has not yet been investigated in many species, such as cattle, sheep, goats, and rabbits. Collectively these data offer persuasive evidence for a possible cholesterol acceptor role for BSA in other mammalian sperm capacitation. This is an area ripe for investigation.

4.2. β -Cyclodextrin

Cyclodextrins are nonbiological molecules found in the female reproductive tract or in oocyte envelopes, but these have been used as highly efficient cholesterol acceptors for cholesterol release as an early event of *in vitro* sperm capacitation and acrosome reaction. Cyclodextrins are cyclic oligosaccharides consisting of α -(1–4)-linked D-glycopyranose units, which are primary degradation products of starch. These compounds act as potent carriers for hydrophobic drugs because of their high water solubility. Cyclodextrins are usually found as hexamers (α), heptamers (β), or octomers (γ). The β -cyclodextrins as compared to α - and γ -cyclodextrins have the highest affinity for inclusion of cholesterol due to their differences in size and hydrophobicity (37,38). Earlier β -cyclodextrins were employed

Table. Effect of different cholesterol acceptors on sperm cholesterol efflux, capacitation, and fertilization rates in different species.

Author	Species	Conc. of agent used	Decrease in cholesterol (%)	Acrosome reaction (%)	Fertilization rate (%)
Bovine serum albumin					
Davis et al., 1979 (16)	Rat	4 mg/mL	~12% decrease	-	-
Go and Wolf, 1985 (18)	Mouse	20 mg/mL	~60% decrease	-	-
Osheroff et al., 1999 (36)	Human	5 mg/mL	~30% decrease	48%	-
Zarintash and Cross, 1996 (63)	Human	26 mg/mL	~26% decrease	-	-
β-Cyclodextrin					
Choi and Toyoda, 1998 (41)	Mouse	0.25 mM	----	45%	53%
		0.75 mM	~50% decrease	-	21%
Visconti et al., 1999 (30)	Mouse	M-β-CD (3 mM)	Sperm = 3 ± 0.6♦ Media = 410 ± 8♦	-	-
		2-OH-p-β-CD (3 mM)	Sperm = 120 ± 3♦ Media = 309 ± 5♦	75%	-
		Control	Sperm = 431 ± 4♦ Media = 11 ± 1♦	-	-
Iborra et al., 2000 (42)	Goat	8 mM	~50%–60% decrease	35%	-
Funahashi, 2002 (64)	Boar	MbCD (0.125–2 mM)	~20%–45% dose-dependent	-	-
		Control	Less than 20%	-	-
Osheroff et al., 1999 (36)	Human	M-β-CD (3 mM)	Sperm = 157 ± 2♦ Media = 295.7 ± 5♦	-	-
		2-OH-p-β-CD (10 mM)	Sperm = 157 ± 2♦ Media = 295 ± 5♦	-	-
		Control	Sperm = 459 ± 4.6♦ Media = 11.3 ± 0.3♦	-	-
Cross, 1999 (43)	Human	2.5–10 mM	89%	26%	-
Bovine seminal plasma					
Thérien et al., 1998 (15)	Mouse	120 µg/mL	~51%	38%	-
		HDL (100 µg/mL)	~39%	15	-
Percoll gradient centrifugation					
Tanphaichitr et al., 1996 (22)	Mouse		Control = 0.235 ± 0.04* PGC = 0.203 ± 0.024*	-	-
Furimsky et al., 2005 (23)	Mouse		Control = 0.338 ± 0.068* PGC = 0.120 ± 0.017*	Control ~90%–92% PGC ~95%–100%	Control = 57 ± 1 PGC = 74 ± 6
Miscellaneous					
Huneau et al., 1994 (60)	Ram	20% estrous sheep serum	Treatment ~60% Control ~14%	-	Treatment ~85% Control = 0.7%
Langlais et al., 1988 (19)	Human	5% human female serum or follicular fluid	Treatment ~50% Control ~14%	-	-
Sugraroek et al., 1991 (61)	Human	3.5% HSA medium	5%	-	-
Benoff et al., 1993 (6)	Human	30 mg/mL HSA	15%	-	-
Ehrenwald et al., 1988 (62)	Human	Incubated protein-free medium	31%	AR was induced by LPC-	-

*: nmol/mg sperm DNA; ♦: ng steroid/10⁶ sperm ± SEM.

to extract cholesterol from erythrocyte and model membranes (37–39). Similarly, derivatives of β -cyclodextrin (methyl- β -cyclodextrins and 2-hydroxyl- β -cyclodextrins) have been used efficiently for cholesterol efflux in cultured cells (40).

Numerous experiments have been conducted to evaluate the effect of cyclodextrins on cholesterol removal and sperm functions in different species (Table). Choi and Toyoda (41) found that cyclodextrins induced capacitation by removing cholesterol from mouse sperm *in vitro*. Visconti et al. (30) and Iborra et al. (42) concluded that β -cyclodextrin derivatives are equally effective cholesterol-binding heptasaccharides, which mediate cholesterol efflux from mouse and goat sperm in a concentration-dependent manner. Cross (43) observed that high cholesterol efflux achieved by using β -cyclodextrins or their derivatives is essential for the acrosome reaction in human sperm. These previous reports indicated that cyclodextrin possesses the ability to bind sperm membrane for cholesterol efflux; it could be a better choice than the other biological hydrophobic compounds for *in vitro* sperm preparation.

Inducing cholesterol in sperm plasma membrane with cyclodextrin molecules (CLCs) (44) enhanced the cryotolerance in rams (45), bulls (46–48), bucks (49), elephants (50), and stallions (51). This might be due to the fact that increased cholesterol in the sperm cell membrane enhanced the osmotic tolerance and inhibited the acrosome reaction (52). On the other hand, fertility after use of CLC-treated frozen-thawed semen is questionable. Although removal of cholesterol from the sperm membrane is essential for capacitation, further studies are needed regarding the effect of cholesterol removal from CLC-treated sperm after cryopreservation and its impact on fertilization or fertility.

4.3. Bovine seminal plasma

Bovine seminal plasma (BSP) contains a family of closely related major proteins designated as BSP-A1, BSP-A2, BSPA3, and BSP-30-kDa. These BSP proteins are secretory products of seminal vesicles; they bind to sperm upon ejaculation and play a role in capacitation (53). Upon ejaculation, the BSP proteins interact with the choline phospholipids on the sperm membrane (54) and bind to heparin in the uterine environment for capacitation (55). Therien et al. (15,21) described how BSP proteins accelerate capacitation of bovine epididymal sperm in the presence of heparin and HDL. Thérien et al. (15) also showed that the BSP proteins and HDL play an important role in the efflux of cholesterol from the plasma membrane of sperm during capacitation. Therefore, sperm capacitation by BSP could be effective in *in vitro* fertilization.

4.4. Percoll gradient centrifugation

Centrifugation through a gradient of Percoll has been used for more than 2 decades to prepare motile sperm from for

intrauterine insemination and *in vitro* fertilization (56,57). It is well known that Percoll gradient centrifugation (PGC)-capacitated mouse sperm has a higher fertilizing ability than washed capacitated sperm or whole populations of sperm (58,59). Tanphaichitr et al. (22) described how washing or PGC had no effect on the cholesterol and phospholipids content of mouse spermatozoa. In contrast, Furimsky et al. (23) reported that PGC lowered the cholesterol content in capacitated mouse sperm, which showed good fertilizing ability compared to the control. However, to the authors' knowledge, the effect of PGC on cholesterol removal and capacitation in other mammalian species has not been studied.

4.5. Miscellaneous agents

Several authors have reported significant cholesterol removal through the addition of various biological components to the incubation medium or through the use of biological fluids. Female biological fluids have also been used for cholesterol efflux from sperm. Huneau et al. (60) used 20% estrous sheep serum in an incubation medium for cholesterol efflux and reported higher cholesterol efflux (60% vs. 14%) and fertility (85% vs. 0.7%) in the treated ram sperm compared to the control. In another study, Langlais et al. (19) incubated human spermatozoa in a medium supplemented with 5% human female serum or follicular fluid. Their results showed increased cholesterol efflux (50%) in the presence of 5% human female serum or follicular fluid, whereas low cholesterol efflux (14%) was observed in spermatozoa incubated with protein-free medium. In contrast, Davis et al. (17) found that cholesterol efflux from sperm membranes was not statistically significant using comparable concentrations of serum albumin. Sugkraroek et al. (61) found that using 3.5% human serum albumin (HSA) medium only caused a 5% cholesterol removal from fertile donors' sperm. Similarly, Benoff et al. (6) reported 15% cholesterol efflux in human sperm when incubated with 30 mg/mL HSA in a medium. Ehrenwald et al. (62) did not detect any acrosome-reacted spermatozoa after 31% cholesterol efflux in 90 min. They obtained acrosome responsiveness only after the addition of lysophosphatidylcholine. These authors concluded that biological agents (estrous sheep serum, HAS, follicular fluid, etc.) might be useful for cholesterol efflux and capacitation in sperm.

5. Conclusion

In summary, the outcome of these studies (Table) provides evidence that BSA and β -cyclodextrins are the best choices for cholesterol efflux and sperm capacitation for *in vitro* fertilization. Other biological substances had little influence on sperm cholesterol efflux, but their effect on cholesterol efflux, capacitation, and fertility should be explored across more species.

Acknowledgment

Thanks to the Scientific and Technological Research Council of Turkey (TÜBİTAK) for providing a fellowship

to Zahid Naseer and Ejaz Ahmad within the framework of the BİDEB 2215 PhD Fellowship Programme for Foreign Citizens.

References

- de Lamirande E, Leclerc P, Gagnon C. Capacitation as a regulatory event that primes spermatozoa for the acrosome reaction and fertilization. *Mol Hum Reprod* 1997; 3: 175–194.
- Yanagimachi R. Requirements of extracellular calcium ions for various stages of fertilization and fertilization-related phenomena in the hamster. *Gamete Res* 1982; 5: 323–344.
- Zhu JJ, Barratt CLR, Cooke ID. Effect of human cervical mucus on human sperm motion and hyperactivation in vitro. *Hum Reprod* 1992; 7: 1402–1406.
- Davis BK. Uterine fluid proteins bind sperm cholesterol during capacitation in the rabbit. *Experientia* 1982; 38: 1063–1064.
- Wolf DE, Hagopian SS, Ishijima S. Changes in sperm plasma membrane lipid diffusibility after hyperactivation during in vitro capacitation in the mouse. *J Cell Biol* 1986; 102: 1372–1377.
- Benoff S, Hurley I, Cooper GW, Mandel FS, Rosenfeld DL, Herschlag A. Head-specific mannose-ligand receptor expression in human spermatozoa is dependent on capacitation associated membrane cholesterol loss. *Hum Reprod* 1993; 8: 2141–2154.
- Fraser LR. Mechanisms regulating capacitation and the acrosome reaction. In: Fenichel P, Parinaud J, editors. *Human Sperm Acrosome Reaction*. Colloque INSERM no. 236. Paris, France: John Libbey Eurotext Ltd.; 1995. pp. 17–33.
- Baldi E, Casana R, Falsetti C, Krausz C, Maggi M, Forti G. Intracellular calcium accumulation and responsiveness to progesterone in capacitating human spermatozoa. *J Androl* 1991; 12: 323–330.
- Bielfeld PR, Anderson A, Mack SR, De Jonge CJ, Zaneveld LJD. Are capacitation or calcium ion flux required for the human sperm acrosome reaction? *Fertil Steril* 1994; 62: 1255–1261.
- Leyton L, Saling P. 95 kd sperm proteins bind ZP3 and serve as tyrosine kinase substrates in response to zona binding. *Cell* 1989; 57: 1123–1130.
- Davis BK. Interaction of lipids with the plasma membrane of sperm cells. I. The antifertilization action of cholesterol. *Arch Androl* 1980; 5: 249–254.
- Yanagimachi R. Mammalian fertilization. In: Knobil E, Neill JD, editors. *The Physiology of Reproduction*. New York, NY, USA: Raven Press; 1994. pp. 189–317.
- Lin Y, Mahan K, Lathrop WF, Myles DG, Primakoff P. A hyaluronidase activity of the sperm plasma membrane protein PH-20 enables sperm to penetrate the cumulus cell layer surrounding the egg. *J Cell Biol* 1994; 125: 1157–1163.
- Ehrenwald E, Foote RH, Parks JE. Bovine oviductal fluid components and their potential role in sperm cholesterol efflux. *Mol Reprod Develop* 1990; 25: 195–204.
- Thérien I, Moreau R, Manjunath P. Major proteins of bovine seminal plasma and high-density lipoprotein induce cholesterol efflux from epididymal sperm. *Biol Reprod* 1998; 59: 768–776.
- Davis BK, Byrne R, Bedigian K. Studies on the mechanism of capacitation: albumin-mediated changes in plasma membrane lipids during in vitro incubation of rat sperm. *Proc Natl Acad Sci U S A* 1979; 668: 257–266.
- Davis BK, Byrne R, Bedigan K. Studies on the mechanism of capacitation: albumin mediated changes in plasma membrane lipids during in vitro incubation of rat sperm cells. *Proc Natl Acad Sci U S A* 1980; 77: 1546–1550.
- Go KJ, Wolf DP. Albumin-mediated changes in sperm sterol content during capacitation. *Biol Reprod* 1985; 32: 145–153.
- Langlais J, Kan FWK, Granger L, Raymond L, Bleau G, Roberts KD. Identification of sterol acceptors that stimulate cholesterol efflux from human spermatozoa during in vitro capacitation. *Gamete Res* 1988; 20: 185–201.
- Yancey PG, Rodriguez WV, Kilsdonk EP, Stoudt GW, Johnson WJ, Phillips MC, Rothblat GH. Cellular cholesterol efflux mediated by cyclodextrins. Demonstration of kinetic pools and mechanism of efflux. *J Biol Chem* 1996; 271: 16026–16034.
- Thérien I, Soubeyrand S, Manjunath P. Major proteins of bovine seminal plasma modulate sperm capacitation by high-density lipoprotein. *Biol Reprod* 1997; 57: 1080–1088.
- Tanphaichitr N, Zheng Y, Kates M, Abdullah N, Chan A. Cholesterol and phospholipid levels of washed and Percoll-gradient centrifuged mouse sperm: presence of lipids possessing inhibitory effects on sperm motility. *Mol Reprod Develop* 1996; 43: 187–195.
- Furimsky A, Vuong N, Xu H, Kumarathanan P, Xu M, Weerachatanukul W, Khalil MB, Kates M, Tanphaichitr N. Percoll gradient-centrifuged capacitated mouse sperm have increased fertilizing ability and higher contents of sulfogalactosylglycerolipid and docosahexaenoic acid-containing phosphatidylcholine compared to washed capacitated mouse sperm. *Biol Reprod* 2005; 72: 574–583.
- Cincik M, Ergur AR, Tutuncu L, Muhcu M, Kilic M, Balaban B, Urman B. Combination of hypoosmotic swelling/eosin Y test for sperm membrane integrity evaluation: correlations with other sperm parameters to predict ICSI cycles. *Arch Androl* 2007; 53: 25–28.
- Crockett EL. Cholesterol function in plasma membranes from ectotherms: membrane-specific roles in adaptation to temperature. *Amer Zool* 1998; 38: 291–304.
- Martinez P, Morros A. Membrane lipid dynamics during human sperm capacitation. *Front Biosci* 1996; 1: 103–117.

27. Langlais J, Zollinger M, Plante L, Chapdelaine A, Bleau G, Roberts KD. Localization of cholesterol sulfate in human spermatozoa in support of a hypothesis for the mechanism of capacitation. *Proc Natl Acad Sci U S A* 1981; 78: 7266–7270.
28. Cheetham JJ, Chen RJB, Epand RM. Interaction of calcium and cholesterol sulphate induces membrane destabilization and fusion: implications for the acrosome reaction. *Biochim Biophys Acta* 1990; 1024: 367–372.
29. Fayrer-Hosken RA, Brackett BG, Brown J. Reversible inhibition of rabbit fertilizing ability by cholesterol sulfate. *Biol Reprod* 1987; 36: 878–883.
30. Visconti PE, Stewart-Savage J, Blasco A, Battaglia L, Miranda P, Kopf GS, Tezón JG. Roles of bicarbonate, cAMP, and protein tyrosine phosphorylation on capacitation and the spontaneous acrosome reaction of hamster sperm. *Biol Reprod* 1999; 61: 76–84.
31. Bavister BD. Substitution of a synthetic polymer for protein in a mammalian gamete culture system. *J Exp Zool* 1981; 217: 45–51.
32. Maurer HR. Towards serum-free, chemically defined media for mammalian cell culture. In: Freshney RI, editor. *Animal Cell Culture: A Practical Approach*. 2nd ed. Oxford, UK: Oxford University Press; 1992. pp. 15–46.
33. Toyoda YM, Yokoyama M, Hoshi T. Studies on the fertilization of mouse eggs in vitro. II. Effects of in vitro preincubation on time of sperm penetration of mouse eggs in vitro. *Japanese J Anim Reprod* 1971; 16: 152–157.
34. Byrd W, Tsu J, Wolf D. Kinetics of spontaneous and induced acrosomal loss in human sperm incubated under capacitating and non-capacitating conditions. *Gamete Res* 1989; 22: 109–122.
35. Cross NL. Decrease in order of human sperm lipids during capacitation. *Biol Reprod* 2003; 69: 529–534.
36. Osheroff JE, Visconti PE, Valenzuela JP, Travis AJ, Alvarez J, Kopf GS. Regulation of human sperm capacitation by a cholesterol efflux-stimulated signal transduction pathway leading to protein kinase A-mediated up-regulation of protein tyrosine phosphorylation. *Mol Hum Reprod* 1999; 5: 1017–1026.
37. Ohtani Y, Irie T, Uekama K, Fukunaga K, Pitha J. Differential effects of alpha, beta and gamma-cyclodextrins on human erythrocytes. *Eur J Biochem* 1989; 186: 17–22.
38. Ohvo H, Slotte JP. Cyclodextrin-mediated removal of sterols from monolayers: effects of sterol structure and phospholipids on desorption rate. *Biochemistry* 1996; 35: 8018–8024.
39. Irie T, Fukunaga K, Pitha J. Hydroxypropylcyclodextrins in parenteral use. I: Lipid dissolution and effects on lipid transfers in vitro. *J Pharma Sci* 1992; 81: 521–523.
40. Kilsdonk EPC, Yancey PG, Stoudt GW, Bangerter FW, Johnson WJ, Phillips MC, Rothblat GH. Cellular cholesterol efflux mediated by cyclodextrins. *J Biol Chem* 1995; 270: 17250–17256.
41. Choi Y, Toyoda Y. Cyclodextrin removes cholesterol from mouse sperm and induces capacitation in a protein-free medium. *Biol Reprod* 1998; 59: 1328–1333.
42. Iborra A, Companyo M, Martinez P, Morros A. Cholesterol efflux promotes acrosome reaction in goat spermatozoa. *Biol Reprod* 2000; 62: 378–383.
43. Cross NL. Effect of methyl- β -cyclodextrin on the acrosomal responsiveness of human sperm. *Mol Reprod Develop* 1999; 53: 92–98.
44. Moore AI, Squires EL, Graham JK. Adding cholesterol to the stallion sperm plasma membrane improves cryosurvival. *Cryobiology* 2005; 51: 241–249.
45. Mocé E, Purdy PH, Graham JK. Treating ram sperm with cholesterol-loaded cyclodextrins improves cryosurvival. *Anim Reprod Sci* 2010; 118: 236–247.
46. Purdy PH, Graham JK. Effect of cholesterol-loaded cyclodextrin on the cryosurvival of bull sperm. *Cryobiology* 2004; 48: 36–45.
47. Amorim EAM, Graham JK, Spizziri B, Meyers M, Torres CAA. Effect of cholesterol or cholesteryl conjugates on the cryosurvival of bull sperm. *Cryobiology* 2009; 58: 210–214.
48. Moraes EA, Graham JK, Torres CAA, Meyers M, Spizziri B. Delivering cholesterol or cholesterol to bull sperm membranes improves cryosurvival. *Anim Reprod Sci* 2010; 118: 148–154.
49. Amidi F, Farshad A, Khor AK. Effects of cholesterol-loaded cyclodextrin during freezing step of cryopreservation with TCGY extender containing bovine serum albumin on quality of goat spermatozoa. *Cryobiology* 2010; 61: 94–99.
50. Kiso WK, Asano A, Travis AJ, Schmitt DL, Brown JL, Pukazhenthil BS. Pretreatment of Asian elephant (*Elephas maximus*) spermatozoa with cholesterol-loaded cyclodextrins and glycerol addition at 48C improves cryosurvival. *Reprod Fertil Develop* 2012; 24: 1134–1142.
51. Spizziri BE, Fox MH, Bruemmer JE, Squires EL, Graham JK. Cholesterol-loaded-cyclodextrins and fertility potential of stallion's spermatozoa. *Anim Reprod Sci* 2010; 118: 255–264.
52. Aksoy M, Akman O, Lehimcioglu NC, Erdem H. Cholesterol-loaded cyclodextrin enhances osmotic tolerance and inhibits the acrosome reaction in rabbit spermatozoa. *Anim Reprod Sci* 2010; 120: 166–172.
53. Manjunath P, Sairam MR. Purification and biochemical characterization of three major acidic proteins (BSP-A1, BSP-A2 and BSP-A3) from bovine seminal plasma. *Biochem J* 1987; 241: 685–692.
54. Desnoyers L, Manjunath P. Major proteins of bovine seminal plasma exhibit novel interactions with phospholipid. *J Biol Chem* 1992; 267: 10149–10155.
55. Parrish JJ, Susko-Parrish JL, Winer MA, First NL. Capacitation of bovine sperm by heparin. *Biol Reprod* 1988; 38: 1171–1180.
56. Kelly MP, Corson SL, Gocial B, Batzer FR, Gutmann JN. Discontinuous Percoll gradient preparation for donor insemination: determinants for success. *Hum Reprod* 1997; 12: 2682–2686.

57. Hammadeh ME, Zavos PM, Rosenbaum P, Schmidt W. Comparison between the quality and function of sperm after semen processing with two different methods. *Asian J Androl* 2001; 3: 125–130.
58. Gravance CG, Champion ZJ, Sax-Gravance SK, Casey PJ. Percentage of normal sperm heads is significantly increased by Percoll separation of semen. *Int J Androl* 1998; 21: 116–119.
59. Prakash P, Leykin L, Chen Z, Toth T, Sayegh R, Schiff I, Isaacson K. Preparation by differential gradient centrifugation is better than swim-up in selecting sperm with normal morphology (strict criteria). *Fertil Steril* 1998; 69: 722–726.
60. Huneau D, Crozet N, Ahmed-Ali M. Estrous sheep serum as a potent agent for ovine IVF: effect on cholesterol efflux from spermatozoa and the acrosome reaction. *Theriogenology* 1994; 42: 1017–1028.
61. Sugkraroek P, Kates M, Leader A, Tanphaichitr N. Levels of cholesterol and phospholipids in freshly ejaculated sperm and Percoll gradient pelleted sperm from fertile and unexplained infertile men. *Fertil Steril* 1991; 55: 820–827.
62. Ehrenwald E, Parks JE, Foote RH. Cholesterol efflux from bovine sperm. I. Induction of the acrosome reaction with lysophosphatidylcholine after reducing sperm cholesterol. *Gamete Res* 1988; 20: 145–157.
63. Zarintash RJ, Cross NL. Unesterified cholesterol content of human sperm regulates the response of the acrosome to the agonist, progesterone. *Biol Reprod* 1996; 55: 19–24.
64. Funahashi H. Effect of methyl- β -cyclodextrin and fertilization promoting peptide on capacitation of boar spermatozoa in a protein-free medium. *J Reprod Develop* 2002; 48: 57–63.