

Comparison of pre- and postpartum serum leptin, ghrelin, and lipid levels in sheep

Ethem Mutlu TEMİZEL^{1*}, Hüseyin CİHAN¹, Pınar LEVENT¹, Ahmet SARIL¹, Yeşim ÖZARDA², Zeki YILMAZ¹

¹Department of Internal Medicine, Veterinary School Medicine, Uludağ University, Bursa, Turkey

²Department of Biochemistry, Faculty of Medicine, Uludağ University, Bursa, Turkey

Received: 30.05.2017 • Accepted/Published Online: 29.03.2018 • Final Version: 08.06.2018

Abstract: Serum leptin and ghrelin play important roles in energy metabolism and inducing appetite during pregnancy and lactation. We investigated the serum levels of leptin and ghrelin and their relationship with the duration of lactation and serum biochemical responses during the prepartum (last week before lambing) and postpartum periods in sheep. They were also evaluated between pregnant and nonpregnant (control ewes) ewes. For these purposes, pregnant (n = 15) and control Awassi ewes (n = 15) that were 2 years of age were used in this study. Blood samples were collected 1 week before the expected date of lambing (baseline) and for 12 weeks after parturitions. Serum leptin, ghrelin, cholesterol, triglyceride, VLDL, LDL, HDL, NEFA, and phospholipid levels were assayed. Leptin and ghrelin levels of the prepartum period in pregnant and control ewes were 4.5 ng/mL and 3.4 ng/mL (P < 0.05) and 75 pg/mL and 166 pg/mL (P < 0.01), respectively. The baseline value of ghrelin (81 pg/mL) decreased to 43 pg/mL within 1 week after parturition (P < 0.01). Serum ghrelin levels increased after 1 week. There was also a significant difference in serum ghrelin levels between control ewes and those a week before parturition in pregnant ewes (P < 0.001). The NEFA level was higher in pregnant than in control sheep (P < 0.001). In conclusion, leptin levels tended to decrease during the lactation period, which could be associated with a negative energy balance. Additionally, ghrelin levels tended to gradually increase during lactation to adapt to a negative energy balance. We think that variations in serum leptin and ghrelin might be used to monitor metabolic adaptation during lactation.

Key words: Leptin, ghrelin, parturition, energy balance, sheep

1. Introduction

Awassi sheep are primarily found in southeastern Turkey as well as the Middle East. Awassi sheep are a breed of dairy sheep bred for multiple lambing and high milk yield in Turkey (1). Pregnancy is the most important metabolic burden in ewes, which have multiple offspring (2,3). In this period, energy and protein intake usually do not cover the energy and protein requirements in many sheep herds (3,4).

Leptin, a product of the *ob* gene, is a protein mainly secreted by white adipose tissue (5). This hormone is responsible for the long-term regulation of energy balance and depressing appetite (6). Leptin plays an important role in energy metabolism during pregnancy and lactation (7,8). Many studies showed that plasma leptin level increased during pregnancy and declined during the transition to lactation (9–12). Carcangiu et al. (13) reported that body condition scores were positively correlated with serum leptin concentration but were negatively correlated with milk yield after delivery in ewes with production levels. Additionally, leptin was negatively correlated with growth hormone (GH) and nonesterified fatty acid (NEFA) in the lactation periods (13).

Otherwise, ghrelin mainly arises from the abomasum in ruminants. Ghrelin induces GH release and stimulates appetite in the brain (14–16). Ghrelin concentration increases prior to meals, but it is suppressed after feeding. Ghrelin also decreases fat oxidation and increases insulin concentration (17–19).

The objective of the present study was to evaluate fluctuations and possible interactions of the leptin, ghrelin, and lipid profile of ewes in the pre- and postparturient periods and control sheep.

2. Materials and methods

2.1. Animals and nutrition

The present study was carried out in the town of Nilüfer in Bursa Province, Turkey. Pregnant Awassi ewes (last week before lambing) that were (n = 15) or were not pregnant (control group) (n = 15), weighing 68.4 ± 10.4 kg (53.4–80.6 kg), and aged 2 years were used in the study. All ewes were selected from among those that were to have twin lambs. During this study period, all sheep were fed a controlled diet and water was provided ad libitum. All animals were housed under the same conditions. The herd consisted of a

* Correspondence: ethem@uludag.edu.tr

total of 150 Awassi sheep. In different compartments of the same pen, all animals were fed alfalfa and foraged wheat bran, bran, sunflower meal, and soybean meal, as well as a concentrated commercial mixture for lambs, which consisted of ground corn, barley, calcium carbonate, and vitamin premix. Blood sampling was performed under the control of the farm health coordinator in accordance with animal welfare guidelines. The study was approved by the Committee of the Animal Experiments of Uludağ University (1.06.2004/2).

2.2. Collection of samples

The study was carried out between February and May. Blood samples were collected from the jugular vein between 1130 and 1430 a week before the expected date of lambing parturition, at baseline, and at 0.5 (day 4 just after the parturition), 1, 2, 3, 4, 6, 8, and 12 weeks after parturitions. Serum was then extracted following centrifugation, frozen, and stored at -80°C until analyzed.

2.3. Biochemical analysis

Serum leptin and ghrelin levels were measured by radioimmunoassay (RIA) using a commercially available kit (Multispecies leptin RIA kit and active ghrelin RIA kit, Linco Research, St. Charles, MO, USA). Since active ghrelin is too unstable to be measured in stored samples (20) and the acidification of plasma prevents the rapid deacylation of ghrelin (21), 1 N hydrogen chloride was added to serum samples before freezing (22).

Serum total cholesterol, high-density lipoprotein cholesterol (HDL-C), very low-density lipoprotein cholesterol (VLDL-C), and triglyceride levels were measured by an automated clinical chemistry analyzer (Architect ci8200; Abbott GmbH Co KG, Wiesbaden, Germany) using commercially available assay kits (Abbott GmbH Co KG). Low-density lipoprotein cholesterol (LDL-C) concentrations were calculated using the formula:

$\text{LDL-C (in milligrams per deciliter)} = \text{total cholesterol} - (\text{HDL} - \text{C} + \text{triglyceride}/5)$. Serum nonesterified fatty acid (NEFA) concentration was measured using a commercially available enzymatic colorimetric assay kit (Wako Chemicals, Neuss, Germany). Serum phospholipids were measured by an enzymatic colorimetric method using a commercially available assay kit (Wako Chemicals) and were reported as choline-containing phospholipids (in milligrams per deciliter).

2.4. Statistical analysis

Data sets were analyzed by analysis of variance (ANOVA), and Tukey's test was chosen as the post hoc multiple comparison test. Bivariate correlation was applied and Pearson's correlation was chosen. Differences were considered significant at a probability level of $P < 0.05$, $P < 0.01$, and $P < 0.001$ in all analyses. All statistical analyses were performed with SigmaStat (SigmaStat GmbH, Erkrath, Germany).

3. Results

There were 7 singletons and 8 twins in the pregnant ewes. Leptin levels tended to decrease during lactation (Figure 1). Baseline values of ghrelin (81 pg/mL) decreased to 43 pg/mL within 1 week of parturition ($P < 0.01$). Serum ghrelin levels tended to increase after a 1-week period (Figure 1). Parturition leptin and ghrelin levels in pregnant and control ewes were 4.5 ng/mL and 3.4 ng/mL ($P < 0.05$) and 75 pg/mL and 166 pg/mL ($P < 0.001$), respectively (Figure 2; Table).

Repeated measurements of cholesterol levels surged during lactation. Cholesterol levels tended to decrease up to week 3 compared to the baseline; thereafter, they returned to a baseline value at week 12. Triglyceride levels were low up to week 12 compared to the baseline level. LDL, HDL, and VLDL levels had a surge during the




Figure 1. Serum leptin and ghrelin levels profile in baseline (birth) and postparturient period.




Figure 2. Serum leptin and ghrelin levels in control (nonpregnant ewes) and preparturient ewes.

Table. Serum leptin, ghrelin and lipid profile in preparturient, postparturient, and control ewes.

	Leptin (nmol/L)	Ghrelin (pmol/L)	Total cholesterol (mg/dL)	Triglyceride (mg/dL)	LDL (mg/dL)	HDL (mg/dL)	VLDL (mg/dL)	Phospholipid (mg/dL)	NEFA (mmol/L)
Control	3.48 ± 0.30 ^A	166.00 ± 7.00 ^a	75.00 ± 7.00	26.00 ± 0.80 ^c	31.00 ± 0.70	38.00 ± 1.40 ^a	5.00 ± 0.20 ^a	112.00 ± 3.00 ^a	0.32 ± 0.03 ^a
Prepartum	4.58 ± 0.20 ^B	75.00 ± 7.00 ^b	70.00 ± 7.00 ^A	40.00 ± 0.80 ^d	32.00 ± 0.70	31.00 ± 1.00 ^b	8.00 ± 0.20 ^b	105.00 ± 2.00 ^b	0.35 ± 0.03 ^b
Baseline	5.00 ± 0.50 ^a	81.00 ± 10.00 ^c	69.00 ± 0.70 ^A	40.00 ± 0.90 ^c	32.00 ± 0.70	31.00 ± 0.30 ^a	8.00 ± 0.20 ^a	105.00 ± 2.00 ^a	0.36 ± 0.02 ^a
0.5 week	2.60 ± 0.40 ^b	43.00 ± 8.00 ^d	57.00 ± 0.70 ^B	16.00 ± 0.50 ^d	32.00 ± 0.70	27.00 ± 0.30 ^b	3.20 ± 0.10 ^b	83.00 ± 4.00 ^b	0.63 ± 0.04 ^b
Week 1	3.00 ± 0.30 ^b	62.00 ± 4.00 ^d	50.00 ± 0.80 ^B	15.00 ± 0.60 ^d	36.00 ± 0.30 ^A	22.00 ± 0.30 ^b	3.00 ± 0.10 ^b	79.00 ± 3.00 ^b	0.38 ± 0.03
Week 2	3.00 ± 0.40 ^b	80.00 ± 9.00	55.00 ± 0.90 ^B	19.00 ± 0.50 ^d	25.00 ± 0.30 ^B	26.00 ± 0.30 ^b	3.30 ± 0.10 ^b	87.00 ± 4.00 ^b	0.36 ± 0.02 ^a
Week 3	2.50 ± 0.50 ^b	50.00 ± 5.00 ^d	51.00 ± 0.70 ^B	14.00 ± 0.40 ^d	27.00 ± 0.90 ^B	25.00 ± 0.70 ^b	3.00 ± 0.10 ^b	91.00 ± 4.00 ^b	0.42 ± 0.04 ^a
Week 4	2.50 ± 0.30 ^b	53.00 ± 6.00 ^d	55.00 ± 0.60 ^B	17.00 ± 0.30 ^d	26.00 ± 0.70 ^B	27.00 ± 0.70 ^b	3.10 ± 0.10 ^b	96.00 ± 0.30 ^b	0.46 ± 0.03 ^a
Week 6	3.20 ± 0.30 ^b	76.00 ± 8.00	62.00 ± 0.80 ^B	20.00 ± 0.50 ^d	25.00 ± 0.30 ^B	32.00 ± 0.70	3.00 ± 0.10 ^b	105.00 ± 4.00	0.30 ± 0.03 ^a
Week 8	3.10 ± 0.30 ^b	97.00 ± 9.00	65.00 ± 0.70 ^B	24.00 ± 0.40 ^d	26.00 ± 0.30 ^B	34.00 ± 0.30 ^a	4.00 ± 0.20 ^b	101.00 ± 3.00	0.33 ± 0.02 ^a
Week 12	2.50 ± 0.40 ^b	120.00 ± 10.00 ^d	68.00 ± 0.70	13.00 ± 0.30 ^d	28.00 ± 0.30 ^B	36.00 ± 0.70 ^a	3.00 ± 0.10 ^b	104.00 ± 4.00	0.41 ± 0.03 ^a

^{ab} indicate statistically significance among column P < 0.001

^{AB} indicate statistically significance among column P < 0.05

^{c,d and cd} indicate statistically significance among column P < 0.01

postparturient period, but these parameters were lower than baseline levels, except at week 1 for LDL and weeks 6, 8, and 12 for HDL (Figure 3; Table)

Serum leptin, triglyceride, VLDL, and NEFA levels were higher in pregnant sheep (before parturient) than in control ewes (P < 0.001). Serum levels of ghrelin, cholesterol, HDL, and phospholipid were lower in pregnant ewes compared to the controls (P < 0.001) (Table).

4. Discussion

In the present study, leptin concentration was high in the prepartum and baseline periods, while it was at the lowest level in week 3, week 4, and week 12 of lactation (P < 0.001). The fluctuating decline in leptin concentration continued until week 12 compared to the baseline.

Observed decreases in serum leptin levels in this study might be related to changes in adipose tissue density in response to parturition in sheep. This low level of leptin may also be associated with the negative energy balance that resulted from increasing metabolic requirements just after parturition, consistent with increasing serum NEFA at the same time point (day 4 after parturition), as reported previously (13,23). Reduced dry matter intake in response to low leptin concentration may be thought to increase the negative energy balance in the sheep studied. Despite the fluctuating course of the NEFA level in the later periods after day 4 of the lactation period, the higher level compared to the initial value may be related to the lower concentration of leptin in this period. However, this may be associated with multifactorial effects, such as nutritional status,




Figure 3. Lipid profile in preparturient and postparturient periods.

breed differences, and other hormonal factors (2,24). The onset of lactation is characterized by a decrease in dry matter consumption versus increased energy demand due to milk production. This leads to the formation of a negative energy balance and a decrease in the body score (2). Leptin concentration is closely related to nutritional status, adipose tissue density, and other hormonal levels (5,25,26). Leptin concentration, which varies according to adipose tissue density, results in decreased dry matter

intake due to the effects on the central nervous system (26–28). Studies in cattle and sheep have indicated a decrease in the postpartum plasma leptin concentration (13,29). However, there is an increase in blood NEFA concentration and a decrease in body score, especially as a sign of negative energy balance in high milk-producing animals (30,31). Taken together, our observations of low serum leptin and an increase in metabolic requirements (meaning negative energy balance) agreed well with the

study by Carcangiu et al. (13) that reported that the high milk yield group had low plasma leptin concentration compared to the low milk yield group.

In the present study, the ghrelin level was low compared to the initial level, but a fluctuating course was seen. At week 2, ghrelin reached the baseline value, but it was accompanied by decreased serum leptin levels at weeks 3 and 4. In the present study, observed increases in the level of NEFA during this period suggest that the decline in both leptin and ghrelin levels may be considered a physiological hormonal response to the negative energy balance. It has been determined that the ghrelin level reached baseline levels at 6 weeks. The fact that the level of NEFA in these periods is close to the values in the baseline and control sheep can be considered a sign of appetite stimulation. Serum leptin and ghrelin significantly affect fetal development during pregnancy (2). As in other mammalian species, ghrelin also stimulates GH hormone from the pituitary gland in ruminants (16). Thus, blood ghrelin concentration may vary during the development and lactation periods. There is also an increase in plasma ghrelin with a decrease in the rumen fold and as a response to hunger (14,15). Kurose et al. (32) reported that the level of ghrelin measured after fasting was higher in animals with more adipose tissue than in weaker sheep.

Ghrelin is known to have important roles in the energy metabolism necessary for the stimulation of appetite, glucose release, cell proliferation, and reproductive activity. The level of maternal ghrelin has been shown to have a great effect on fetal development, especially in the late period of pregnancy (33,34). Fuglsang et al. (35) reported that the ghrelin level was higher in the early and middle periods of pregnancy compared to the baseline level, whereas it decreased significantly in the late gestation period. In accordance with these observations (33–35), in the present study, the ghrelin level was significantly higher in the control sheep than in the late gestation sheep. The difference in the ghrelin level in the sheep in the control and prepartum periods in the present study suggests that ghrelin may be associated with an increase in the ghrelin mRNA level versus the lower blood level of ghrelin in the late period of pregnancy (35).

Leptin plays an important role in regulating fat metabolism (36). Leptin has a role in the regulation of acute triglyceride metabolism, HDL-cholesterol clearance, lipid oxidation, and the lipogenic effects of insulin at the cellular level (37,38). Various studies have suggested that leptin is positively associated with cholesterol and HDL. Otherwise, a weak positive correlation with triglyceride concentration has been reported (36,38,39). In a study by Comba et al. (24), the relationship between leptin level and lipid profile was evaluated in different sheep breeds that were not pregnant and 2–3 years old. In that study

(24), there was a positive correlation between leptin concentration and VLDL, LDL, total cholesterol, and triglyceride concentrations. Cohen et al. (40) reported that triglyceride levels were elevated in animals that received exogenous leptin. In the present study, there were no statistically significant differences between postpartum leptin and triglycerides, VLDL, and total cholesterol levels, but a positive relationship was found between them. The level of triglycerides was low until week 12 after birth ($P < 0.01$), but the level of LDL was low from week 2 to week 12 based on the baseline value ($P < 0.05$). Total cholesterol level was low until week 8 compared to the baseline ($P < 0.05$). These findings suggest that low postpartum leptin levels may be associated with negative energy balance, suggesting that the low level of leptin could also induce changes in the lipid profile in the postparturient period, in agreement with a previous study (41).

Total cholesterol and phospholipids play an important role in the membrane and basic substance of the cell. For this reason, various studies reported that the levels of both have not changed (42,43). However, it was noted by Caldeira and Portugal (31) that total cholesterol and phospholipid levels were elevated in sheep with high body condition scores. Another study, by Nielsen and Jakobsent (44), noted that total cholesterol, triglyceride, and phospholipid levels were high in goats with high feed consumption. In the present study, phospholipid and total cholesterol levels decreased between day 4 and week 4, and the leptin level was also lower than the baseline levels in this period. The increasing level of NEFA in this period compared to the initial level suggests the presence of a negative energy balance. The present study findings suggest that low levels of leptin may be associated with a decrease in total cholesterol and phospholipid levels, particularly in the early stages of lactation, which agreed with the findings reported by Bitman and Wood (45). In the present study, the total cholesterol level was low in ewes in the prepartum period compared to control ewes, whereas various authors reported that total cholesterol levels were reduced throughout the late pregnancy period due to rising steriogenesis (46–48).

In conclusion, this study reveals the changes in serum leptin and ghrelin levels in the pre- and postnatal periods in sheep and the relationship between these changes and the serum lipid profile. Our findings show that serum leptin and ghrelin levels may be effective in clarifying the changing physiological conditions in the postpartum period. Our findings in this study provide important data for studies in which negative energy balance and hormonal relationships are assessed in ewes.

Acknowledgment

This study was supported in part by the Research Fund of Uludağ University (V-2003/86).

References

1. Yilmaz O, Kor A, Ertugrul M, Wilson R. The domestic livestock resources of Turkey: goat breeds and types and their conservation status. *Anim Gene Res* 2012; 51: 105-116.
2. Fthenakis G, Arsenos G, Brozos C, Fragkou I, Giadinis N, Giannenas I, Mavrogianni VS, Papadopoulos E, Valasi I. Health management of ewes during pregnancy. *Anim Reprod Sci* 2012; 130: 198-212.
3. Rook JS. Pregnancy toxemia of ewes, does, and beef cows. *Vet Clin North Am Food Anim Pract* 2000; 16: 293.
4. Brozos C, Mavrogianni V, Fthenakis G. Treatment and control of peri-parturient metabolic diseases: pregnancy toxemia, hypocalcemia, hypomagnesemia. *Vet Clin North Am Food Anim Pract* 2011; 27: 105-113.
5. Caro JF, Sinha MK, Kolaczynski JW, Zhang PL, Considine RV. Leptin: the tale of an obesity gene. *Diabetes* 1996; 45: 1455-1462.
6. Nowroozi-Asl A, Aarabi N, Rowshan-Ghasrodashti A. Ghrelin and its correlation with leptin, energy related metabolites and thyroidal hormones in dairy cows in transitional period. *Pol J Vet Sci* 2016; 19: 197-204.
7. Woodside B, Abizaid A, Walker C. Changes in leptin levels during lactation: implications for lactational hyperphagia and anovulation. *Horm Behav* 2000; 37: 353-365.
8. Brogan RS, Mitchell SE, Trayhurn P, Smith MS. Suppression of leptin during lactation: contribution of the suckling stimulus versus milk production. *Endocrinol* 1999; 140: 2621-2627.
9. Tomimatsu T, Yamaguchi M, Murakami T, Ogura K, Sakata M, Mitsuda N, Kanzaki T, Kurachi H, Irahara M, Miyake A et al. Increase of mouse leptin production by adipose tissue after midpregnancy: gestational profile of serum leptin concentration. *Biochem Biophys Res Commun* 1997; 240: 213-215.
10. Ehrhardt R, Slepatis R, Bell A, Boisclair Y. Maternal leptin is elevated during pregnancy in sheep. *Domestic Animal Endocrinol* 2001; 21: 85-96.
11. Block S, Butler W, Ehrhardt R, Bell A, Van Amburgh M, Boisclair Y. Decreased concentration of plasma leptin in periparturient dairy cows is caused by negative energy balance. *J Endocrinol* 2001; 171: 339-348.
12. Liefers S, Veerkamp R, te Pas M, Delavaud C, Chilliard Y, van der Lende T. Leptin concentrations in relation to energy balance, milk yield, intake, live weight, and estrus in dairy cows. *J Dairy Sci* 2003; 86: 799-807.
13. Carcangiu V, Giannetto C, Luridiana S, Fazio F, Mura M, Parmeggiani A, Piccione G. Lactation influences the serum level of leptin and growth hormone during the daily bathyphase in ewes. *Biol Rhythm Res* 2016; 1016: 1-7.
14. Itoh F, Komatsu T, Yonai M, Sugino T, Kojima M, Kangawa K, Hasegawa Y, Terashima Y, Hodate K. GH secretory responses to ghrelin and GHRH in growing and lactating dairy cattle. *Domestic Animal Endocrinol* 2005; 28: 34-45.
15. De Vriese C, Delporte C. Ghrelin: A new peptide regulating growth hormone release and food intake. *Int J Biochem Cell Biol* 2008; 40: 1420-1424.
16. Hayashida T, Murakami K, Mogi K, Nishihara M, Nakazato M, Mondal M, Horii Y, Kojima M, Kangawa K, Murakami N. Ghrelin in domestic animals: Distribution in stomach and its possible role. *Domestic Animal Endocrinol* 2001; 21: 17-24.
17. Sugino T, Yamaura J, Yamagishi M, Ogura A, Hayashi R, Kurose Y, Kojima M, Kangawa K, Hasegawa Y, Terashima Y. A transient surge of ghrelin secretion before feeding is modified by different feeding regimens in sheep. *Biochem Biophys Res Commun* 2002; 298: 785-788.
18. Takahashi H, Kurose Y, Kobayashi S, Sugino T, Kojima M, Kangawa K, Hasegawa Y, Terashima Y. Ghrelin enhances glucose-induced insulin secretion in scheduled meal-fed sheep. *J Endocrinol* 2006; 189: 67-75.
19. Tschöp M, Smiley DL, Heiman ML. Ghrelin induces adiposity in rodents. *Nature* 2000; 407: 908-913.
20. Hosoda H, Kojima M, Matsuo H, Kangawa K. Ghrelin and des-acyl ghrelin: two major forms of rat ghrelin peptide in gastrointestinal tissue. *Biochemical Biophys Res Commun* 2000; 279: 909-913.
21. Nakai Y, Hosoda H, Nin K, Ooya C, Hayashi H, Akamizu T, Kangawa K. Plasma levels of active form of ghrelin during oral glucose tolerance test in patients with anorexia nervosa. *Eur J Endocrinol* 2003; 149: R1-3.
22. Yilmaz Z, Ilcol Y, Golcu E. Serum leptin and ghrelin levels in response to methylprednisolone injection in healthy dogs. *Res Vet Sci* 2007; 82: 187-194.
23. Kadokawa H, Blache D, Martin G. Plasma leptin concentrations correlate with luteinizing hormone secretion in early postpartum Holstein cows. *J Dairy Sci* 2006; 89: 3020-3027.
24. Comba A, Mert H, Coma B. Leptin levels and lipids profile determination in different sheep breeds. *Pak Vet J* 2016; 36: 169-173.
25. Delavaud C, Ferlay A, Faulconnier Y, Bocquier F, Kann G, Chilliard Y. Plasma leptin concentration in adult cattle: Effects of breed, adiposity, feeding level, and meal intake. *J Anim Sci* 2002; 80: 1317-1328.
26. Zhang Y, Proenca R, Maffei M, Marisa B, Lori L, Friedman J. Positional cloning of the mouse obese gene and its human homologue. *Nature* 1994; 372: 425-431.
27. Ahima J, Flier R. Leptin. *Annu Rev Physiol* 2000; 62: 413-437.
28. Considine R, Caro J. Leptin and the regulation of body weight. *J Biochem Cell Biol* 1997; 29: 1255-1272.
29. Kadokawa H, Blache D, Yamada Y, Martin G. Relationships between changes in plasma concentrations of leptin before and after parturition and the timing of first post-partum ovulation in high-producing Holstein dairy cows. *Reprod Fertil Dev* 2000; 12: 405-411.

30. Temizel EM, Batmaz H, Keskin A, Orman A, Gencoglu H, Catik S, Topal O. Butaphosphan and cyanocobalamin treatment of pregnant ewes: metabolic effects and potential prophylactic effect for pregnancy toxemia. *Small Rumin Res* 2015; 125: 163-172.
31. Caldeira RM, Portugal AV. Interrelationships between body condition and metabolic status in ewes. *Small Rumin Res* 1991; 6: 15-24.
32. Kurose Y, Iqbal J, Rao A, Murata Y, Hasegawa Y, Terashima Y, Kojima M, Kangawa K, Clarke IJ. Changes in expression of the genes for the leptin receptor and the growth hormone-releasing peptide/ghrelin receptor in the hypothalamic arcuate nucleus with long-term manipulation of adiposity by dietary means. *J Neuroendocrinol* 2005; 17: 331-340.
33. Nakahara K, Nakagawa M, Baba Y, Sato M, Toshinai K, Date Y, Nakazato M, Kojima M, Miyazato M, Kaiya H et al. Maternal ghrelin plays an important role in rat fetal development during pregnancy. *Endocrinol* 2006; 147: 1333-1342.
34. Maillard V, Ram C, Froment P. Ghrelin in female and male reproduction. *Int J Pept* 2010; 2010: 2016.
35. Fuglsang J, Skjærbæk C, Espelund U, Frystyk J, Fisker S, Flyvbjerg A. Ghrelin and its relationship to growth hormones during normal pregnancy. *Clin Endocrinol* 2005; 554-559.
36. Di Palo R, Campanile G, Prandi A, Baruselli P, Vecchio D, Carvalho N, Zicarelli L. Plasma leptin levels in Murrah buffalo heifers fed diet with two different energy levels. *Ital J Anim Sci* 2005; 4: 301-303.
37. Koh KK, Park SM, Quon MJ. Contemporary reviews in cardiovascular medicine leptin and cardiovascular disease. *Circulation* 2013; 11: 3238-3249.
38. Halle M, Berg A, Garwers U, Grathwohl D, Knisel W, Keul J. Concurrent reductions of serum leptin and lipids during weight loss in obese men with type II diabetes. *Am J Physiol Endocrinol Metab* 1999; 277: 277-282.
39. Ban-Tokuda T, Orden EA, Barrio AN, Lapitan RM, Delavaud C. Effects of species and sex on plasma hormone and metabolite concentrations in crossbred Brahman cattle and crossbred water buffalo. *Livestock Science* 2007; 107: 244-252.
40. Cohen SM, Werrmann JG, Tota MR. ¹³C NMR study of the effects of leptin treatment on kinetics of hepatic intermediary metabolism. *Proceedings of the National Academy of Sciences of the United States of America* 1998; 95: 7385-7390.
41. Prieur X, Tung Y, Griffin J, Farooqi I, O'Rahilly S, Coll A. Leptin regulates peripheral lipid metabolism primarily through central effects on food intake. *Endocrinology* 2008; 149: 5432-9.
42. Berridge M. Cell Signalling Through Phospholipid Metabolism. *J Cell Sci* 1986; 4: 137-153.
43. McMurray W, Magee W. Phospholipid metabolism. *Annu Rev Biochem* 1972; 41: 129-160.
44. Nielsen M, Jakobsen K. Changes in mammary uptake of free fatty acids, triglyceride, cholesterol and phospholipid in relation to milk synthesis during lactation in goats. *Comprehensive Biochemical Physiology* 1994; 109: 857-867.
45. Bitman J, Wood DL. Changes in milk fat phospholipids during lactation. *J Dairy Sci* 1989; 73: 1208-1216.
46. Tainturier D, Braun J, Rico A, Thouvenot J. Variations in blood composition in dairy cows during pregnancy and after calving. *Res Vet Sci* 1984; 37: 129-131.
47. Krajnicakova M, Bekeova E, Hendrichovscky V, Maracek I. Concentrations of total lipid, cholesterol and progesterone during oestrus synchronization and pregnancy in sheep. *Vet Med* 1993; 38: 349-357.
48. Piccione G, Caola G, Giannetto C, Grasso F, Runzo SC, Zumbo A, Pennisi P. Selected biochemical serum parameters in ewes during pregnancy, post-parturition lactation and dry period. *Anim Sci Pap Rap* 2009; 27: 321-330.